

MATERIA MEDICA

A MANUAL
FOR STUDENTS

ISAMBARD OWEN


M15554

THE LIBRARY
OF THE
MANCHESTER



MEDICAL SOCIETY

Geo. Folkner & Sons, Manchester



22101624809

7175.
Jup
112
046
d

Presented by

Dr. E. Bosdin Leech.

Manchester.

December 30th., 1926.



Digitized by the Internet Archive
in 2014

<https://archive.org/details/b20399728>

34800
John Adolph Albert.

MATERIA MEDICA.

A MANUAL

FOR THE USE OF STUDENTS.

BY

ISAMBARD OWEN, M.D.,

LECTURER ON MATERIA MEDICA AND THERAPEUTICS TO ST. GEORGE'S HOSPITAL;

ASSISTANT-PHYSICIAN TO THE BROMPTON HOSPITAL FOR CONSUMPTION

AND DISEASES OF THE CHEST.

LONDON:

J. & A. CHURCHILL, 11, NEW BURLINGTON STREET.

1883.

15843

26317887

PRINTED BY WHITING AND CO., LIMITED, 30 & 32, SARDINIA STREET,
LINCOLN'S-INN-FIELDS, W.C.

MISS 54

WELLCOME INSTITUTE	
Y	
Coll.	weim Omec
Call	
No.	QVX
	1883
	097m

PREFACE.

THIS little book has no pretensions to be considered an original or complete treatise on *Materia Medica*. It is simply intended to aid Students, by placing the outlines of the subject before them in such a manner that they may be clearly grasped, and readily committed to memory. It is to be regarded rather as a supplement to, than as a substitute for, the systematic works.

The contents of the former "*Tables of Materia Medica*", of which five editions were called for in eight years, are incorporated, in a revised form, in the present Manual.

I have great pleasure in thanking Dr. Frederick J. Hicks, of the Brompton Hospital, for his material assistance in revising the sheets of Section III; and Mr. A. F. Damon, of St. George's Hospital, for much valuable information on Pharmaceutical points.

41, GLOUCESTER GARDENS, W.

May 1st, 1883.



TABLE OF CONTENTS.

	PAGE
THE PHARMACOPŒIA	1
The Drugs	1
The Preparations	3
The Weights and Measures	11
 SECTION I.—DRUGS FURNISHED DIRECTLY BY THE ANIMAL KINGDOM ...	12
First Group	12
Second Group	13
Third Group	14
 SECTION II.—DRUGS FURNISHED DIRECTLY BY THE VEGETABLE KINGDOM	16
First Group :—Fungi and Lichens	17
Second „ Plants entire	18
Third „ Green Tops. Twigs	18
Fourth „ Flowering Tops. Strobile	19
Fifth „ Flowers	20
Sixth „ Buds	21
Seventh „ Woods	22
Eighth „ Barks	23
Ninth „ Leaves	25
Tenth „ Roots and Rhizomes	28
Eleventh „ Fruits. Glands from Fruit	34
Twelfth „ Seeds. Gossypium	39
Thirteenth „ Fixed Oils	42
Fourteenth „ Volatile Oils	44
Fifteenth „ Resins	48
Sixteenth „ Oleo-Resins	50
Seventeenth „ Balsams	51
Eighteenth „ Gums	52
Nineteenth „ Gum-Resins	53

Twentieth Group :—Concrete Juices. Galls	55
Twenty-first „ Active Principles of Definite Chemical Com- position—				
α. Carbohydrates	60
β. Acids	61
γ. Neutral Principles	61
δ. Alkaloids	62
Extraction of Alkaloids	65
Botanical Synopsis of the Natural Orders and Plants furnishing Official Drugs	72
Non-Official Drugs from the Vegetable Kingdom—				
Plants or parts of Plants	84
Exudations and Concrete Juices	86
Oleo-Resin	86
Fixed Oil	87
Volatile Oils	87
Active Principles of Definite Composition	88
SECTION III.—MINERALS AND PRODUCTS OF MANUFACTURE	91
First Group :—Non-metallic Elements and their Compounds	92
Second „ Mineral Acids	96
Third „ Alcohol and its Derivatives. Amylic Alcohol and its Derivatives	99
Fourth „ Tars and their Derivatives. Organic Acids	105
Fifth „ Pyroxylin, Glycerine ; their Derivatives, and Mis- cellanea	111
Sixth „ Metals and Metallic Compounds	113
Potassium Compounds	114
Sodium Compounds	118
Ammonium Compounds	123
Iron Compounds	125
Mercurial Compounds	129
Calcium Compounds	132
Zinc Compounds	134
Lead Compounds	136
Bismuth Compounds	138
Antimonial Compounds...	139
Arsenical Compounds	141

Sixth Group—*continued*.

Magnesium Compounds...	142
Alum Compounds	143
Lithium Compounds	144
Silver Compounds	145
Copper, Cadmium, and Cerium Compounds...			146
Official Preparations	147

SYNOPSIS OF THE COMPOUND PREPARATIONS OF THE PHARMACOPŒIA, AND

THEIR INGREDIENTS	152
-------------------	-----	-----	-----	-----

POPULAR DESIGNATIONS FOR PHARMACOPEIAL PREPARATIONS	...	162
---	-----	-----

PHRASES USED IN PRESCRIBING	164
-----------------------------	-----	-----	-----

SYNOPSIS OF DOSES	165
-------------------	-----	-----	-----

PROPORTION OF ACTIVE INGREDIENTS IN THE PHARMACOPEIAL PREPARATIONS	174
--	-----

INCOMPATIBILITY	181
-----------------	-----	-----	-----

INDEX	183
-------	-----	-----	-----

Twentieth Group :—Concrete Juices. Galls	55
Twenty-first „ Active Principles of Definite Chemical Com- position—				
<i>a.</i> Carbohydrates	60
<i>β.</i> Acids	61
<i>γ.</i> Neutral Principles	61
<i>δ.</i> Alkaloids	62
Extraction of Alkaloids	65
Botanical Synopsis of the Natural Orders and Plants furnishing				
Official Drugs	72
Non-Official Drugs from the Vegetable Kingdom—				
Plants or parts of Plants	84
Exudations and Concrete Juices	86
Oleo-Resin	86
Fixed Oil	87
Volatile Oils	87
Active Principles of Definite Composition	88
SECTION III.—MINERALS AND PRODUCTS OF MANUFACTURE	91
First Group :—Non-metallic Elements and their Compounds	92
Second „ Mineral Acids	96
Third „ Alcohol and its Derivatives. Amylic Alcohol and its Derivatives	99
Fourth „ Tars and their Derivatives. Organic Acids	105
Fifth „ Pyroxilin, Glycerine ; their Derivatives, and Mis- cellanea	111
Sixth „ Metals and Metallic Compounds	113
Potassium Compounds	114
Sodium Compounds	118
Ammonium Compounds	123
Iron Compounds	125
Mercurial Compounds	129
Calcium Compounds	132
Zinc Compounds	134
Lead Compounds	136
Bismuth Compounds	138
Antimonial Compounds...	139
Arsenical Compounds	141

Sixth Group—*continued*.

Magnesium Compounds...	142
Alum Compounds	143
Lithium Compounds	144
Silver Compounds	145
Copper, Cadmium, and Cerium Compounds...			146
Officinal Preparations	147

SYNOPSIS OF THE COMPOUND PREPARATIONS OF THE PHARMACOPŒIA, AND
THEIR INGREDIENTS

...	152
-----	-----	-----	-----	-----

POPULAR DESIGNATIONS FOR PHARMACOPŒIAL PREPARATIONS

...	162
-----	-----

PHRASES USED IN PRESCRIBING

...	164
-----	-----	-----	-----	-----

SYNOPSIS OF DOSES

...	165
-----	-----	-----	-----	-----

PROPORTION OF ACTIVE INGREDIENTS IN THE PHARMACOPŒIAL PREPARATIONS

174

INCOMPATIBILITY

...	181
-----	-----	-----	-----	-----

INDEX

...	183
-----	-----	-----	-----	-----	-----

ERRATA.

Page 7, eighteenth line from the bottom :—*for* $\frac{1}{10}$ *read* $\frac{1}{16}$.

Page 127, fourteenth line :—omit *

MATERIA MEDICA.

THE PHARMACOPŒIA.

THE BRITISH PHARMACOPŒIA of 1867, with the additions of 1874, defines 393 drugs, and prescribes 530 preparations, under Latin and English names.*

It prescribes the mode of production of a large number of the manufactured articles included among the drugs.

It specifies the physical properties and the chemical tests by which the drugs may be recognised and their purity certified.

It lays down tables of weights and measures.

It authorises a scale of doses for all the drugs and preparations intended for internal administration.

THE DRUGS.

Of the 393 drugs, 19 are furnished directly by the animal kingdom, and 197 by the vegetable kingdom, of nature. The remaining 177 are native minerals and products of industrial or chemical manufacture.

The following twenty-nine of the drugs are not themselves used in medicine, but are defined in the Pharmacopœia as being the source of other officinal drugs, or as required for some of the processes prescribed :—

(Minerals and Manufactured Products.)

Natural Water	or	<i>Aqua.</i>
Chalk (<i>unpurified</i>)	„	<i>Creta.</i>
Marble (source of Carbonic Acid)	„	<i>Marmor Album.</i>
Iron	„	<i>Ferrum.</i>
Copper	„	<i>Cuprum.</i>
Bismuth	„	<i>Bismuthum.</i>
Purified Bismuth	„	<i>Bism. Purificatum.</i>
Zinc	„	<i>Zincum.</i>
Granulated Zinc	„	<i>Zincum Granulatum.</i>
Purified Silver	„	<i>Argentum Purificatum.</i>
Black Oxide of Manganese	„	<i>Manganesi Oxidum Nigrum.</i>
Black Antimony	„	<i>Antimonium Nigrum.</i>
Nitrate of Ammonia	„	<i>Ammonia Nitras.</i>
Nitrate of Soda	„	<i>Sodæ Nitras.</i>
Nitrate of Lead	„	<i>Plumbi Nitras.</i>

* Drugs and preparations contained in the Pharmacopœia are termed "Officinal" or "Official". Throughout this book, the use of **Clarendon type**, or of the Latin language, in the absence of any statement to the contrary, signifies the officinal character.

Sulphate of Mercury	or	<i>Mercurii Sulphas.</i>
Acetate of Soda	„	<i>Sodæ Acetatus.</i>
Bichromate of Potash	„	<i>Potassæ Bichromas.</i>
Ferrocyanide of Potassium	„	<i>Potassæ Prussias Flava.</i>
Bone-Black*	„	<i>Carbo Animalis.</i>
Bone-Ash	„	<i>Os Ustum.</i>
Amylic Alcohol (<i>Fusel Oil</i>)	„	<i>Alcohol Amylicum.</i>

(Vegetables.)

Bebeeru Bark (source of Sulphate of Beberia)	or	<i>Neetandra Cortex.</i>
Cevadilla (source of Veratria)	„	<i>Subadilla.</i>
Podophyllum Root (source of Podophyllin)	„	<i>Podophylli Radix.</i>
Scammony Root (source of Scammony Resin)	„	<i>Scammonia Radix.</i>
Squirting Cucumber (source of Elaterium)	„	<i>Ecbalii Fructus.</i>
“Worm Seed” (source of Santonin)	„	<i>Santonica.</i>

(Animal Products.)

Unclarified Honey	or	<i>Mel.</i>
---------------------------	----	-------------

The following thirty-one are used merely as vehicles :—

(Animal Kingdom.)

Milk	or	<i>Lae.</i>
Clarified Honey	„	<i>Mel Depuratum.</i>
Sugar of Milk	„	<i>Saccharum Lactis.</i>
Lard	„	<i>Adcps Preparatus.</i>
Clarified Mutton Suct	„	<i>Sevum Preparatum.</i>
Bees-wax	„	<i>Cera Flava.</i>
White-wax	„	<i>Cera Alba.</i>

(Vegetable Kingdom.)

Wheat-flour	or	<i>Farina Triticæ.</i>
Linseed-meal	„	<i>Linæi Farina.</i>
Starch	„	<i>Amylum.</i>
Loaf-sugar	„	<i>Saccharum Purificatum.</i>
Gum Arabic	„	<i>Acaciæ Gummi.</i>
Tragacanth	„	<i>Tragacantha.</i>
Almond Oil	„	<i>Oleum Amygdalæ.</i>
Olive Oil	„	<i>Oleum Olivæ.</i>
Hips	„	<i>Rosæ Caninæ Fructus.</i>
Cocoa-butter	„	<i>Oleum Theobromæ.</i>
Canada Balsam	„	<i>Terebinthina Canadensis.</i>
Gutta-percha	„	<i>Gutta-percha.</i>

(Minerals and Manufactured Products.)

Distilled Water	or	<i>Aqua Destillata.</i>
Proof Spirit	„	<i>Spiritus Tenuior.</i>
Rectified Spirit	„	<i>Spiritus Rectificatus.</i>
Sherry	„	<i>Vinum Xericum.</i>
Orange Wine	„	<i>Vinum Aurantii.</i>
Crude Ether†	„	<i>Æther.</i>
Glycerine	„	<i>Glycerinum.</i>
Acetic Acid‡	„	<i>Acidum Aceticum.</i>
Treacle	„	<i>Theriacæ.</i>
Bread-crumbs	„	<i>Mica Panis.</i>
Soft Soap	„	<i>Sapo Mollis.</i>
Curd Soap	„	<i>Sapo Animalis.</i>

Rosin, and Castile Soap (*Resina*, and *Sapo Durus*) are used mainly, but not exclusively, as vehicles.

* Animal Charcoal (*Carb. An. Purificatus*) is used, but mainly as a pharmaceutical (filtering) agent.

† Washed Ether (*Æther Purus*) is used medicinally.

‡ Glacial Acetic Acid, and Dilute Acetic Acid, are used medicinally.

The following four are colouring agents :—

Cochineal	or	<i>Coccus.</i>
Red Sandal-wood	„	<i>Pterocarpī Lignum.</i>
Red Poppy	„	<i>Papaver Rhæas.</i>
Saffron	„	<i>Crocus.</i>

Red-rose petals (*Rosa Gallica*) are used chiefly as a colouring agent, but possess astringent properties also.

The following eight are merely flavouring agents :—

Orange-flower Water	or	<i>Aqua Aurantii Floris.</i>
Cabbage-Rose	„	<i>Rosa Centifolia.</i>
Elder flowers	„	<i>Sambuci Flores.</i>
Raisins	„	<i>Uvæ.</i>
Mulberry-juice	„	<i>Mori Succus.</i>
Jordan Almond	„	<i>Amygdala Dulcis.</i>
Liquorice Root	„	<i>Glycyrrhizæ Radix.</i>
Hemidesmus Root	„	<i>Hemidesmi Radix.</i>

The following twenty-four may also be classed as flavouring agents, though the first possesses bitter, the rest stimulant and carminative properties, which make them therapeutically useful as vehicles :—

Bitter Orange Peel, fresh	or	<i>Aurantii Fructus.</i>
„ „ „ dried	„	<i>Aurantii Cortex.</i>
Lemon Peel	„	<i>Limonis Cortex.</i>
Caraway	„	<i>Carui Fructus.</i>
Cardamoms	„	<i>Cardamomum.</i>
Cinnamon	„	<i>Cinnamomi Cortex.</i>
Coriander	„	<i>Coriandri Fructus.</i>
Cloves	„	<i>Caryophyllum.</i>
Dill	„	<i>Anethi Fructus.</i>
Nutmeg	„	<i>Myristica.</i>
Pimento	„	<i>Pimenta.</i>

And the Essential Oils of

Anise, Caraway,	or	<i>Ol. Anisi, Carui,</i>
Cinnamon, Cloves,	„	<i>Cinnamomi, Caryophylli,</i>
Coriander, Dill,	„	<i>Coriandri, Anethi,</i>
Lavender, Lemon,	„	<i>Lavandulæ, Limonis,</i>
Nutmeg, Peppermint,	„	<i>Myristicæ, Menthæ Piperitæ,</i>
Pimento, Rosemary, and	„	<i>Pimentæ, Rosmarini,</i>
Spearmint	„	<i>Menthæ Viridis.</i>

The remaining 297 drugs are reputed to possess more or less active medicinal properties.

THE PREPARATIONS.

The preparations prescribed by the Pharmacopœia are of thirty-two different kinds.

Extracts (*Extracta*) consist either (α , β) of the expressed juices of fresh vegetables, clarified and concentrated; or (γ , δ) of the soluble ingredients of dry drugs, separated by macerating them in various menstrua, and evaporating the strained solution to the desired consistence.

a. The Extracts of Aconite, Belladonna, Hemlock, Henbane, and Lettuce are prepared from fresh plants as follows :—

1. The fresh juice, obtained by bruising in a mortar,
Heated to 130 deg. F.
A coagulum of green colouring matter separates, is strained off, and put aside.
2. The strained juice,
Heated to 200 deg. F.
A coagulum of albumen separates.

3. The juice filtered from albumen,
Evaporated to a thin syrup;
The green colouring matter returned;
Evaporated at 140 deg. F. to a consistence "suitable for forming pills".

These are termed "Green Extracts".

β. The Extracts of Colchicum and Taraxacum from fresh drugs as follows:—

1. The fresh juice, obtained by crushing and expressing,
Decanted from deposit;
Heated to 212 deg. F.
A coagulum of albumen separates.
2. The filtered liquid,
Evaporated at 160 deg. F. to a similar consistence.

The Acetic Extract of Colchicum is prepared in a similar way, but Acetic Acid is added to the crushed corms before expressing the juice, and evaporation is not carried so far.

γ. In making the extracts of dry drugs, the menstrua used are the following:—

Cold water, for the Extracts of Opium, Calumba, Quassia, Rhatany, and Liquorice.

Also for the Liquid Extracts of Yellow-Bark, Opium, Bael, Liquorice, and Ergot.

Water at 160 deg. F., for the Liquid Extract of Sarsaparilla.

Boiling water, for the Extracts of Chamomile, Aloes, Pareira, Poppy, Logwood, and Gentian.

Also for the Liquid Extract of Pareira.

Diluted Spirit, for the Extract of Rhubarb.

Proof Spirit, for the Extracts of Stramony and Colocynth.

Rectified Spirit, for the Extracts of Nux Vomica, Calabar Bean, and Indian Hemp.

Rectified Spirit first, and cold water afterwards, for the Extract of Jalap.
Rectified Spirit first, and boiling water afterwards, for the Extract of Hop.

Ether, for the Liquid Extract of Fern.

Ext. Mezerei Æthereum is an Ethereal Extract of a *Spirituos Extract of Mezcreon*.

Stramony and Ergot are cleared of their oils by ether before making the extracts.

In making the Extract of Poppy, the gummy matters in the solution are precipitated by Rectified Spirit before evaporation.

Extract of Chamomile has the essential oil of Chamomile added to it.

The Extract of Colocynth is a "Compound Extract"; having other drugs added to it during evaporation.

- δ. The *Liquid Extracts (Extracta Liquida)* are evaporated only so far as to produce a highly concentrated solution; except in the case of the Liquid Extract of Fern, the fluid form of which is given by the oily matters of the drug itself, the ether being all distilled off. To the rest a little spirit is added to prevent decomposition.

Ext. Opii, Liq. is simply an infusion of the Extract, with a little spirit.

Extracts of Nux Vomica, Calabar-bean, Mezereon, and Indian Hemp are left in a softer state than the others, which are evaporated to a consistence "suitable for forming pills"; except the Extracts of Rhatany, Logwood, and Aloes, which are evaporated to dryness.

Juices (*Succi*)* are the expressed juices of fresh vegetables preserved by the addition of one-third of their volume of Rectified Spirit, and strained after standing for a week.

The Pharmacopœia prescribes five, of
Belladonna, Hemlock, Henbane, Dandelion, and Broom.

Essences (*Essentiæ*) are solutions of Essential (volatile) Oils in four times their volume of Rectified Spirit.

The Pharmacopœia prescribes two only, of
Anise and Peppermint.

Infusions (*Infusa*) are extractive solutions of such ingredients of vegetables as are soluble in water, made by pouring boiling water on the broken-up drug, macerating for periods varying from a quarter of an hour to four hours, and straining clear. The Pharmacopœia prescribes twenty-eight.

The infusions of Cusparia and Chiretta are made with water at 120 deg. F. only.

The infusions of Quassia and Calumba with cold water, so as not to dissolve out the starch that these drugs contain.

Infusion of Koussou is administered unstrained.

Acid Infusion of Roses is made with boiling water acidulated with Sulphuric Acid.

Decoctions (*Decocta*) are similar preparations, made by boiling the drugs in water for periods varying from five to twenty minutes, and straining clear. Some are ordered to be strained *while hot*, as part of their active ingredients is deposited on cooling.

The Pharmacopœia prescribes fourteen.

In preparing Decoction of Pomegranate Root, the liquid is boiled down to half its original bulk.

Tinctures (*Tincturæ*).

The Tinctures of Quinine, Iodine, Iron Salts, and Chloroform are simple solutions in spirit. The Tincture of Indian Hemp is a solution of the extract in spirit.

The rest of the Pharmacopœial Tinctures are extractive solutions prepared by macerating animal or vegetable drugs in spirit, in some cases for two days, in others for seven, and straining or percolating.

It is to these extractive Tinctures, which are generally deeply tinged by the colouring matters of the drug, that the name of *Tincture* properly applies.

The Pharmacopœia prescribes sixty eight tinctures altogether.

Proof Spirit is the menstruum used for most of the Tinctures.

The stronger Rectified Spirit is used in preparing the Tinctures of

Aconite, Arnica, Assafœtida,	Indian Hemp,
Benzoin, Cubebs,	Lavender, Larch,
Capsicum, Castor,	Myrrh, Nux Vomica,
Hellebore, Kino,	Pellitory, Tolu,
Iodine, Iron Salts,	Fresh Orange Peel, and Ginger.

A mixture of Rectified Spirit and Tr. Cardam. Co. for Tr. Chloroformi Co.

Tincture of Orange Peel for Tr. Quiniæ.

Spirit of Ether for Tr. Lobeliæ Ætherca.

* Mulberry, Lemon, and Buckthorn Juices (*Mori Succus*, *Jamoni Succus*, *Rhamni Succus*) are natural drugs, and not preparations. The prepared juices are termed, "*Succus Belladonnae*, *Succus Conii*", etc.

Sal. Volatile for Tr. Guaiaci Ammoniata.

„ „ Valerianæ „

A mixture of Rectified Spirit with one-fourth of Liq. Amm. Fortior for Tr. Opii Ammoniata.

Proof Spirit for Tr. Quiniæ Ammoniata, Liq. Ammoniaë being added subsequently.

Wines (*Vina*) are similar preparations made with white wines. The Pharmacopœia prescribes nine,

Sherry (*Vin. Xericum*) is prescribed for all but Vinum Quiniæ and Vinum Ferri Citratis, which are made with Orange Wine (*Vin. Aurantii*).

Spirits (*Spiritus*) are solutions of colourless substances in Rectified Spirit. The Pharmacopœial *Spirits* (thirteen in number) are solutions of volatile oils (two being ammoniated), of Camphor, Chloroform, Ether, and Nitrous Ether.

Sp. Armoraciæ Co., Sp. Amm. Aromaticus, and Sp. Amm. Fœtidus are prepared by distillation.

The preparation of Sp. Ætheris Nitrosi is detailed in Section iii.

The rest are prepared by direct solution, in the following proportions :—
Oils, 1 in 50; Chloroform, 1 in 20; Camphor, 1 in 10; Ether, 1 in 3.

Spiritus Rectificatus (Rectified Spirit), Sp. Tenuior (Proof Spirit), and Sp. Vini Gallici (French Brandy) are not *preparations*. (See “Alcohol”.)

Vinegars (*Acta*) are extractive solutions in Acetic Acid, prepared by maceration and straining.

The Pharmacopœia prescribes two only.

In making Acetum Cantharidis, a mixture of the glacial with the ordinary acid is used.

For Acetum Scillæ, the Dilute Acid; a little spirit being added after straining, and the whole filtered again.

Solutions (*Liquores*) are properly watery solutions of definite chemical compounds. The menstruum of the Pharmacopœial *Solutions* is not in all cases pure water. (See under “Alkaloids”, “Iodine”, and “Metallic Compounds”.) The Pharmacopœia prescribes forty solutions, including the three following, and the effervescing solutions.

Liquor Epispasticus is an extractive solution prepared from Cantharides with Acetic Acid and Ether.

Liquor Gutta-Percha is a solution of Gutta-Percha in Chloroform.

Collodion is a solution of Gun-Cotton (*Pyroxylin*) in Ether and Rectified Spirit.

The “Effervescing Solutions” (*L. Effervescentes*) are weak solutions of the Bicarbonates of Potash and Soda, and of Carbonate of Lithia, charged with Carbonic Acid under a pressure of seven atmospheres, and corked down,—in short, the “Potash”, “Soda”, and “Lithia” waters of ordinary use. Liq. Magnesiæ Carbonatis and Liq. Mag. Citratis also contain Carbonic Acid, developed under pressure, and require to be kept corked.

Waters (*Aquæ*) consist of water saturated with essential oils or similar fragrant substances sparingly soluble. The Pharmacopœia prescribes thirteen.

Essential Oils are extracted by distilling the vegetable products containing them with water. From the milky fluid which condenses in the receiver the oil separates, leaving a clear “*distilled water*”, which

retains the minute portion of the oil it is capable of dissolving. The Pharmacopœia prescribes processes for distilling these *waters* from vegetable drugs or essential oils, but they are practically procured as above.

Aqua Camphoræ and Aqua Chloroformi are prepared simply by shaking the two drugs up with water.

Syrups (*Syrupi*). The Pharmacopœial syrups are prepared by adding large proportions of sugar or syrup to

the fresh juices of *Lemon*, *Mulberry*, and *Buckthorn* fruit;

Solutions of *Chloral*, of *Phosphate*, and of *Iodide*, of *Iron*;

Specially made infusions of *Hemidesmus*, *Poppy*, *Red Poppy*, *Red Rose*, and *Senna*;

A special decoction of *Tolu*;

A special tincture of *Rhubarb*;

The official Tinctures of *Orange Peel* and *Ginger*;

„ „ Vinegar of *Squill*;

„ „ *Orange Flower* Water.

To some a little spirit is added for preservation.

Simple Syrup (*Syrupus*) is prepared by dissolving sugar in half its weight of water, the product giving 6 parts of sugar in 7.

Mucilages (*Mucilagines*) are solutions of colloid substances in water.

The Pharmacopœia prescribes three, of

Gum-arabic, Starch, and Tragacanth.

Glycerines (*Glycerina*) are solutions in glycerine.

The Pharmacopœia prescribes four such solutions, of

Borax, of Tannic, Gallic, and Carbolic Acids,

each prepared by dissolving 1 part in 4, making $4\frac{1}{2}$.

Also a translucent jelly, *Glycerinum Amyli*, for use as an ointment, prepared by heating 1 part of starch with 8 of glycerine.

Oils (*Olea*) are solutions in fixed oils. The Pharmacopœia prescribes one only, *Oleum Phosphoratum*;—Almond Oil saturated with Phosphorus, containing $\frac{1}{16}$ grain in 10 minims.

Honey (*Mel*).

Mel Boracis is a solution of Borax in 7 parts of Clarified Honey. Oxymel is a mixture of Clarified Honey with an eighth part of Acetic Acid and the same of water.

Oxymel Scillæ is a mixture of Clarified Honey with Vinegar of Squill, evaporated to a thick consistency.

Dilute Acids (*Acida Diluta*) are Acids diluted with water to a strength convenient for dispensing. The Pharmacopœia prescribes

Ac. Sulphuricum Dil., containing $8\frac{1}{2}$ p.c. of the "Acidum".

„ Aceticum Dil., „ $12\frac{1}{2}$ p.c. „ „

„ Nitricum Dil., „ 20 p.c. „ „

„ Hydrochloricum Dil., „ 30 p.c. „ „

„ Phosphoricum Dil., „ 13.8 p.c. of Phosphoric Acid, $H_3 PO_4$.

„ Hydrocyanicum Dil., „ 2 p.c. of Prussic Acid HCN.

(The last two are prescribed in the Pharmacopœia in the diluted form only.)

Also:—Ac. Sulphuricum Aromaticum,—7 p.c. of the "Acidum" in Rectified Spirit, flavoured with Cinnamon and Ginger.

* * The above preparations, it will be noted, simply place the drugs in a convenient form for dispensing. They are for the most part simple, containing but a single drug, or its active part, besides the vehicle. Some are directed to be flavoured and coloured with various ingredients; and a few are compound preparations (see the "Synopsis" later in the book), *i.e.*, they contain the active principles of more than one drug.

* * The following are forms of preparation ready for internal or external administration without further dispensing. Many are compound (see the "Synopsis").

(For internal use.)

Compound Powders (*Pulveres Compositæ*) are intimate admixtures of dry drugs, finely powdered.

Confections or electuaries (*Confectiones*) are similar mixtures of dry or pulpy drugs, brought to the consistence of a soft extract with sugar, syrup, or honey.

Pill-masses (*Pilulæ*) are composed of extracts or other tenacious substances, or are directed to be made up with Castile soap, treacle, syrup, confection of roses, or of hips, beeswax, or castor oil.

Some are scented with particular essential oils, as a means of distinguishing them apart.

The Pharmacopœia gives no directions for rolling these masses into pills. As usually rolled, each pill is made by 5 grains of the pill-mass. The more active pills, *Pil. Hydrargyri*, *Pil. Phosphori*, *Pil. Plumbi c. Opio*, *Pil. Saponis Co.*, are rolled in various sizes.

Lozenges (*Trochisci*). The Pharmacopœial lozenges are small tablets, made up with sugar and gum. They are ten in number; of

Bismuth.	Catechu.
Reduced Iron.	Opium.
Chlorate of Potash.	Morphia.
Bicarbonate of Soda.	Morphia and Ipecacuanha.
Tannic Acid.	Ipecacuanha.

Mixtures* (*Misturæ*). Under this title, the Pharmacopœia prescribes modes of suspending or emulsifying certain drugs insoluble in water; also a preparation of Epsom Salts and Senna, and two aromatic bitter infusions, one of which is made to take up a minute quantity of iron. (See the "Synopsis of Compound Preparations".)

Inhalations (*Vapores*). The Pharmacopœia prescribes methods of disengaging for inhalation.

Prussic Acid,—by mixing 10 to 15 min. of *Ac. Hydrocyan. Dil.*, with a drachm of water;

Chlorine.—by moistening 2 oz. of Chlorinated Lime with cold water.

And of volatilizing for the same purpose

Creasote, by mixing 12 min. with 8 oz. of boiling water in an "inhaler" which allows air to be drawn through the fluid;

Iodine, by mixing a drachm of *Tr. Iodi* with an ounce of water, and heating gently;

Conia, by mixing a drachm of Extract of Hemlock with a fluid drachm of *Liq. Potassæ* and ten of water, and placing 20 min. of the mixture on a sponge in an "inhaler" which allows the vapour from hot water to be drawn over it.

* Draughts (*Haustus*) are single doses of mixtures.

Enemas or Clysters (*Enemata*). The Pharmacopœia prescribes methods of preparing for administration *per rectum*

Opium, by mixing 30 min. of Tr. Opii with 2 oz. of Mucilage of Starch,—*Enema Opii*;

Tobacco, by making an infusion of 20 grains in 8 oz. of boiling water,—*Enema Tabaci*.

And the following purgative Clysters :—

Enema Assofetidæ.—30 grains rubbed into an emulsion with 4 oz. of water.

Enema Aloës.—40 grains rubbed together with half a pint of Mucilage of Starch and 15 grains of Potassic Carbonate.

Enema Terebinthinæ.—An ounce of Ol. Terebinth. mixed with three-fourths of a pint of Mucilage of Starch.

Enema Magnesie Sulphatis.—An ounce of Epsom Salts dissolved in three-fourths of a pint of Mucilage of Starch, and an ounce of Olive Oil added.

Suppositories (*Suppositoria*) are small cones of greasy matters with an intimate admixture of active drugs, for administration *per rectum*. The Pharmacopœia prescribes four made with cocoa-butter, lard, and wax, viz. :—

Supp. Morphie,—Hydrargyri,—Acidi Tannici,—Plumbi Co. (Opium and Acetate of Lead) ;

And three made with Curd Soap and Starch, viz. :—

Supp. Ac. Carbolici cum Sapone,—Morphie c. S.,—Ac. Tannici c. S.

Hypodermic Injections (*Injectiones Hypodermicæ*) are strong solutions in plain water, without spirit or appreciable amounts of free acid, for injection into the areolar tissue under the skin. The Pharmacopœia prescribes but one, *Inject. Morphie Hypod.*, containing 5 gr. of the acetate in each drachm, or $\frac{1}{12}$ gr. in each minim.

(For external application.)

Lotions (*Lotiones*) are solutions or suspensions of drugs in water or spirit, for continuous external application. The Pharmacopœia prescribes but two, *Lotio Hydrargyri Nigra*, and *Lotio Hydr. Flava*, containing Mercurous and Mercuric Oxide respectively, suspended in water.

Liniments (*Linimenta*) are either strong tinctures, which can be painted on the skin and allowed to dry ; or mixtures containing oily matters, which enable them to be rubbed into a part. (See “Synopsis of Compound Preparations.”)

Ointments (*Unguenta*) are admixtures of active drugs with greasy matters, preserving a soft but solid consistence, for smearing over or rubbing into the skin. (See “Synopsis of Compound Preparations.”)

Unguentum Simplex is a mixture of white wax, lard and almond oil, without any active drug.

Poultices (*Cataplasmata*). The Pharmacopœia prescribes methods of making

Linseed Poultice (*Cat. Lini*) by mixing 4 oz. of Linseed Meal gradually with half a pint of boiling water, and stirring in half an ounce of olive oil ;

Hemlock Poultice (*Cat. Conii*), by mixing an ounce of powdered Hemlock leaf with 3 of Linseed Meal, and stirring gradually into half a pint of boiling water ;

Mustard Poultrice (*Cat. Sinapis*), by mixing $2\frac{1}{2}$ oz. of Linseed Meal with half a pint of boiling water, and stirring in $2\frac{1}{2}$ oz. of mustard ;

Chlorine Poultrice (*Cat. Sodæ Chloratæ*), by mixing 4 oz. of Linseed Meal with 8 oz. of boiling water, and adding 2 fl. oz. of Liq. Sod. Chloratæ ;

Charcoal Poultrice (*Cat. Carbonis*), by macerating 2 oz. of bread-crumbs for 10 minutes in half a pint of boiling water, stirring in $1\frac{1}{2}$ oz. of linseed meal, then $\frac{1}{2}$ oz. of wood charcoal, and sprinkling a second $\frac{1}{2}$ oz. of charcoal on the surface ;

Yeast Poultrice (*Cat. Fermentii*), by mixing 6 oz. of Yeast with the same amount of water at 100° F., stirring in $1\frac{1}{2}$ oz. of wheat flour, and setting the mass near a fire till it "rises".

Plasters (*Emplastra*). The Pharmacopœial plasters are all made with resins, wax, Castile soap, or lead-soap, which when gently warmed, soften and become adhesive to the skin. *Emplastrum Plumbi* consists entirely of lead-soap, and is used for adhesive purposes simply ; the others contain an admixture of more or less active drugs. (See "Synopsis of Compound Preparations.")

The Pharmacopœia gives no directions for *spreading* the plasters. When required for use, they are softened by heat, and evenly spread on thin leather or on any suitable tissue.

Papers (*Chartæ*). *Charta Epispastica* (blistering-paper) is made by coating cartridge paper with a Cantharides plaster, made with white wax, spermaceti, olive oil, and a little rosin and Canada Balsam.

Charta Sinapis (mustard leaf) is cartridge paper coated with mustard by the aid of gutta-percha solution. It requires to be immersed in warm water (see "*Sinapis*") for use.

* * * The nature of *Benzoated Lard*, of *Flexible Collodion*, and of the *distilled, purified, clarified, granulated, and dehydrated* drugs, is explained in the body of the work as each occurs.

The Pharmacopœia does not prescribe any of the following :—

Gargles (*Gargarismata*). Mixtures which it is customary to thicken with syrup, gum, or glycerine, for application to the fauces and pharynx.

Eye-washes (*Collyria*). Lotions which should be free from acid or caustic alkali, much spirit, or other irritating substances.

Linctuses (*Linctus*). Mixtures of syrupy, oily, or mucilaginous matters, sometimes medicated with active drugs, intended to be taken in small doses to allay irritation about the throat and larynx.

Medicated Pessaries (*Pessus*), preparations analogous to suppositories, for local application of drugs to the vaginal mucous membrane.

Urethral Suppositories, similar preparations made in the form of bougies for application to the urethra.

The following have been introduced by the leading druggists, but are not officinal :—

Chloroforms (*Chloroforma*). Tinctures made with Chloroform instead of spirit, intended for use in liniments.

Tablets. Single doses of alkaline or metallic salts, moulded by means of strong pressure into concrete masses that are easily swallowed.

Lamels (*Lamellæ*), made by rolling out a thin sheet of gelatine with an intimate mixture of active drugs, and dividing it into little squares, each of which contains a single dose of the drug. They may be swallowed as they are, or dissolved in water.

Discs (*Disci*), similar preparations, but of smaller size, containing a single dose of an alkaloid for hypodermic injection. To be dissolved in water when required.

Ophthalmic Discs. Similar discs, medicated with Atropin or Eserin, for insertion under the eyelids.

Capsules (*Capsulæ*). Single doses of drugs which possess nauseous flavours, enclosed in capsules of gelatin, to be swallowed whole.

THE WEIGHTS AND MEASURES.

WEIGHTS.		MEASURES.	
	1 grain.		1 minim.
437½ grains	= 1 ounce.	60 minims	= 1 fluid drachm.
16 ounces, or 7,000 grains	= 1 pound.	8 fluid drachms	= 1 fluid ounce.
		20 fluid ounces	= 1 pint.
		8 pints	= 1 gallon.

The grain in the above system is adopted from the "Troy" scale, the ounce from the "avoirdupois" or Imperial standard. The pint and gallon are the Imperial measures.

1 minim is the measure of	0.91 grains of water.
1 fluid drachm	" " 54.68 " "
1 fluid ounce	" " 437.5 " or 1 ounce, of water.
1 pint	" " 8750.0 " or 1½ pound "
1 gallon	" " 70,000.0 " or 10 pounds "

In the Metrical System :—

1 gramme	= 15.432 grains.
1 litre	= 1 pint, 15 oz., 2 drms., 11 min.

A litre is the measure of a thousand grammes (1 kilo.) of water.

A millilitre (its thousandth part) is therefore the measure of 1 gramme of water, and is equivalent to 16.93, or practically 17, minims.

A cubic centimetre is equivalent to a millilitre.

SYMBOLS.

The following are customary :—

Gr. (granum), ℥ (minimum), ℥ (drachma), ℥ (uncia), lb. (librum), O (Octarius—pint).

℥ (scrupulum) and ℥ (drachma) are legally used in prescribing solids to denote 20 grains and 60 grains respectively.

SECTION I.

DRUGS FURNISHED DIRECTLY BY THE ANIMAL
KINGDOM. 19 OFFICINAL.

DIVIDED INTO THREE GROUPS.

FIRST GROUP.

The following nine well known substances are introduced into the Pharmacopœia, all but the last being used simply as vehicles.

Lac. Fresh cows' milk.

Mel. Honey.

Mel Depuratum. Clarified Honey, *i.e.*, Honey melted over a water bath and strained through flannel.

Adeps Præparatus. Clarified Lard.

Sevum Præparatum. Clarified Mutton Suet.

Cera Flava. Bees-wax (Yellow).

Cera Alba. White-wax, *i.e.*, Bees-wax bleached by exposure to moisture, air, and light.

Albumen Ovi. The Whites of Fowls' Eggs.

Ovi Vitellus. The Yolks " "

Lard melts at about 100-deg. F.

Mutton-suet " 103 "

Bees-wax " 140 "

White-wax " 150 "

Suet is soluble in boiling Rectified Spirit.

Lard " Ether, and Oil of Turpentine.

White-wax " Oil of Turpentine.

Bees-wax " " leaving about a fourth undissolved.

Neither form of wax yields anything to cold Rectified Spirit.

OFFICINAL PREPARATIONS.

Oxymel. Clarified Honey, with an eighth part of Ac. Aceticum and the same amount of water.

Adeps Benzoatus. Clarified Lard heated with Gum-Benzoin, of which it takes up about 7 grains to the ounce.

Unguentum Simplex. A mixture of White-wax, Lard, and Almond Oil.

Mistura Spiritus Vini Gallici:—The Yolks of two Eggs, mixed with 4 oz. of Brandy, the same amount of Aq. Cinnamoni, and $\frac{1}{2}$ oz. of sugar.

*** *Ovi Vitellus* is introduced only to make the last-named preparation. *Lac* only for *Mistura Scammonii*. *Mel Depuratum* furnishes *Oxymel*, *Oxymel Scillæ*, and *Mel Boracis*. *Lard*, *Suet*, and *Wax* are used in preparing Ointments, Plasters, Suppositories, and Papers. No pharmaceutical use for *white of egg* is suggested in the Pharmacopœia.

SECOND GROUP :—ONE LIVING ANIMAL.

Hirudo. The Leech. Two species are defined, *Sanguisuga Medicinalis* (Speckled Leech) and *S. Officinalis* (Green Leech).

Both are wrinkled transversely and have olive-green backs with six rusty-red longitudinal stripes. The belly in the latter is olive-green also ; in the former it is greenish-yellow, with black spots.

Leeches are collected in France, Spain, Italy, and Hungary. They are preserved in pans of fresh soft water.

The Leech attaches itself to the skin by muscular sucking-discs, one at each extremity. On the anterior, or smaller, of the two opens the mouth, which is tri-radiate and surrounded by three cartilaginous jaws, each armed with a double row of minute teeth, by means of which the animal saws into the skin.

On an average, a Leech draws $1\frac{1}{2}$ drachm of blood ; at most 4 drachms. When satiated, it usually drops off spontaneously.

Bleeding is generally encouraged afterwards by fomentations. The amount of afterflow varies greatly, according to the region and the individual. If not watched, it may, in some cases, go on indefinitely.

THIRD GROUP :—NINE DRUGS.

The first is merely a colouring agent; the last is chiefly used as a vehicle. The remainder possess more or less valuable medicinal properties. Cantharis, Pepsin, and Cod-liver Oil are important drugs.

Coccus. *Cochineal.* The female Cochineal Insect, *Coccus Cacti*, killed and dried. Mexico and Teneriffe.

Contains *Carmin*.

If collected during pregnancy, coated with a white "bloom", forming *Silver Cochineal*.

Cantharis. *Cantharides.* The Spanish Fly, *Cantharis Vesicatoria*, killed and dried. Hungary.

About .4 p. e. of *Cantharidin*, $C_5H_{12}O_2$, the active principle.

Cantharidin is soluble in Chloroform, Ether, and strong Acetic Acid.

It sublimes unchanged.

Castoreum. *Castor.* The preputial (sebaceous) follicles of the Beaver, *Castor Fiber*, with their contained secretion. Hudson's Bay.

The secretion contains a *volatile oil*.

Moschus. *Musk.* The secretion of the preputial follicles of the Musk-deer, *Moschus Moschiferus*, dried. Central Asia.

Contains a *volatile oil*.

Fel Bovinum Purificatum. *Ox-gall.* An extract made with Rectified Spirit from fresh Ox-bile.

Contains all the ingredients of the bile but the mucus.

Soluble in water.

Pepsin. The contents of the gastric glands of a pig, sheep, or calf, mixed with some epithelium, dried and powdered. Obtained by scraping the inner surface of the stomach (after washing) with a blunt knife, and drying the scrapings at a low temperature.

Insoluble in water or spirit.

Contains *Pepsin*, the active principle of the Gastric Juice.

The properties of Pepsin are destroyed by a heat over 120 deg. F.

Cetaceum. *Spermaceti.* Nearly pure *Cetin*, $C_{32}H_{64}O_2$, with a little oily matter, separated from the oil contained in a cavity in the head of the Sperm whale, *Physeter Macrocephalus*. Pacific and Indian Oceans.

Crystalline.

Fuses at 112 deg. F.

Soluble in fixed oils.

Oleum Morrhue. *Cod-liver Oil.* Oil extracted from the fresh liver of the Cod (*Gadus Morrhua*) at a heat not exceeding 180° F. North Sea.

Pale yellow.

If prepared at higher temperatures, or from livers that have been kept, it has a brown tint, deeper according to the degree of temperature and the staleness of the livers. The pale yellow is the only official oil, and is generally preferred in medicine.

Contains 80 p. e. of *Olein*, 15 p. e. of *Margarin*, some biliary principles, and traces of Iodine and Bromide.

Other species of *Gadus* yield the same oil, but in less amount.

Saccharum Lactis. *Milk Sugar.* On evaporating whey, the sugar separates and crystallizes on pieces of cord or wood placed in the fluid.

Hard crystals of pure *Lactose*, $C_{12}H_{24}O_{12}$.
Soluble in 6 parts of water, not in Rectified Spirit.

* * Besides the above nineteen official drugs, the Animal Kingdom furnishes none that can be said to be commonly used in medicine. *Pancreatin*, the active agent of the Pancreatic secretion, is used in preparing fatty matters as aliments. *Dugong Oil* has been used as a substitute for Cod-liver Oil. *Bone Ash (Os Ustum)*, the source of Phosphorus and Phosphates, and *Bone Black (Carbo Animalis)*, the source of purified Animal Charcoal, are considered in Section III of this book.

OFFICIAL PREPARATIONS.

Tinctura Cocci.

Contains Carmine.

Tinctura Castorei.

Red colour.

Tinctura Cantharidis. Strength, 1 in 80.

Acetum Cantharidis. An extractive solution in Acetic Acid, strengthened by addition of the Glacial Acid, made at a temperature of 200° F. 1 in 10.

Liquor Epispasticus. *Blistering Fluid.* An extractive solution of Cantharides in Acetic Acid and Ether. 1 in $2\frac{1}{2}$.

Emplastrum Cantharidis. 1 in 3.

Emplastrum Calefaciens. 1 of Cantharides in 25.

Charta Epispastica. *Blistering Paper.* Cartridge paper coated with a plaster of Cantharides. 1 in 9.

Unguentum Cantharidis. 1 in 7.

Unguentum Cetacei.

The constituents of the various Ointments, Plasters, etc., will be found in the "Synopsis of Compound Preparations".

* * Milk Sugar is little used, except as a vehicle for Powders, as in the official *Pulvis Elaterii Co.*

SECTION II.

DRUGS FURNISHED DIRECTLY BY THE VEGETABLE
KINGDOM. 197 OFFICINAL.

First	Group . . .	Fungi and Lichens, 3.
Second	„ . . .	Plants Entire, 3.
Third	„ . . .	Green Tops, 2 ; Twigs, 1.
Fourth	„ . . .	Flowerings Tops, 2 ; Strobile, 1.
Fifth	„ . . .	Flowers 7.
Sixth	„ . . .	Buds, 2.
Seventh	„ . . .	Woods, 4.
Eighth	„ . . .	Barks, 13.
Ninth	„ . . .	Leaves, 13.
Tenth	„ . . .	Roots and Rhizomes, 25 ; Corm., 1 ; Bulb, 1 ; Tubers, 1.
Eleventh	„ . . .	Fruits, 26 ; Glands from Fruit, 1.
Twelfth	„ . . .	Seeds, 14 ; Gossypium.
Thirteenth	„ . . .	Fixed Oils, 7.
Fourteenth	„ . . .	Volatile Oils, 23.
Fifteenth	„ . . .	Resins, 7.
Sixteenth	„ . . .	Oleo-Resins, 4.
Seventeenth	„ . . .	Balsams, 4.
Eighteenth	„ . . .	Gums, 2.
Nineteenth	„ . . .	Gum-Resins, 6.
Twentieth	„ . . .	Concrete Juices, 8 ; Excrescence, 1.
Twenty-first	„ . . .	Active Principles of Definite Composition (Carbohydrates, Acids, Neutral Principles, Alkaloids), 14.

The Official Preparations are described at the end of each group.

Pharmacopœial Prescriptions for the extraction of Alkaloids and Active Principles.

Botanical Synopsis of the Natural Orders and Plants furnishing the Official Drugs, with their Geographical Sources and Preparations.

The Chief Non-official Drugs furnished by the Vegetable Kingdom.

FIRST GROUP:—FUNGI AND LICHENS. 3.

Ergot is the only important drug in this group.

Ergota. *Ergot of Rye.* (*Spurred Rye*, or *Secale Cornutum*.)

The compact mycelium (sclerotium) forming a stage in the life-history of the fungus, *Claviceps Purpurea*.

The fungus is parasitic on the ovary of *Secale Cereale* (common rye).

The sclerotium grows from beneath the ovary, and replaces the grain within the paleæ.

Contains 10 p.c. of *Ergotin*, a brown bitter extract of doubtful composition, and 35 p.c. of a *fixed oil*.

Ergotin is soluble in water and alcohol.

Cerevisiæ Fermentum. *Yeast.*

Consists almost entirely of round or oval cells, forming the fungus *Torula Cerevisiæ*. During the fermentation of malt infusion they multiply actively, being the immediate cause of the conversion of sugar into alcohol.

Cetraria. *Iceland Moss.* *Cetraria Islandica*, a lichen, forming a foliaceous thallus.

Contains 80 p.c. of *starchy* matters, and a bitter principle, *Cetrariæ Acid*, only slightly soluble in water.

OFFICINAL PREPARATIONS.

Extractum Ergotæ Liquidum (see p. 4).

Contains the ergotin, but no oil. 1 fl. oz. corresponds to 1 oz. of Ergot. The preparation ordinarily used.

Infusum Ergotæ.

One-fortieth the strength of the Liquid Extract.

Tinctura Ergotæ.

One-fourth the strength of the Liquid Extract.

Decoctum Cetrariæ.

A pint corresponds to an ounce of the moss. Contains the starches, but not much of the bitter.

Cataplasma Fermenti.

Made with 14 oz. of flour, and six fluid ounces of yeast mixed with the same of water at 100° F. Placed near a fire to "rise".

SECOND GROUP :—PLANTS ENTIRE. 3.

Lobelia is the only important drug in this group.

Chirata. *Chiretta* (*Ophelia Chirata*. Ord. Gentianaceæ).

Stems, roots, small flowers and capsules.

Contains a *bitter* principle, soluble in water and alcohol.

Lactuca. *Wild Lettuce* (*Lactuca Virosa*. Ord. Compositæ).

Collected while flowering.

Contains *Lactucin*, a crystalline principle resembling mannite, soluble in water, and other unimportant principles.

Lobelia. *Indian Tobacco* (*Lobelia Inflata*. Ord. Lobeliaceæ).

Stem, root, leaves, flowers and capsules.

Contains a liquid alkaloid, *Lobelina*, and a *volatile oil*.

OFFICIAL PREPARATIONS.

Extractum Lactucæ (p. 3).

Infusum Chiratæ.

Tinctura Chiratæ.

Tinctura Lobeliæ.

Tinctura Lobeliæ Ætherea (p. 5). } Same strength.

THIRD GROUP :—GREEN TOPS, 2. TWIGS, 1.

Sabina. *Savin Tops*. From *Juniperus Sabina* (Coniferæ).

Young branches enveloped in imbricated leaves, dried.

Contains **Oleum Sabinæ** (see "Volatile Oils", Fourteenth Group), *Gallic Acid*, and a *resin*.

Scoparium. *Broom Tops*. From *Sarothamnus Scoparius* (Leguminosæ).

Young branches, green and dried.

Contains *Scoparin*, a neutral principle, and *Sparteia*, a volatile bitter liquid alkaloid, both soluble in water and alcohol.

Dulcamara. *Bitter Sweet, Woody Nightshade*. From *Solanum Dulcamara* (Solanaceæ).

Young twigs, after leaf-fall, dried.

Contains an alkaloid, *Solanina*, and a bitter-sweet glucoside, *Dulcamarin*.

OFFICIAL PREPARATIONS.

Succus Scoparii.

Tinctura Sabinæ.

Decoctum Scoparii.

Unguentum Sabinæ.

Infusum Dulcamaræ.

FOURTH GROUP :—FLOWERING TOPS, 2. STROBILE, 1.

The first two are important.

Cannabis Indica. *Indian Hemp, Bhang, Gunjah, Hashish.* From *Cannabis Sativa* (Cannabinaceæ).

Twigs, leaves, remains of flowers, and a few ripe fruits, compressed.

The plant is ordinary Hemp, grown in India. Grown in Europe, it has not the same medicinal properties. The Indian is alone officinal.

Contains a *resin* ("Cannabin"), soluble in Rectified Spirit, not in water; and a little *volatile oil*. The former is the active principle.

Cusso. *Kousso.* From *Brayera Anthelmintica* (Rosaceæ).

Twigs, flowers, and unripe fruits.

Contains a *volatile oil*, an acrid *resin*, *Tannic Acid*, and *Koussin*, a crystallizable principle soluble in water.

OFFICIAL PREPARATIONS.

Extractum Cannabis Indicæ. Contains the resin (p. 4).

Tinctura Cannabis Indicæ. A solution of the extract in Rectified spirit, $2\frac{3}{4}$ gr. in the drachm.

The resin is precipitated on dilution with water.

Infusum Cusso. Unstrained, containing the powdered drug suspended.

(STROBILE.)

Lupulus. *Hop.* From *Humulus Lupulus* (Cannabinaceæ).

Bracts and achenes, and minute glands forming a yellow powder, "*Lupulin*," which amounts to about a sixth of the whole.

Lupulin contains 2 p. c. of *volatile oil*, and 10 p. c. of a bitter principle, *Humulin*, soluble in water and alcohol.

The bracts contain *Tannic Acid*.

OFFICIAL PREPARATIONS.

Extractum Lupuli. (See p. 4).

Infusum Lupuli.

Tinctura Lupuli.

FIFTH GROUP :—FLOWERS. 7.

Chamomile is the only drug of any importance in this group. Red-rose is used medicinally in the Acid Infusion. The remainder are colouring and flavouring agents.

Anthemis. *Chamomile.* From *Anthemis Nobilis* (Compositæ).

Both single and double flower-heads.

Contains **Oleum Anthemidis** (see "Volatile Oils", Fourteenth Group), and a *bitter* principle. The former is chiefly in the disk, therefore most abundant in the "single" flowers.

Aqua Aurantii Floris. *Orange-flower Water.* From *Citrus Aurantium* and *Bigaradia*, *Sweet* and *bitter Orange*. (Aurantiaceæ).
Distilled in France.

Contains a *volatile oil*, and some free *Acetic Acid*.

Crocus. *Saffron.* From *Crocus Sativus* (Iridaceæ).

The stigmata, with part of the styles, only.

Contains orange-red colouring matter, *Polycroite*, soluble in water and alcohol.

Rhæas. *Red Poppy.* From *Papaver Rhæas* (Papaveraceæ).

The petals.

Contains a red colouring matter, soluble in water and alcohol.

Rosa Centifolia. *Cabbage Rose.* From *Rosa Centifolia*. (Rosaceæ.)

The petals, fresh.

Contains a volatile oil, "*Attar of Roses*", soluble in alcohol, slightly in water.

Rosa Gallica. *Red Rose.* From *Rosa Gallica*.

The petals.

Contains *Tannic* and *Gallie Acids*, red colouring matter, a trace of *volatile oil*.

Sambucus. *Elder Flower.* From *Sambucus Nigra* (Caprifoliaceæ).

The fresh flowers.

Contains a *volatile oil*.

OFFICIAL PREPARATIONS.

Extractum Anthemidis. Strengthened with Ol. Anthemidis.

Aqua Aurantii Floris.

Aqua Rosæ (from Rosa Centifolia). Colourless.

Aqua Sambuci.

Infusum Anthemidis.

Infusum Rosæ Acidum. Made from Rosa Gallica, with 6 min. of Ac. Sulphuricum Dil. in the ounce.

Tinctura Croci. Orange red.

Syrupus Aurantii Floris. Aq. Aur. Flor. sweetened.

Syrupus Rosæ Gallicæ.

Syrupus Rhæados. Red.

Confectio Rosæ Gallicæ.

SIXTH GROUP :—BUDS. 2.

Santonica is the source of Santonin. Clove is little more than a flavouring agent.

Caryophyllum. *Cloves.* From Caryophyllus Aromaticus (Myrtaceæ).
Contains **Oleum Caryophylli** (see "Volatile Oils", Fourteenth Group).

Santonica. *Worm Seed.* From Artemisia Pauciflora (Compositæ).
Unexpanded flower-heads.
Contains *Santonin* (**Santoninum**, see Twenty-first Group), and a *volatile oil*.

OFFICIAL PREPARATION.

Infusum Caryophylli.

SEVENTH GROUP :—WOODS. 4.

Sandal Wood is merely a colouring agent. The others are important.

Guaiacum. *Lignum Vitæ.* From *Guaiacum Officinale* (Zygophyllaceæ).

Contains a *resin* (**Guaiaci Resina**, see Fifteenth Group), soluble in Rectified Spirit, ether, and alkaline solutions, not in water.

Hæmatoxylon. *Logwood.* From *Hæm. Campechianum* (Leguminosæ).

Contains a red crystalline colouring matter, *Hæmatoxylin*, soluble in alcohol, slightly in water: and *Tannic Acid*.

Pterocarpus. *Red Sandal Wood.* From *Pterocarpus Santalinus* (Leguminosæ).

Contains a red crystalline colouring matter, *Santalin*, soluble in alcohol, alkaline solutions, and Acetic Acid, not in water; and *Gallic Acid*.

Quassia. From *Picræna Excelsa* (Simarubaceæ).

Contains a crystalline neutral bitter principle, *Quassin*. Some Starch. No Tannin.

OFFICIAL PREPARATIONS.

Extractum Hæmatoxyli.

Extractum Quassiæ.

Infusum Quassiæ. Made with cold water, on account of the starch in the wood.

Tinctura Quassiæ.

Decoctum Hæmatoxyli. Flavoured with cinnamon.

*** *Pterocarpus* is used to colour Tr. *Lavandulæ Co.* There are no preparations of *Guaiac Wood*; of the *Resin*, a mixture and an ammoniated tincture (See Fifteenth Group). *Guaiac Wood* is used in preparing Dec. *Sarsæ Co.*

EIGHTH GROUP :—BARKS. 13.

The Cinchona Barks and Bebeeru Bark are of high importance. Canella is merely a flavouring agent. The rest are of minor importance.

Canella Alba. From Canella Alba (Canellaceæ).

Contains a *volatile oil* and a *bitter* principle.

Cascarilla. From Croton Eluteria (Euphorbiaceæ).

Contains a *volatile oil* and a *bitter* principle.

Cinchona Flava. *Yellow Bark.* From Cinch. Calisaya (Rubiaceæ).

Two kinds :—quilled and flat.

Contains the alkaloids—*Quinia*, *Cinchonin*, *Quinidin*, *Cinchonidin*; *Quinic*, *Quinovic*, and *Quino-tannic Acids*.

The active principles, as found in the bark, are soluble in alcohol, and sparingly in water.

Quino-tannic Acid gives a green precipitate with Ferric Salts.

The Sulphate of Quinia is officinal, as **Quiniæ Sulphas** (see “Alkaloids”, in the Twenty-first Group).

Yellow Bark yields from 2.2 to 3.8 p. c. of Quinia, and a similar amount of other alkaloids.

Cinchona Pallida. *Pale Bark.* From Cinchona Condaminea.

Quills only.

The same active principles, but in small amount. Yields .6 to 1.4 p. c. of alkaloids, chiefly *Cinchonin*.

Cinchona Rubra. *Red Bark.* From Cinchona Succirubra.

Both quills and flat pieces.

The same active principles, in variable amount. The best bark yields 2.6 p. c. of *Quinia*, and 4 p. c. of other alkaloids (about as much *Cinchonin* as *Quinia*).

Cinnamomum. *Cinnamon.* From Cinnam. Zeylanicum (Lauraceæ).

Contains **Oleum Cinnamomi** (see “Volatile Oils”, Fourteenth Group) and *Tannic Acid*.

Cusparia. *Angustura Bark.* From Galipea Cusparia (Rutaceæ).

Contains a *volatile oil*, and a bitter principle, *Angusturin*, soluble in alcohol, less so in water, precipitated by Tannic Acid.

Distinguished from Nux Vomica Bark, the chief *adulteration*, by the blood-red colour the Brucia in the latter strikes with Nitric Acid.

Granati Radix. *Pomegranate Bark.* From Punica Granatum (Myrtaceæ).

The bark of the root.

Contains 20 p. c. of *Tannic Acid*.

Larix. *Larch Bark.* From Larix Europæa (Coniferæ.)

Contains *resin*, *Tannic Acid*, and *Larixinic Acid*, crystalline and volatile, soluble in alcohol, sparingly in water.

Mezereon. From Daphne Mezereon (Thymelaceæ).

Contains an acrid *resin* and an acrid *volatile oil*, the former soluble in alcohol and ether.

Nectandra. *Bebeeru Bark.* From *Nectandra Rodiæi* (Lauraceæ).

Contains about $2\frac{1}{2}$ p. c. of the alkaloid *Beberia*.

Beberia Sulphas is officinal (see Twenty-first Group).

Quercus. *Oak Bark.* From *Quercus Pedunculata* (Cupuliferæ).

From the small branches and young stems only.

Contains about 6 p. c. of *Tannic Acid*.

Ulmus. *Elm Bark.* From *Ulmus Campestris* (Ulmaceæ).

The liber only.

Contains about 3 p. c. of *Tannic Acid*, and a large amount of *mucilage*.

OFFICINAL PREPARATIONS.

Extractum Mezerei Æthereum (See p. 4).

Pulvis Cinnamomi Co. Equal parts of cinnamon, cardamoms, and ginger.

Aqua Cinnamomi.

Infusum Cascarillæ.

Infusum Cuspariæ. Made with water at 120° F., not to disperse the volatile oil.

Decoctum Granati Radicis (See p. 5).

Decoctum Quercus.

Decoctum Ulmi. Mucilaginous.

*** Decoctions of Pomegranate and Oak Bark are practically solutions of Tannic Acid.

Tinctura Cascarillæ.

Tinctura Cinnamomi.

Tinctura Laricis.

(Of *Cinchona Flava*.)

Extractum Cinchonæ Flavæ Liquidum. 1 fl. oz. corresponds to 4 oz. of the bark.

Infusum Cinchonæ Flavæ. } Made with similar proportions
Decoctum Cinchonæ Flavæ. } of bark.

Tinctura Cinchonæ Flavæ.

(Of *Cinchona Pallida*.)

Tinctura Cinchonæ Co. With Serpentry; flavoured with orange peel, coloured with saffron and cochineal. A much weaker preparation than *Tr. Cinchonæ Flavæ*; contains very little Quinia.

* * * Of *Cinchona Rubra*, no preparations are officinal. It is used as a source of Quininc. *Nectandra* is used only as a source of *Beberia*. *Canella Alba* is used to flavour *Vin. Rhei*.

NINTH GROUP :—LEAVES. 13.

Nearly all highly important drugs.

Aconiti Folia. *Aconite Leaves.* From *Aconitum Napellus*, *Monkshood* (Ranunculacæ).

Contain the alkaloid **Aconitia** (see Twenty-first Group), but in less amount than the root **Aconiti Radix** (see Tenth Group).

Belladonnæ Folia. *Belladonna Leaves.* From *Atropa Belladonna*, *Deadly Nightshade* (Solanacæ).

Contain the alkaloid **Atropia** (see Twenty-first Group).

Belladonnæ Radix is also officinal (see Tenth Group).

Buchu. *Bucco.* From *Barosma Betulina*, *Crenulata*, and *Serratifolia* (Rutacæ).

Three kinds of leaf.

Contain a *volatile oil*, and a *bitter principle* soluble in water, not in alcohol.

Conii Folia. *Hemlock Leaves.* From *Conium Maculatum* (Umbelliferæ).

Contain the liquid volatile alkaloid *Conia*, $C_8 H_{15} N$, but in less amount than the fruit, **Conii Fructus** (see Eleventh Group).

Also a *volatile oil*, and a variable amount of *Methyl-Conia*, $CH_3 \cdot C_8 H_{14} N$. *Conia* is separated from its combination by adding Caustic Alkali to any part of the plant. It has an odour of *mice*. It is officinal in the form of an *Inhalation* (see pages 8 and 27).

Digitalis. From *Digitalis Purpurea*, *Foxglove* (Scrophulariaceæ).

Contain the neutral principle *Digitalin* (**Digitalinum**, see Twenty-first Group), soluble in alcohol, sparingly in water.

Hyoscyamus. *Henbane.* From *Hyoscyamus Niger* (Solanacæ).

Contain an alkaloid *Hyoscyamia*.

Laurocerasus. *Cherry Laurel.* From *Prunus Laurocerasus* (Rosacæ).

Contain the glucoside *Amygdalin*, which yields a *volatile oil*, and a small amount of *Prussic Acid*, on distillation.

Matica. *Matico.* From *Artanthe Elongata* (Piperacæ).

A thick down on the under surface.

Contain a *volatile oil*, and a *bitter principle*, *Artanthic Acid*, soluble in water and alcohol.

Senna Alexandrina. *Alexandrian Senna.* From *Cassia Lanceolata* and *Obovata* (Leguminosæ).

Two kinds of leaf.

Contain a yellowish-red neutral principle, *Cathartin*, of bitter and nauseous taste, soluble in water and alcohol. Also a *volatile oil* of nauseous odour and taste.

Cathartin is precipitated by Tannic Acid.

Senna Indica. *Tinnevelly Senna.* From *Cassia Elongata*.

The same constituents.

Stramonii Folia. *Stramony Leaves.* From *Datura Stramonium* (Solanaceæ).

Contain an alkaloid, *Daturia*, which appears to be identical with *Atropia*.

The proportion of alkaloid is much less than in *Belladonna*.

Stramonii Semina are also officinal (see Twelfth Group).

Tabacum. *Tobacco.* From *Nicotiana Tabacum* (Solanaceæ).

Unmanufactured tobacco.

Contain a liquid alkaloid, *Nicotia*, $C_{10}H_{14}N_2$, of acrid odour and taste.

Uva Ursi. *Bearberry.* From *Arctostaphylos Uva Ursi* (Ericaceæ).

Contain about 36 p. c. of *Tannic Acid* and 1.5 of *Gallic Acid*, and a little *volatile oil*.

OFFICIAL PREPARATIONS.

Extractum Aconiti (p. 3).

The remaining preparations of Aconite, the *Tincture* and *Liniment*, as well as the Alkaloid *Aconitia*, are prepared from the Root (see Tenth Group).

Extractum Belladonnæ.

Extractum Conii.

Extractum Hyoscyami.

Succus Belladonnæ.

Belladonna Liniment and the Alkaloid *Atropia* are prepared from *Belladonna Root* (see Tenth Group).

Succus Conii.

Succus Hyoscyami.

Infusum Buchu.

Tincture of Hemlock is prepared from the fruit (see Eleventh Group).

Infusum Digitalis.

The preparations of Stramony, the *Extract* and *Tincture*, are made from the seeds (see Twelfth Group).

Infusum Maticæ.

Infusum Sennæ.

Stramony leaves are only used in the form of cigarettes.

Infusum Uvæ Ursi.

Tinctura Belladonnæ.

The three kinds of Senna leaves are used indifferently.

Tinctura Buchu.

Tinctura Digitalis.

Tinctura Hyoscyami.

Tinctura Sennæ. Flavoured with raisins, caraway, and coriander.

Syrupus Sennæ. Flavoured with oil of coriander.

Confectio Sennæ. With coriander seed, figs, tamarinds, cassia, prunes, extract of liquorice, sugar and water.

Mistura Sennæ Co. A solution of Epsom Salts (*Magnesiæ Sulphas*) in Infusion of Senna, flavoured with Tincture of Senna, Extract of Liquorice, and Tr. Cardam. Co. One-fifth of an ounce of the salt in each fluid ounce.

Pil Conii Co. With Ipecacuanha and Treacle.

Cataplasma Conii. An ounce of Hemlock leaves in a poultice made with three ounces of Linseed (p. 9).

Vapor Coniæ. A drachm of Extract of Hemlock, a drachm of Liq. Potassæ, and ten of water, mixed. Twenty minims to be placed on a sponge over hot water, and the steam inhaled.

Unguentum Belladonnæ. One part of the extract in five.

Emplastrum Belladonnæ. Equal parts of the extract and resin plaster.

Aqua Laurocerasi.

Enema Tabaci. An infusion of 20 grains in 8 fluid ounces of water.

* * * Aqua Laurocerasi is, practically, a solution of Hydrocyanic Acid, of very uncertain strength.

* * * The strength of the preparations of Hemlock is very variable, as Conia is not only volatile, but is liable to decomposition in presence of air. Succus Conii is the most trustworthy. Tr. Conii, according to Dr. John Harley, is almost inert.

TENTH GROUP :—ROOTS AND RHIZOMES. 25.

CORM, 1. BULB, 1. TUBERS, 1.

Liquorice and Hemidesmus are merely flavouring agents ; Horse-radish, Arnica, Pellitory, Sassafras, Serpentry, and Sumbul are of small importance. The remainder are important drugs, and some of high importance.

Aconiti Radix. *Aconite Root.* From *Aconitum Napellus*, *Monkshood* (Ranunculaceæ). Root.

Contains **Aconitia** (see Twenty-first Group) in greater amount than the leaves **Aconiti Folia** (p. 25).

Armoracia. *Horse-radish.* From *Cochlearia Armoracia* (Cruciferae). Root.

Contains, or develops on being scraped, an acrid *volatile oil*, apparently identical with oil of mustard.

Arnica. From *Arnica Montana* (Compositæ). Rhizome and rootlets.

Contains a *volatile oil*, and an acrid *resin* soluble in alcohol.

Belladonnæ Radix. From *Atropa Belladonna*, *Deadly Nightshade* (Solanaceæ). Root.

Contains **Atropia** (see Twenty-first Group).

Belladonnæ Folia are officinal (see p. 25).

Calumba. From *Jateorrhiza Calumba* (Menispermaceæ). Root.

Contains a bitter alkaloid, *Berberia*, and a neutral bitter principle, *Calumbin*, slightly soluble in water and alcohol. Also *Starch*.

Colchicum. From *Colchicum Autumnale*, *Meadow Saffron* (Melanthaceæ). Corms, fresh and dried.

Contains an alkaloid, *Colchicia*.

Colchici Semina are officinal (see Twelfth Group).

Filix Mas. *Male Fern.* From *Aspidium Filix Mas* (Filices). Rhizome with rootlets and bases of petioles.

Contains a *fixed oil*, a *volatile oil*, and a *resin*, all soluble in ether.

Gentiana. *Gentian.* From *Gentiana Lutea* (Gentianaceæ). Root.

Contains a bitter principle *Gentianite*, soluble in water ; also *sugar*, *pectin*, and *gum*.

Glycyrrhiza. *Liquorice.* From *Glycyrrhiza Glabra* (Leguminosæ). Rhizome.

Contains a sweet glucosido, *Glycyrrhizin*, soluble in water and alcohol, precipitated by acids.

Hemidesmus. From *Hemidesmus Indicus* (Asclepiadaceæ). Root.

Ipecacuanha. From *Cephaelis Ipecacuanha* (Rubiaceæ). Root.

Contains an alkaloid, *Emetia*, and an astringent acid, *Cephaëlic Acid*, which strikes green with Ferric Salts.

Jalapa. *Jalap.* From *Exogonium Purga* (Convolvulaceæ). Tubers.

Contains 18 p.c. of **Resina Jalapæ** (see Fifteenth Group.)

Krameria. *Rhatany.* From *Krameria Triandra* (Polygalaceæ). Root.

Contains 40 p.c. of *Tannic Acid*, and a red astringent matter, soluble in water and alcohol.

Pareira. From *Cissampelos Pareira* (Menispermaceæ). Root.

Contains a bitter alkaloid *Pelosia*, possibly identical with *Beberia*.

Podophyllum. From *Podoph. Peltatum* (Ranunculaceæ). Rhizome.

Contains **Podophylli Resina** (see Fifteenth Group).

Pyrethrum. *Pellitory.* From *Anacyclus Pyrethrum* (Compositæ). Root

Contains *acid oils* and *resins*, and some *Tannic Acid*.

Rheum. *Rhubarb.* From undetermined species of *Rheum* (Polygonaceæ).

Root, deprived of its outer coat.

Three kinds, *Russian* or "*Turkey*", *East Indian*, and *Dutch-Trimmed*, or *Batavian*, all from China and Central Asia.

Contains* *Chrysophanic Acid* as yellow granules, sparingly soluble in cold water or alcohol, more so in hot, yielding red soluble chrysophanates with alkaline solutions; insoluble chrysophanates with lime-water, magnesia, and various metallic solutions.

Several *resins*, soluble in alcohol, not in water; and possibly a volatile oil.

Tannic and *Gallic Acids* and some *bitter* principle.

Calcic Oxalate, in masses of acicular crystals (*raphides*):—35 p.c. in Russian Rhubarb.

Sarsa. *Jamaica Sarsaparilla.* From *Smilax Officinalis* (Smilacæ). Roots, folded in bundles.

Contains a *volatile oil*, and a crystalline principle, *Smilacin*, soluble in alcohol, not in water. No starch.†

Sassafras. From *Sassafras Officinale* (Lauraceæ). Root.

Contains a *volatile oil*.

* The purgative principle of Rhubarb has not yet been isolated.

† Other varieties of Sarsaparilla contain Starch.

Scammonia Radix. *Scammony Root.* From *Convolvulus Scammonia* (Convolvulaceæ). Root.

Contains **Scammonia Resina** (see Fifteenth Group).

Scammonium, the *gum-resin* obtained from the root during life, is official (see Nineteenth Group).

Scilla. *Squill.* From *Scilla Maritima* (Liliaceæ). Bulb, sliced.

Contains an acrid *resin*, and a *bitter* principle, soluble in water, alcohol, and acetic acid. (Composition practically unknown.)

Senega. From *Polygala Senega* (Polygalaceæ). Root-stock and roots.

Contains a glucoside, *Senegin* or *Polygalic Acid*, soluble in water or alcohol; also a little *Tannic Acid*.

Serpentaria. *Serpentary.* From *Aristolochia Serpentaria* (Aristolochiaceæ). Rhizome and rootlets.

Contains a *volatile oil* and a *bitter principle*, soluble in water and alcohol.

Sumbul. From *Euryangium Sumbul* (Umbelliferae). Root.

Contains a *volatile oil*, two *balsamic resins*, one soluble in ether, one in alcohol, *starch* and *sumbulic acid*.

Taraxacum. *Dandelion.* From *Taraxacum Dens Leonis* (Compositæ). Root.

Yields a milky juice, which contains *caoutchouc*, *gum*, *albumen*, *sugar*, and a bitter crystalline substance, *Taraxacin*, soluble in alcohol, sparingly in water.

Valeriana. *Valerian.* From *Valeriana Officinalis* (Valerianaceæ). Rhizome and rootlets, smelling of Valerianic Acid.

Contains a hydrocarbon, *Borneene*, $C_{10}H_{16}$, and *Valerol*, $C_6H_{10}O$.

On exposure to air, valerol is oxidized to form *Valerianic Acid*, $C_5H_{10}O_2$.*

On distilling the root with water, a *volatile oil*, consisting of Borneene, Valerol, Valerianic Acid, and probably other substances, is obtained.

The above principles are soluble in alcohol, sparingly in water. They are all volatile.

Veratrum Viride. *Green Hellebore.* From *Veratrum Viride* (Melanthaceæ). Rhizome and rootlets.

Contains two active alkaloids, *Viridia* and *Veratroidea*, combined with *Gallic Acid*.

* In making the official Valerianates, the acid is produced by oxidizing Fusel-oil (amylic alcohol). See Sodæ Valerianas, in Section III.

Zingiber. *Ginger.* From *Zingiber Officinale* (*Zingiberaceæ*). Rhizome, scraped.

Contains a *volatile oil*, a pungent *resin*, soluble in alcohol, not in water, and a great deal of *starch*.

OFFICIAL PREPARATIONS.

Extractum Calumbæ.

- „ Colchici (see p. 4).
- „ Colchici Aceticum (see p. 4).
- „ Gentianæ.
- „ Glycyrrhizæ.
- „ Jalapæ.
- „ Krameriæ.
- „ Pareiræ.
- „ Rhei.
- „ Taraxaci (see p. 4).

Extractum Filicis Liquidum (p. 4).

- „ Glycyrrhizæ Liquidum.
- „ Pareiræ Liquidum.
- „ Sarsæ Liquidum.

Succus Taraxaci.

Infusum Calumbæ, made with *cold* water, on account of the starch.

- „ Gentianæ Co., flavoured with orange and lemon peels.
- „ Krameriæ.
- „ Rhei.
- „ Senegæ.
- „ Serpentariæ.
- „ Valerianæ.

Decoctum Pareiræ.

- „ Sarsæ.
- „ Sarsæ Co., made with Sassafras, Guaiac Wood, Mezereon, and Liquorice-root.
- „ Taraxaci.

Tinctura Aconiti.

- ” **Arnicae.**
- ” **Calumbæ.**
- ” **Gentianæ Co.**, flavoured with orange peel and cardamoms.
- ” **Jalapæ.**
- ” **Krameria.**
- ” **Pyrethri.**
- ” **Rhei**, flavoured with cardamoms and coriander, and coloured with saffron.
- ” **Scillæ.**
- ” **Senegæ.**
- ” **Serpentaria.**
- ” **Sumbul.**
- ” **Valerianæ.**
- ” **Valerianæ Ammoniata** (see p. 6).
- ” **Veratri Viridis.**
- ” **Zingiberis.**
- ” **Zingiberis Fortior**, four times the strength of Tr. Zingib.

Acetum Scillæ, a “Tincture” made with dilute Acetic Acid (p. 6); about the same strength as Tr. Scillæ.

Spiritus Armoraciæ Co. (p. 6).

Vinum Colchici.

- ., **Ipecacuanhæ**, nearly 1 gr. in the drachm.
- ., **Rhei**, flavoured with Canella Alba.

Syrupus Hemidesmi.

- ” **Rhei**, flavoured with Coriander.
- ” **Scillæ**, made with Vinegar of Squill.
- ” **Zingiberis**, made with Tr. Zingib. Fort.

Oxymel Scillæ. A mixture of Honey and Vinegar of Squill, concentrated.

Pulvis Glycyrrhizæ Co. Equal parts of Senna and Liquorice-root, with three times the amount of Sugar.

Pulvis Ipecacuanhæ Co. *Dover's Powder.* Equal parts of Opium and Ipecacuanha, with four times the amount of Potassæ Sulphas.

Pulvis Rhei Co. *Gregory's Powder.* Rhubarb, with thrice the amount of Light Magnesia and half the amount of Ginger.

Pulvis Jalapæ Co. Five parts of Jalap, one of Ginger, nine of Cream of Tartar.

ELEVENTH GROUP:—FRUITS, 26. GLANDS FROM FRUIT, 1.

- a. The following twelve and their preparations are of minor importance, being little more than vehicles and flavouring agents (see p. 3).

Anethum. *Dill.* From *Anethum Graveoleus* (Umbelliferæ).
Achenes.

Contains **Oleum Anethi** (see "Volatile Oils", Fourteenth Group).

Aurantii Fructus. *Bitter Orange.* From *Citrus Bigaradia* (Aurantiaceæ).
Fresh fruit. The peel only is used (see p. 35).

The outer part of the peel contains a *volatile oil*, the inner a *bitter principle* and a little *Gallic Acid*.

Aurantii Cortex. *Bitter Orange Peel.*

The outer part of the peel of the same, dried.

Aqua Aurantii Floris is officinal (see p. 20).

Caruum. *Caraway.* From *Carum Carui* (Umbelliferæ).
Achenes.

Contain **Oleum Carui** (see "Volatile Oils", Fourteenth Group).

Coriandrum. *Coriander.* From *Coriandrum Sativum* (Umbelliferæ).
Adhering Mericarps.

Contains **Oleum Coriandri** (see "Volatile Oils", Fourteenth Group).

Fœniculum. *Fennel.* From *Fœniculum Dulce* (Umbelliferæ).
Achenes.

Contains a *volatile oil*.

Limonis Succus. *Lemon-juice.* From *Citrus Limonum* (Aurantiaceæ).
The fresh juice

Contains *Citric Acid*, about 4 gr. in the drachm, some *Malic Acid*, and a little *bitter principle*; also some acid salts of Potassium.

Limonis Cortex. *Lemon-peel.*

The outer part of the same, fresh.

Contains **Oleum Limonis** (see "Volatile Oils", Fourteenth Group), a *bitter principle*, and some *Gallic Acid*.

Mori Succus. *Mulberry-juice.* From *Morus Nigra* (Moraceæ)

Contains *Tartaric Acid*, sugar, and red colouring matter.

Pimenta. *Pimento*, or *All-Spice*. From *Eugenia Pimenta* (Myrtaceæ),
Unripe berries, dried.

Contains **Oleum Pimentæ** (see "Volatile Oils", Fourteenth Group)
and *Tannic Acid*.

Rosa Canina. *Hips*. From *Rosa Canina*, *Dog-rose* (Rosaceæ).

Contains *Citric* and *Malic Acids* and their salts, sugar, and a little *Tannin*.
The hairy achenes are irritating to the bowels.

Uvæ. *Raisins*. From *Vitis Vinifera* (Vitaceæ).

Spanish Raisins only.

Contain *glucose* and *Acid Potassic Tartrate*

OFFICIAL PREPARATIONS.

Aqua Anethi.

„ **Carui.**

„ **Fœniculi.**

„ **Pimentæ.**

Infusum Aurantii. From the dried peel.

„ **Aurantii Co.** Ditto; flavoured with Lemon-peel and Cloves.

Tinctura Aurantii. From the dried peel.

„ **Aurantii Recentis.** From the outer part of the peel of
the fresh fruit.

„ **Limonis.** From the peel.

Syrupus Aurantii. Syrup, flavoured with Tr. Aurantii.

„ **Limonis.** Made with the fresh juice.

„ **Mori.** Ditto. Violet red.

Confectio Rosæ Caninæ. The "seeds" (achenes) are excluded.
Liable to candy on keeping.

- * * Coriander is used to flavour several preparations.
 Raisins to flavour Tr. Cardam. Co., and Tr. Sennæ.
 Conf. Ros. Caninæ is the basis of Pil. Quiniæ.
 Lemon-juice is often used as a solution of Citric Acid in making effervescent draughts. The juice of Lemons is the source of **Acidum Citricum** (see Section III).
Vinum Aurantii (see "Alcohol", Section III) is not a Pharmacopœial preparation, but the "*Orange Wine*" of the Grocers' shops.
-

β. The remainder are used for their medicinal properties, but are of variable importance.

Bela. *Bael-fruit.* From *Egle Marmalos* (Aurantiaceæ).

Half ripe fruit, dried.

Contains an *astringent principle*, supposed to be allied to Tannic Acid, but as yet incompletely studied.

Capsicum.* From *Capsicum Fastigiatum* (Solanaceæ).

Membranous pods, varying much in size.

Contains a red *acid oil*, volatilizable by heat, soluble in alcohol, sparingly in water.

Cassia. From *Cassia Fistula* (Leguminosæ).

The pulp from the pods.

Contains *sugar*, *mucilage*, and *pectin*. The active principle has not been isolated.

Colocynthis. *Colocynth.* From *Citrullus Colocynthis* (Cucurbitaceæ).

The fruit, freed from rind and seeds, dried.

Contains a bitter purgative principle, viz., the glucoside *Colocynthin*, soluble in water and alcohol.

Conii Fructus. *Hemlock Fruit.* From *Conium Maculatum* (Umbellifereæ).

Unripe fruit, dried.

Contains *Conia* and *Methyl-Conia* (see **Conii Folia**, p. 25).

Cubeba. *Cubebs.* From *Cubeba officinalis* (Piperaceæ).

Contains about 10 p.c. of **Oleum Cubebæ** (see "Volatile Oils", Fourteenth Group) a principle *Cubebin*, allied to Piperin, soluble in alcohol, and resins.

* Cayenne Pepper consists of powdered Capsicums. Capsicum is the chief ingredient of Curry-powders.

Ecbalium. *Squirting Cucumber.* From Ecb. Officinatum (Cucurbitaceæ).

A pepo, which, when ripe, separates from its stalk, and expels, through the aperture left at the insertion of the stalk, the seeds and the juice surrounding them.

Gathered before complete ripeness and separation from the stalk.

The sediment strained from the juice is alone used, under the name of **Elaterium** (see Twentieth Group).

Ficus. *Figs.* From Ficus Carica (Moraceæ).

Dried Figs.

The numerous achenes ("seeds") contained in the pulp are a mechanical stimulant to the intestines.

Kamala. *Wurrus.* From Rottleria Tinctoria (Euphorbiaceæ).

Minute superficial glands from the Capsules, forming a red powder.

Contains 80 p.c. of *resins* soluble in alcohol, a trace of *volatile oil* and of *tannin*.

Papaver. *Poppy Capsules.* From Papaver Somniferum (Papaveraceæ).

Nearly ripe, dried. A star-like stigma at the apex. Variable in size.

The capsule contains a minute quantity of the active principles found in Opium.

The seeds contain a fixed oil, *Poppy Oil*, which has no medicinal properties.

The same capsules in an unripe state, yield **Opium** (see Twentieth Group).

Piper. *Black Pepper.* From Piper Nigrum (Piperaceæ).

Berries ("Peppercorns").

"White Pepper" consists of decorticated peppercorns.

Contains a *volatile oil*, an acid *resin*, and a tasteless nitrogenised principle, *Piperin*, $C_{17}H_{19}NO_3$, which is colourless when quite free from the oil.

Prunum. *Prunes.* From Prunus Domestica (Rosaceæ).

Dried Plums.

Contains an undetermined purgative principle, *sugar*, and *malic acid*.

Rhamni Succus. *Buckthorn Juice.* From Rhamnus Cathartica (Rhamnaceæ).

The juice of the berries.

Contains a *purgative principle*, not yet accurately determined, and a *green colouring matter*.

Sabadilla. *Cevadilla.* From Asagraea Officinalis (Melanthaceæ).

Follicles, dried.

Contains **Veratria** (see "Alkaloids" in Twenty-first Group) and *Gallic Acid*.

Tamarindus. *Tamarind.* From *Tamarindus Indica* (Leguminosæ).

Pulp preserved in sugar. The West Indian is alone officinal.

Contains free *Acids*:—*Citric*, *Tartaric*, and *Malic*; *Acid Potassic Tartrate*, *sugar*, and *pectin*.

OFFICIAL PREPARATIONS.

Extractum Belæ Liquidum.

Extractum Colocynthis Co. A spirituous extract, mixed, before evaporation, with extract of Socotrine Aloes, Scammony resin, and hard soap; flavoured with powdered Cardamoms.

Extractum Papaveris. Of the Capsules alone, without the seeds.

Decoctum Papaveris. Of the seeds as well as the Capsules; contains mucilage and oil.

Syrupus Papaveris. Of the capsules only (p. 6).

Syrupus Rhamni. Flavoured with Ginger and Pimento (p. 6).

Tinctura Capsici.

Tinctura Conii. The only preparation of Hemlock fruit, the *extract* and *juice* being made from the leaves, the *Pill-mass* and *Inhalation* of Conia from the extract.

* * Said by Dr. John Harley to be an almost inert preparation.

Tinctura Cubebæ.

Confectio Piperis. Made with Honey, flavoured with Caraway.

Pil. Colocynthis Co. Made of Colocynth with twice the amount of Barbadoes Aloes and of Scammony, mixed with a little Sulphate of Potash, and scented with Oil of Cloves.

Pil. Col. cum Hyoscyamo. The same, with half as much of Extract of Henbane.

* * Cassia, Figs, Prunes, and Tamarind form part of Confection of Senna.

Cevadilla is used only as a source of Veratria, Ecballium as a source of Elaterium (see Twentieth Group).

TWELFTH GROUP:—SEEDS. 14. GOSSYPIUM.

. *Nux Vomica*, *Physostigma*, and *Stramonium*, are important and active drugs. The remainder are of minor importance as medicinal drugs, though some are in extensive use as vehicles.

Amygdala Dulcis. *Sweet, or Jordan, Almond.* From *Amygdala Communis* (Rosaceæ).

Large and long, easily distinguished from the Bitter Almond.

Contains **Oleum Amygdalæ** (see "Fixed Oils", Thirteenth Group), an albuminoid *Emulsin*, soluble in cold water, precipitated by boiling or by alcohol, sugar and gum.

Amygdala Amara. *Bitter Almond.* From the same species, var. *Amara*. Smaller and more ovate.

Contains, besides the above, a white crystalline glucoside, *Amygdalin*, $C_{20}H_{27}NO_{11}$, soluble in water and alcohol, slightly bitter. When water is added to the almond, the *Amygdalin* decomposes in the presence of the *Emulsin*, furnishing *glucose*, *Hydrocyanic Acid*, and volatile *Oil of Bitter Almonds* (Benzoyl Hydride,) C_7H_6O .*

Areca. *Betel nut.* From *Areca Catechu* (Palmaceæ).

Contains *Tannic* and *Gallic Acids*, with a red colouring matter, soluble in alkaline solutions and boiling water.

Cardamomum. *Cardamoms.* From *Elettaria Cardamomum* (Zingiberaceæ).

Usually kept in the pericarps, but the seeds alone are official.

Contains a *volatile oil*, about $4\frac{1}{2}$ p.c., and *colouring matter*.

Colchici Semina. *Colchicum Seeds.* From *Colch. Autumnale* (Melanthaceæ).

Contain an alkaloid, *Colchicia*, resembling *Veratria* in properties.

The *Corm* also is official (see p. 28).

Farina Triticci. *Wheat-flour.* From *Triticum Vulgare* (Graminaceæ).

Contains 52-75 p.c. of *starch* (see Twenty-first Group), 10-23 p.c. of *gluten* and *albuminoids*, 6-13 p.c. of *dextrin* and *glucose*, with *Potassium* and other *salts*.

Hordeum Decorticatum. *Pearl Barley.* From *Hordeum Distichon* (Graminaceæ).

Barley, husked in a mill, the grains polished.

Contains on an average 64 p.c. of *starch*, 14 of *gluten*, *albuminoids*, etc., 2 of *fatty matter*, 2 of *saline matter*.

* $C_{20}H_{27}NO_{11} + 2H_2O = 2C_6H_{12}O_6 + HCN + C_7H_6O$. Commercial Oil of Bitter Almonds retains the *Hydrocyanic Acid*, to the extent of 4-8 p.c. The oil itself is not poisonous, and may be obtained in a pure state.

Lini Semina. *Linseed.* From *Linum Usitatissimum*, *Flax* (Linaceæ).

Contains 20 p.c. of **Oleum Lini** (see "Fixed Oils", Thirteenth Group) in the kernel. The seed-coats contain *mucilage*.

Lini Farina. *Linseed-meal.* The "Oil-cake" or residue of Linseed after the expression of the oil, ground to powder.

Contains the *mucilage*.

Crushed Linseed, with the oil in it, grows rancid on keeping. The meal, from which the oil has been expressed, keeps well.

Myristica. *Nutmeg.* From *Myristica Officinalis* (Myristicaceæ).

The kernel invested with the closely adherent inner seed coat (dark brown), which dips down into the substance of the nutmeg, producing a veined appearance on section.

The "Arillode" of the nutmeg constitutes *Mace*.

Contains the concrete fixed oil **Oleum Myristicæ Expressum** (see Thirteenth Group) and the volatile oil **Oleum Myristicæ** (see Fourteenth Group).

Nux Vomica. From *Strychnos Nux Vomica* (Loganiaceæ).

Contains the alkaloids **Strychnia** (see Twenty-first Group) and *Brucia*, with *Strychnic* or *Igasuric Acid*.

The red colour which *Brucia* strikes with Nitric Acid is a test for parts of the *Nux Vomica* plant. See *Cusparia Bark*, p. 23.

Physostigma. *Calabar Bean.* *Eseré.* From *Physostigma Venenosum* (Leguminosæ).

Contains an alkaloid, *Eserin* (see "Non-officinal Drugs").

Sinapis. *Mustard.* From *Sinapis Alba* and *Nigra*. (Cruciferae.)

The seeds of both plants, powdered.

Both white and black seeds contain about 28 p.c. of a mild *fixed oil*, and an albuminoid substance, *Myrosin*, resembling Emulsin (see "*Amygdala Amara*" above).

White mustard seeds contain a glucoside, *Sinalbin*, which yields glucose and an acrid *fixed oil* in presence of Myrosin and water.

Black mustard-seeds contain the glucoside *Myronic Acid*, as Potassium Myronate, $\text{KC}_{10}\text{H}_{18}\text{NS}_2\text{O}_{10}$. In presence of water at about 120 deg. F., Myrosin determines decomposition of the Myronate, forming glucose, Hydric-Potassic Sulphate, and an acrid volatile oil, **Oleum Sinapis** (see Fourteenth Group).

Stramonii Semina. *Stramony.* From *Datura Stramonium* (Atropaceæ).

Contain an alkaloid, *Daturia*, which appears to be identical with Atropia; and a *fixed oil*.

Stramonii Folia are also officinal (see p. 26).

Gossypium. *Cotton Wool.* From various species of *Gossypium* (Malvaceæ).

Fine hairs, formed of cells placed end to end;* attached to the seed-coat.

Nearly pure *Cellulose*. Soluble in strong alkaline solutions. See *Pyroxylin* in Sec. III.

* Flax-fibres are formed of cells with tapering ends, overlapping.

OFFICIAL PREPARATIONS.

Extractum Nucis Vomicæ.

Extractum Physostigmatis.

Extractum Stramonii (p. 4).

Infusum Lini. Flavoured with Liquorice Root. Contains Mucilage, holding some of the oil.

Decoctum Hordei. (*Barley-water*) 2 oz. in a pint and a half.

Tinctura Cardamomi Co. A tincture of Cardamom, Caraway, Raisins, and Cinnamon, coloured with Cochineal.

Tinctura Colchici Seminum.

Tinctura Nucis Vomicæ.

Tinctura Stramonii.

Pulvis Amygdalæ Co. Sweet Almonds, powdered, with Sugar and Gum.

Mistura Amygdalæ. An emulsion made by rubbing Pulv. Amyg. Co. with water, strained through muslin. $2\frac{1}{2}$ oz. of the powder to the pint.

Cataplasma Lini. A poultice made with 4 oz. of Linseed-meal, half an ounce of olive oil being added when it is made.

Cataplasma Sinapis. A poultice made with $2\frac{1}{2}$ oz. each of Linseed-meal and mustard.

Charta Sinapis. Cartridge paper coated with *Black Mustard* by means of Liq. Gutta-percha.

* * * **Nux Vomica** Seeds are softened by steaming before being powdered for use.

The other Colchicum preparations, the *Extract*, *Acetic Extract*, and *Wine*, are made from the *Corm* (pp. 30, 31).

Nutmeg enters into several compound preparations. (See the "Synopsis".) *Spirit* of Nutmeg is made with the Volatile Oil (see Fourteenth Group).

Both Sweet and Bitter Almonds yield Ol. Amygdalæ (pp. 39, 42); the former only is used for the Compound Powder and the Mixture.

Wheat-flour is used to make Yeast-poultice (p. 17).

Pyroxylin (*Gun-cotton*) is prepared from Cotton wool. (See Section III.)

THIRTEENTH GROUP:—FIXED OILS. 7.

Croton and Castor Oils are important drugs; Nutmeg-butter is used for external application; the rest are vehicles.

Fixed oils are composed of Glyceric Salts of Acids of the Acetic or the Oleic Series. They cannot be distilled unchanged. With caustic alkalies, and some metallic oxides, they yield the corresponding alkaline or metallic salts, and glycerine. (See Glycerinum; Sapo Mollis, under "Potassium"; Sapo Durus and Animalis, under "Sodium"; Emplastrum Plumbi, under "Lead"; in Section III.)

They are extracted by expression, and in the case of the liquid oils, are of finest quality when expressed without the aid of heat. Heat either produces partial decomposition, or causes them to be contaminated with other constituents of the plant. Cold expression is expressly demanded by the Pharmacopœia in the case of Linseed oil.

The concrete oils, which are solid at ordinary temperatures, require the aid of heat in expression.

Oleum Amygdalæ. *Almond Oil.* From the seeds (Almonds) of *Amygdalus Communis*, sweet and bitter varieties (Rosaceæ) see p. 39.

Chiefly *Glyceric Oleate (Olein)*.

Ol. Crotonis. *Croton Oil.* From the seeds of *Croton Tiglium* (Euphorbiaceæ).

Composition undetermined. *Crotonic Acid* has been isolated.

Ol. Lini. *Linseed Oil.* From the seeds of *Linum Usitatissimum* (Linaceæ) (p. 40).

Chiefly *Glyceric Lin-oleate* and *Margarate*.

Ol. Myristicæ Expressum. *Nutmeg-butter.* From the kernels of *Myristica Officinalis* (Myristicaceæ) p. 40.

A concrete oil: contains a little volatile Oil of Nutmeg.

Myristic Acid has been isolated.

Ol. Olivæ. *Olive, or Sweet, Oil.* From the fleshy fruit of *Olea Europæa* (Oleaceæ).

28 p.c. *glyceric Margarate* (Margarin), 72 p.c. *glyceric Oleate* (Olein).

Ol. Ricini. *Castor Oil.* From the seed of *Ricinus Communis* (Euphorbiaceæ).

Glyceric Ricinate, Ricin-oleate and Ricin-stearate, with some acrid resin.

Ol. Theobromæ. *Cocoa-butter.* From the seeds of *Theobroma Cacao*. (Byttneriaceæ).

A concrete oil. Melts at 122° F.

The above oils are all soluble in ether, and insoluble in water.

The liquid oils are miscible one with the other in all proportions.

Castor oil mixes with absolute alcohol in all proportions, and is soluble in twice its volume of Rectified Spirit.

Croton oil, when freshly expressed without the aid of heat, is soluble in equal parts of alcohol, but by keeping loses its solubility.

The other fixed oils are only slightly soluble in alcohol. Admixture of any one with Castor Oil renders it more soluble in alcohol than the oil alone.

By prolonged exposure to air the fluid oils partly decompose, forming acrid products, and "becoming rancid."

Olive oil is the least liable to such decomposition.

Linseed and Castor oils become solid by prolonged exposure. They are therefore "drying oils."

At a temperature of 36° F., the margarin in Olive and Linseed oils solidifies and separates.

OFFICIAL PREPARATIONS.

Linimentum Crotonis. A solution in equal parts of Oil of Cajeput and Rectified Spirit, 1 in 8.

* * Almond Oil is used in forming various ointments (see "Compound Preparations") and is the basis of *Oleum Phosphoratum*.

Olive Oil is used in forming various ointments, liniments, and plasters; also *Charta Epispastica*, *Cataplasma Lini*, and *Enema Magnesic Sulphatis*.

Castor Oil enters into *Flexile Collodion*, *Compound Calomel Pill*, and *Lin. Sinapis Co*.

Nutmeg-butter into *Empl. Calefaciens* and *Empl. Picis*.

Cocoa-butter is used as a basis for Suppositories (see p. 9).

FOURTEENTH GROUP :—VOLATILE OILS. 23.

The volatile oils of plants consist of Hydrocarbons, or of mixtures of Hydrocarbons with oxygenated derivatives, or of the oxygenated compounds alone.

Either a *Terpene*, $C_{10}H_{16}$, or a hydrocarbon of the composition $C_{15}H_{24}$, enters into the constitution of nearly all.

Oils of Turpentine, Savin, and Copaiba consist purely of Terpenes.

Oil of Juniper is a mixture of a Terpene with a $C_{15}H_{24}$ —hydrocarbon.

Oil of Cubebs is almost entirely $C_{15}H_{24}$ —hydrocarbons, with a little Terpene.

Oils of Lemon, Nutmeg, and Rosemary are Terpenes with a little oxygenated compound; the second contains also *Cymene* $C_{10}H_{14}$.

Oils of Caraway, Cajeput, Chamomile, Lavender, Spearmint, Peppermint, and Rue are mixtures of Terpenes with oxygenated compounds.

Oils of Dill, Anise, Cinnamon, and Pimento contain hardly any Hydrocarbon.

Oil of Coriander contains none; but only oxygenated compounds.

Oil of Cloves is almost entirely oxygenated, but contains a little $C_{15}H_{24}$ —hydrocarbon.

(Oxygenated Compounds.)

Camphor possesses the composition $C_{10}H_{16}O$. It is usually described as a concrete volatile oil, and for pharmaceutical purposes may be so considered. Its precise chemical position is not determined.

Oils of Lavender and Rosemary in cold weather deposit a solid substance of the same composition as, and supposed to be identical with, Camphor. It is also said to be contained in Oil of Cloves.

A liquid body of similar composition enters into the constitution of Oils of Spearmint, Caraway, Dill, and Nutmeg, and is variously known as *Menthol*, *Carvol*, and *Myristicol*. It is regarded as an alcohol.

The term *Menthol* is also applied to "*Peppermint Camphor*" (*Menthylic Alcohol*, $C_{10}H_{20}O$), which is sometimes deposited by Oil of Peppermint in the cold. It is obtained abundantly from the Oil of *Mentha Javanica* (see "Non-Official Drugs").

Amongst other oxygenated constituents that have been investigated, may be mentioned *Anethol*, $C_{10}H_{12}O$, which forms the bulk of Oil of Anise; *Cinnamic Aldehyde*, C_9H_8O , which largely enters into Oil of Cinnamon; *Angelical Aldehyde*, which is found in Oil of Chamomile; and *Euclic Aldehyde*, $C_{11}H_{22}O$, in Oil of Rue.

Oil of Mustard differs widely from all the rest, being *Allylic Sulphocyanide*, C_3H_5NS . It is not a naturally existing oil, but in a sense an artificial product (see p. 40).

All the volatile oils are soluble in alcohol and fixed oils, and to a slight extent in water.

They can be distilled without decomposition, and volatilize at ordinary temperatures, their vapours being powerfully odorous.

They are obtained by distillation. (See p. 6.)

This group may be divided into two divisions; the first containing thirteen oils used little more than as flavouring agents, the second containing ten which possess active medicinal properties.

(1st division, unimportant.)

- Oleum Anethi.** *Oil of Dill.* From the fruit (p. 34) of *Anethum Graveolens* (Umbelliferæ).
- Ol. Anisi.** *Oil of Anise.* From the fruit of *Pimpinella Anisum*, *Anise*, (Umbelliferæ), and *Illicium Anisatum*, *Star Anise*, (Magnoliaceæ).
- Ol. Carui.** *Oil of Caraway.* From the fruit (p. 34) of *Carum Carui* (Umbelliferæ).
- Ol. Caryophylli.** *Oil of Cloves.* From the sun-dried buds (p. 21) of *Caryophyllus Aromaticus* (Myrtaceæ).
- Ol. Cinnamomi.** *Oil of Cinnamon.* From the inner bark (p. 23) of the shoots of *Cinnamomum Zeylanicum* (Myrtaceæ).
- Ol. Coriandri.** *Oil of Coriander.* From the dried fruit (p. 34) of *Coriandrum Sativum* (Umbelliferæ).
- Ol. Lavandulæ.** *Oil of Lavender.* From the flowers of *Lavandula Vera* (Labiataæ).
- Ol. Limonis.** *Oil of Lemons.* From the fresh peel (p. 34) of the fruit of *Citrus Limonum* (Aurantiaceæ).
- Ol. Menthæ Piperitæ.** *Oil of Peppermint.* From the fresh-flowering plant of *Mentha Piperita* (Labiataæ).
- Ol. Menthæ Viridis.** *Oil of Spearmint.* From the fresh-flowering plant of *Mentha Viridis* (Labiataæ).
- Ol. Myristicæ.** *Oil of Nutmeg.* From the kernel of the fruit (p. 40) of *Myristica Officinalis* (Myristicaceæ).
- Ol. Pimentæ.** *Oil of Pimento.* From the unripe fruit (p. 35) of *Eugenia Pimenta* (Myrtaceæ).
- Ol. Rosmarini.** *Oil of Rosemary.* From the flowering tops of *Rosmarinus Officinalis* (Labiataæ).

OFFICIAL PREPARATIONS.

Aqua Menthæ Piperitæ.**Aqua Menthæ Viridis.****Essentia Anisi.****Essentia Menthæ Piperitæ.**

Tinct. Lavandulæ Co. A weak tincture of Cinuamon, Nutmeg, and Red Saudal Wood, with a little oil of Lavender, and a trace of Oil of Rosemary.

Spiritus Lavandulæ.**Spiritus Menthæ Pip.****Spiritus Myristicæ.****Spiritus Rosmarini.**

} All 1 volume of oil in 50.

* * * The other flavouring waters, Aq. Anethi, Aq. Aurant. Floris, Aq. Carui, Aq. Cinnamomi, Aq. Fœuiculi, Aq. Pimentæ, Aq. Rosæ, Aq. Sambuci, are also preparations of volatile oils, but are ordered to be distilled directly from the raw drugs, and not from the oils. (See p. 6.)

Ol. Anthemidis forms part of Extractum Anthemidis.

(2nd division, important.)

Camphora. *Camphor.* Sublimed from the wood of *Camphora Officinarum* (Lauracæ).

Solid and crystalline.

Oleum Anthemidis. *Oil of Chamomile.* From the flowers (p. 20) of *Anthemis Nobilis* (Compositæ).

Ol. Cajuputi. *Oil of Cajeput.* From the leaves of *Melaleuca Minor* (Myrtacæ).

Ol. Copaibæ. *Oil of Copaiva.* From Copaiva. (See "Oleo-resins," Sixteenth Group.)

Ol. Cubebæ. *Oil of Cubebs.* From the unripe fruit (p. 36) of *Cubeba Officinalis* (Piperacæ).

Ol. Juniperi. *Oil of Juniper.* From the unripe fruit of *Juniperus Communis* (Coniferæ).

Ol. Rutæ. *Oil of Rue.* From the leaves and unripe fruit of *Ruta Graveolens* (Rutacæ).

- Ol. Sabinæ.** *Oil of Savin.* From the fresh tops (p. 18) of *Juniperus Sabina* (Coniferæ).
- Ol. Sinapis.** *Oil of Mustard.* From the seeds (p. 40) of *Sinapis Nigra* (Cruciferæ), after the expression of the fixed oil.
Formed in the process of distillation by the action of Myrosin on Myronic Acid (see p. 40).
- Ol. Terebinthinæ.** *Oil of Turpentine.* From the Oleo-Resin (*Turpentine*) exuding from the cut trunks of *Pinus Palustris*, *Tæda*, and *Pinaster*. (See "**Resina**" in Fourteenth Group.)

OFFICINAL PREPARATIONS.

Aq. Camphoræ. Water saturated with Camphor.

Tinctura Camphoræ Co. A preparation of Opium. (See Twentieth Group.)

Spiritus Camphoræ. $5\frac{1}{2}$ gr. in the drachm.

Spiritus Cajuputi. 1 part of oil in 50.

Spiritus Juniperi. Do.

Linimentum Camphoræ. A strong solution in Olive Oil.

Linimentum Camphoræ Co. A solution of Camphor and Ammonia in spirit, scented with Oil of Lavender.

Linimentum Sinapis Co. A solution of Oil of Mustard, Castor Oil, Camphor, and Ethereal Extract of Mezereon, in Rectified Spirit.

Linimentum Terebinthinæ. Camphor and Soft Soap dissolved in Oil of Turpentine.

Linimentum Tereb. Aceticum. Equal parts of Oil of Turpentine, Acetic Acid, and Camphor Liniment.

Unguentum Terebinthinæ. Half oil of Turpentine.

Confectio Terebinthinæ. With Honey and Liquorice-root; 1 in 4.

Enema Terebinthinæ. An ounce of the oil in 15 of Starch-Mucilage.

FIFTEENTH GROUP :—RESINS. 7.

All important except Mastic, which is rarely used. Rosin is chiefly important as a vehicle.

The composition and chemical nature of resins are as yet incompletely investigated. They are oxygenated compounds; amorphous, solid and brittle at ordinary temperatures, softening and fusing by heat. They possess acid properties, forming soluble salts ("resin soaps") with alkalies, and insoluble salts with some metallic oxides.

They are for the most part soluble in alcohol and ether, insoluble in water. *From solution in alcohol they are precipitated in fine division by water; from solution in alkali by acids.*

The "Resins" obtained from plants are combinations of several resinous substances.

Guaiaci Resina. *Guaiacum Resin.* From the stem of *Guaiacum Officinale* (Zygophyllacæ).

In "tears" or lumps. Exudes naturally, or from artificial incisions in the living trunk; but ordinarily melted out of billets of Guaiac wood (p. 22).

Soluble in spirit and ether, and in solutions of caustic alkalies.

Yields a little extractive matter to water. Discoloured green, then blue and brown, by nitric acid, ozone, chlorine, Ferric and Cupric chlorides; blue by gluten.

Jalapæ Resina. *Jalap Resin.* From **Jalapa** (p. 29).

Extracted by Rectified Spirit. Water is added before distilling off the spirit, to retain and wash away the gummy matters, &c.

Soluble in Rectified Spirit and alkaline solutions, partially in ether.

Mastiche. *Mastic.* From the stem of *Pistachia Lentiscus* (Anacardiaceæ). Exudes from artificial incisions.

Contains a little volatile oil.

90 p.c. is soluble in Rectified Spirit.

Pix Burgundica. *Burgundy Pitch.* From the stem of *Abies Excelsa*, *Spruce Fir.* (Coniferæ.)

The resinous portion of the oleo-resin (Turpentine) from the above tree, separated by melting and straining.

Still contains a little volatile oil, giving it its odour.

Almost entirely soluble in Rectified Spirit, and in twice its weight of Glacial Acetic Acid.

The genuine article is seldom found.

Resina. Rosin. From the oleo-resin (ordinary Turpentine) collected from the stem of *Pinus Palustris*, *Pinus Tæda*, and *Pinus Pinaster*.

The residue left after the distillation of **Ol. Terebinthinæ** (p. 47).

Consists of Pinic Acid, soluble in cold Rectified Spirit.

Sylvic	„	„	warm	„
Pimaric	„	„	boiling	„

All of the formula $C_{20}H_{30}O_2$.

All soluble in ether and volatile oils; uniting by fusion with wax and fixed oils; forming greasy salts (*resin-soaps*) with caustic alkalies.

Contains also a little water and volatile oil.

Podophylli Resina. Podophylline. From **Podophylli Radix** (p. 30).

Procured in fine powder by making a strong liquid extract from the rhizome with Rectified Spirit, and pouring it into water acidulated with Hydrochloric Acid.

Soluble in Rectified Spirit and solutions of Ammonia.

Scammoniæ Resina. Scammony Resin. From **Scammoniæ Radix** (p. 29) or **Scammonium** (see “Gum-Resins,” Nineteenth Group).

Procured in the same way as Jalap Resin.

Soluble in alcohol and ether.

OFFICINAL PREPARATIONS.

Tinctura Guaiaci Ammoniata. A solution of the Resin in Sp. Amm. Aromaticus.

Mistura Guaiaci. The resin suspended (an emulsion) by Gum Arabic in sweetened Cinnamon water.

Mistura Scammonii. The resin similarly suspended in milk.

Unguentum Resinæ. See “Synopsis of Comp. Prep.”

Emplastrum Resinæ.	„	„
---------------------------	---	---

Emplastrum Picis.	„	„
--------------------------	---	---

Pilula Scammonii Co. $1\frac{1}{2}$ grain each of Scammony Resin and Jalap Resin in 5 grains, with Ginger; on a basis of curd soap.

SIXTEENTH GROUP :—OLEO-RESINS. 4.

Copaiba is the only one of any great importance.

Olco-resins are exudations formed by a combination of resins with volatile oils.

Copaiba. *Copaiva*. From the trunk of *Copaifera Multijuga* (Leguminosæ).
Exudes from artificial incisions.

Contains 40 p.c. of **Ol. Copaibæ** (p. 46); the rest chiefly the resin
Copaivic Acid $C_{20}H_{30}O_2$.
Soluble in Rectified Spirit, ether, fixed and volatile oils.

Elemi. Said to be from *Canarium Communc* (Amyridaceæ). Source not exactly known.

Neither the oily nor the resinous portion have been accurately examined.
Almost entirely soluble in Rectified Spirit.

Terebinthina Canadensis. *Canada Balsam*. From the stem of
Abies Balsamea (Coniferæ).

Exudes from artificial incisions.

Nearly the same as ordinary Turpentine, containing *resin* and an *oil* resembling **Ol. Terebinthinæ** (p. 47).

Solidifies with Magnesia, resinous salts of Magnesia being formed, which absorb the volatile oil.

On exposure to air, the oil passes off, leaving the resin as a transparent varnish.

Thus Americanum. *Frankincense*. From *Pinus Palustris* and *Tæda* (Coniferæ).

Concrete American Turpentine; a natural exudation. Solid.

Much the same composition as ordinary Turpentine, which is collected in artificial hollows made in the trunk, and is semi-fluid, as it contains more oil.

OFFICIAL PREPARATIONS.

Unguentum Elemi.

* * * *Canada Balsam* enters into *Charta Epispastica* and *Collodium Flexile*; *Frankincense* is used in making Pitch Plaster, to improve its colour and consistence.

SEVENTEENTH GROUP :—BALSAMS. 4.

Balsams are resinous or oleo-resinous exudations, containing one or other of the volatile Aromatic Acids, *Benzoic*, $C_7 H_6 O_2$, or *Cinnamic*, $C_9 H_8 O_2$.

Of the four officinal Balsams, Benzoin contains the former, the others the latter, acid.

In Benzoin, the remaining constituents are resinous simply; in the others they comprise both resinous and oily matters, and a neutral principle, *Styracin*.

Tolu and Tr. Benzoini Co. are in frequent use.

Balsamum Peruvianum. *Peruvian Balsam.* From the trunk of *Myroxylon Pereiræ* (Leguminosæ).

Exudes after removal of the Bark.

Soluble in Rectified Spirit.

Balsamum Tolutanum. *Tolu.* From the trunk of *Myroxylon Toluifera* (Leguminosæ).

Exudes after incising the bark. Soft when fresh; hardens on keeping, by decomposition of the oil into resin and Cinnamic Acid.

Soluble in Rectified Spirit.

Yields Cinnamic Acid to water.

Benzoinum. *Benzoin.* From the trunk of *Styrax Benzoin* (Styracaceæ).

Exudes after incising the bark.

See **Acidum Benzoicum** (Twenty-first Group).

Soluble in Rectified Spirit and solutions of caustic alkalies.

Styrax Præparatus. *Purified Storax.* From the inner bark of *Liquidambar Orientale* (Liquidambaraceæ).

Extracted by hot water and pressure; purified by melting or solution in Rectified Spirit, and straining.

Soluble in Rectified Spirit.

The volatile oil, *Styrol*, $C_8 H_8$, has aromatic properties.

OFFICINAL PREPARATIONS.

Tinctura Tolutana. Prepared from Tolu.

Tinctura Benzoini Co. *Friar's Balsam.* A tincture of Benzoin, Storax, Socotrine Aloes, and Tolu.

Syrupus Tolutanus. Prepared from Tolu; practically a Syrup of Cinnamic Acid (p. 7).

* * Storax enters only into Tr. Benzoini Co.

For Adeps Benzoatus, see p. 13.

EIGHTEENTH GROUP:—GUMS. 2.

Important as vehicles.

Gums are carbo-hydrates, *i.e.* substances whose molecules are formed of six, or some simple multiple of six, atoms of Carbon, with Hydrogen and Oxygen in the proportions of H_2 , O.

Acaciæ Gummi. *Gum Arabic.* From the stems of undetermined species of *Acacia* (Leguminosæ).

A natural exudation.

Consists almost entirely of the carbohydrate *Arabin*, $C_{12} H_{22} O_{11}$.

Soluble in equal parts of water, hot or cold; precipitated from solution by alcohol in excess. Insoluble in spirit, ether, or oils.

The solution forms an opaque white jelly with Plumbic Subacetate.

Tragacantha. *Tragacanth.* From the stem of *Astragalus Verus* (Leguminosæ).

Exudes naturally, or from artificial incisions in the bark.

Consists of *Arabin*, and its modification *Metarabin*, which is insoluble in water, but absorbs it and swells up.

Metarabin is soluble in Rectified Spirit.

OFFICINAL PREPARATIONS.

Mucilago Acaciæ. 4 oz. of gum in 6 fl. oz. of water. A thick solution, used to suspend insoluble powders, oils, etc., in mixtures.

"Oils require about three-fourths their weight, balsams and spermaceti equal parts, resins two parts, and musk 5 times its weight" (Pereira). The preparation develops Acetic Acid on being kept.

Mucilago Tragacanthæ. 60 grains of *Tragacanth* in half a pint of water. A similar preparation.

Pulvis Tragacanthæ Co. Equal parts of *Tragacanth*, Gum, and Starch, with three parts of sugar.

A vehicle for heavy powders.

NINETEENTH GROUP :—GUM-RESINS. 6.

All important drugs except Galbanum.

Gum-resins consist of Resins and Gums, with, in most cases, a volatile oil.

The resins generally form 65-80 p.c. of the whole, the gum 20-25 p.c., the oil about 4 p.c. In Scammony the resin is from 80 to 90 p.c., the gum from 6 to 20 p.c.

Of the pharmacopœial gum-resins, Gamboge and Scammony contain no oil.

They form perfect emulsions when rubbed with water, the resin and oil being suspended by the gum.

Ammoniacum. *Ammoniac.* From all parts of *Dorema Ammoniacum* (Umbelliferae).

Oozes from the punctures of insects as a milky juice, which hardens in the air. In two forms, "tears" and lumps.

Assafoetida. From the root of *Narthex Assafoetida* (Umbelliferae).

Oozes as a milky juice, which hardens in the sun, from incisions in the top of the root, producing "tears"; or from the surface of the root, cut off transversely, producing lumps.

The volatile oil contains sulphur, being *Allylic Sulphide* ($C_3H_5)_2S$.

Cambogia. *Gamboge.* From *Garcinia Morella*, var. *Pedicellata* (Guttiferae).

Exudes from the broken twigs or pedicels; generally run into hollow bamboo sticks to harden.

The resin is *Gambogic Acid*.

Galbanum. From *Ferula Galbaniflua* (Umbelliferae).

Mode of collection not known. In "tears" or lumps.

Myrrha. *Myrrh.* From the stem of *Balsamodendron Myrrha* (Amyridaceae).

Exudes naturally from the bark. In "tears" or lumps.

The resin and oil are soluble in Rectified Spirit.

Scammonium. *Scammony.* From the root of *Convolvulus Scammonia* (Convolvulaceae). Exudes from artificial excisions.

The resin is **Scammoniae Resina** (p. 49).

Scammoniae Radix is officinal (p. 30).

OFFICINAL PREPARATIONS.

Pulvis Scammonii Co. Scammony, with three-fourths its amount of Jalap, one-fourth of Ginger.

Confectio Scammonii. Scammony and Ginger, with Syrup and Honey, flavoured with Oils of Caraway and Cloves.

Pil. Assafœtidæ Co. Equal parts of Assafœtida, Galbanum, and Myrrh. Made up with Treacle.

Pil. Aloes et Assafœtidæ, see next page.

Pil. Cambogiæ Co. Equal parts of Gamboge, Barbadoes Aloes, and Pulv. Cinnamomi Co., with 2 parts of hard soap and a little syrup.

Pil. Aloes et Myrrhæ, see next page.

Tinctura Myrrhæ.

Tinctura Assafœtidæ. 1 in 8.

Spiritus Ammoniae Fœtidus. A weaker preparation of Assafœtida, 1 in 13, a tenth part of the fluid being Liq. Ammon. Fortior.

Mistura Ammoniaci. An emulsion made by rubbing the gum-resin with water, strained.

Enema Assafœtidæ. An emulsion formed by rubbing 30 gr. of the gum-resin in 4 oz. of water.

Emplastrum Galbani. Contains an equal amount of Ammoniacum.

Emplastrum Ammoniaci cum Hydrargyro. (See "Mercury", in Section III).

TWENTIETH GROUP:—CONCRETE JUICES. 8. EXCRESCENCE. 1.

Manna is now seldom used ; Gutta-percha is used as a vehicle ;
the others are highly important.

a

Aloe Barbadensis. *Barbadoes Aloes.* From the fleshy leaf of Aloe Vulgaris (Liliaceæ).

The juice flowing from the cut leaves, concentrated by boiling, and solidified. Inferior kinds are extracted from the leaves by pressure or boiling.

Contains a crystalline principle *Aloïn*, $C_{16}H_{18}O_7$, orange yellow, soluble in Rect. Spirit and boiling water, slightly in cold water ; *Aloetic Acid*, which strikes olive-brown with ferric salts ; a *resin*, which is soluble in boiling water, as well as in spirit ; and about 60 p.c. of *bitter matters*, not yet accurately examined.

Aloe Socotrina. *Socotrine Aloes.* From the leaves of Aloe Perryi. The juice from the cut leaves dried in the sun.

Composition similar.

OFFICINAL PREPARATIONS.

Extractum Aloes Barbadensis.

Extractum Aloes Socotrinæ.

Decoctum Aloes Co. A decoction from the Extract of Socotrine Aloes, the Extract of Liquorice, and Myrrh, with Potassic Carbonate ; coloured with Saffron, and flavoured with Tr. Cardam. Co.

Tinctura Aloes. A Tincture of Socotrine Aloes and Extract of Liquorice.

Vinum Aloes. From Socotrine Aloes, Ginger, and Cardamoms.

Pilula Aloes Barbadensis. 2 gr. in 5, with hard soap and Confection of Roses, scented with Oil of Caraway.

Pilula Aloes et Ferri. $\frac{3}{4}$ gr. of Ferri Sulphas and 1 gr. of Barbadoes Aloes in 5 ; with Pulv. Cinn. Co. and Confection of Roses.

Pilula Aloes et Assafœtidæ. $1\frac{1}{4}$ gr. of Socotrine Aloes and of Assafœtida in 5 ; with Hard Soap and Confection of Roses.

Pilula Aloes et Myrrhæ. $1\frac{3}{4}$ gr. of Socotrine Aloes, and half as much Myrrh, in 5 ; with Saffron and Confection of Roses.

Pilula Aloes Socotrinæ. Similar to Pil. Aloes Barb. ; scented with Oil of Nutmeg.

Enema Aloes. 40 gr. of either variety rubbed with 15 of Potassic Carbonate in half a pint of Mucilage of Starch.

β

Catechu Pallidum. *Pale Catechu.* From the leaves and young shoots of *Uncaria Gambir* (Rubiaceæ).

An extract made in Singapore and the Eastern Archipelago by boiling the leaves and shoots and evaporating to dryness.

Contains about 49 p.c. of a modification of *Tannic Acid*, which gives a green precipitate with Ferric salts, and becomes insoluble and dark red on exposure to air and moisture; and a crystalline principle, *Catechin*, soluble in boiling water, very slightly in cold.

Galla. *Galls.* From *Quercus Tinctoria*, *Dyer's Oak* (Cupuliferæ).

Excrescences on the small twigs, formed around the ova of the insect *Diplopis Gallæ Tinctoriæ*, which it deposits in a puncture made in the bark.

Parenchymatous in structure, with a few vascular bundles.

Two kinds, "blue" and "white", the latter perforated by the aperture of the insect's escape.

Yields 35 p.c. of *Tannic Acid*, 5 p.c. of *Gallic Acid* (see Tannic and Gallic Acids in Section III).

Kino. From the trunk of *Pterocarpus Marsupium* (Leguminosæ).

Similar composition to Catechu, but contains less matter insoluble in cold water.

OFFICINAL PREPARATIONS.

Pulvis Catechu Co. Catechu, Kino, Rhatany, Cinnamon, and Nutmeg.

Pulvis Kino Co. Kino, with a little Cinnamon and 1 gr. of Opium in the scruple.

Infusum Catechu.

Tinctura Catechu, flavoured with Cinnamon.

Tinctura Gallæ.

Tinctura Kino.

Trochisci Catechu. 1 gr. in each.

Unguentum Gallæ.

Unguentum Gallæ cum Opio. 4 gr. of Opium in the drachm.

γ

Manna. From the stem of *Fraxinus Ornus* and *Rotundifolia* (Oleaceæ).

Exudes from artificial incisions.

Contains 60-80 p.c. of *Mannite*, or Manna-Sugar, $C_6 H_{14} O_6$, with a little *resin*, and some *bitter extractive matter*, in which the active properties probably lie.

Elaterium. From the fruit of *Ecbalium Officinatum* (Cucurbitaceæ).

A sediment from the juice surrounding the seeds (see **Ecbalii Fructus**, p. 37) brought into the market directly, though the Pharmacopœia prescribes a mode of preparing it.

Contains 20 to 30 p.c. of *Elaterin* or *Momordicin* $C_{10} H_{14} O_2$, and some *resin*.

OFFICIAL PREPARATION.

Pulvis Elaterii Co. 1 gr. in 10, with Sugar of Milk.

δ

Gutta-percha. From *Isonandra Gutta* (Sapotaceæ).

Collected from the felled trees. Coagulates spontaneously in air. Sliced and cleansed in England.

80 p.c. of the hydrocarbon $C_{20} H_{32}$, a milk white solid; the rest resins, formed by its oxidation.

Soluble in chloroform and Oil of Turpentine, sparingly in absolute alcohol; unaffected by dilute acetic, hydrochloric or hydrofluoric acids, or alkaline solutions.

Softens with heat.

OFFICIAL PREPARATION.

Liquor Gutta-percha. A solution in Chloroform, purified from the insoluble matters that contaminate commercial Gutta, by shaking with Carbonate of Lead, which carries them to the bottom.

Opium. From the unripe capsules of *Papaver Somniferum* (*Papaveracæ*).

Exudes from artificial incisions and inspissates spontaneously, forming "tears", which are collected and pressed into masses.

The Pharmacopœia prescribes Opium obtained in Asia Minor only. Of this there are two kinds.

1. *Smyrna (Turkey or Levant) Opium*.—Irregular round or flattened lumps, from $\frac{1}{4}$ lb. to 2 lbs. in weight; usually covered with the fruits of a species of rumex. On fracture, they are seen to be made up of agglutinated "tears".
2. *Constantinople Opium*.—Either large flattened cakes, or small lenticular masses, from $\frac{1}{4}$ lb. to $\frac{1}{2}$ lb. in weight, enclosed in a poppy leaf, and marked with its midrib.

No. 2 is unequal in quality, and usually contains less morphia than No. 1. Its odour is less powerful.

Egyptian, Persian, English, and East Indian Opium are not officinal, and are usually inferior to the Asia Minor Opium.

From Opium have been isolated some fourteen alkaloids, of which the most important are *Morphia*, *Codeia*, *Papaverin*, *Cryptopia*, *Thebaia*, and *Narcotin*: two neutral crystalline bodies, *Narceia* and *Meconin*; and two acids:—*Meconic* $C_7 H_4 O_7$, and *Thebolactic* (probably the same as *lactic*).

Resins, gums, and extractive matters, with some *caoutchouc, volatile oil, fatty matters and salts* form the basis.

The best Opium should yield about 12 p.e. of *Morphia*, 8 p.e. of *Meconic Acid*, and the same of *Narcotin*. None of the other definite principles above mentioned form as much as 1 per cent.

Morphia is officinal in the forms of the *Acetate* and *Hydrochlorate* (see Twenty-first Group.) For *Codeia* see "non-officinal drugs".

OFFICINAL PREPARATIONS.

Extractum Opii.

Extractum Opii Liquidum (see p. 4).

Tinctura Opii.

Tinctura Opii Ammoniata. A tincture of Opium and Benzoic Acid, made with a menstruum 4 parts Rectified Spirit, 1 part Liq. Ammon. Fort.; flavoured with Oil of Anise, and coloured with Saffron.

Tinctura Camphoræ Co. A weaker Tincture of Opium, Benzoic Acid, and Camphor; made with proof spirit, and flavoured with oil of Anise.

Vinum Opii. Made from the Extract of Opium with Sherry; flavoured with Cinnamon and Cloves

Pulvis Cretæ Aromaticus cum Opio. Pulv. Cretæ Aromat.
(see Sect. III) with 1 grain of Opium in 40.

Pulvis Opii Co. Opium with Black Pepper, Ginger, and Caraway ;
and a little Tragacanth.

Confectio Opii. 192 grains of the above Powder mixed with a fluid
ounce of Syrup.

Pil. Saponis Co. Opium in Castile Soap. 1 grain in 6.

Pil. Plumbi cum Opio. Acetate of Lead and Opium, with Con-
fection of Roses.

Trochisci Opii. Made with the Extract ; flavoured with Tr. of Tolu.

Enema Opii. Half a drachm of Tr. Opii in 2 ounces of Starch Mucilage.

Suppositoria Plumbi Co. Made with Acetate of Lead and Opium.

Linimentum Opii. Equal parts of Tr. Opii and Lin. Saponis Co.

Ung. Gallæ cum Opio. See p. 56.

Emplastrum Opii. On a basis of Resin Plaster.

* * * *Opium is contained in three other Official Preparations. They are printed
in italics in the subjoined table.*

Official Preparations containing Opium. Amount of preparation containing,
or corresponding to, 1 grain of solid Opium.

Extr. Opii	½ gr.
Pulv. Opii Co.	}	10 gr.
<i>Pulv. Ipecac. Co.</i> (p. 32)						
<i>Pulv. Kino Co.</i> (p. 56)	20 gr.
Pulv. Cretæ Arom. c. Op.	}	40 gr.
Confect. Opii.						
Lozenges	Five.
Pil. Saponis Co.	6 gr.
Pil. Pb. c. Opio	8 gr.
<i>Pil. Ipecac. c. Scillā</i> (p. 33)	23 gr.
Ext. Opii Liq.	}	11 min.
Vinum Opii						
Tinct. Opii	14½ min.
Tr. Op. Ammon.	96 min.
Tr. Camph. Co.	½ oz.
Enema Opii	1 oz.
Lin. Opii	½ drm.
Empl. Opii	10 gr.
Ung. Gallæ c. Opio	15 gr.
Suppos. Pb. Co.	One suppository.

TWENTY-FIRST GROUP:—ACTIVE PRINCIPLES OF DEFINITE
CHEMICAL COMPOSITION. 14.

All important.

a. Two Carbohydrates, used as vehicles.

Amylum. *Starch.* $C_6H_{10}O_5 + 2H_2O$.

On exposing wheat-flour in contact with water for a week or two, fermentation takes place, and Acetic Acid is formed, which disaggregates and partly dissolves the gluten. The gluten, sugar, gum, acid, etc., being washed away, starch remains, nearly pure, containing only some mineral salts. It is moulded and dried. Wheat-starch alone is official.

Rendered anhydrous at 230 deg. F. in dry air. Converted into dextrin and sugar at 320 deg. F. in closed vessels.

Insoluble in cold water, alcohol, and ether. Forms a mucilage with water at 195 deg. F.

Converted into dextrin and sugar by prolonged boiling in water, by saliva, malt extract, and mineral acids.

Saccharum Purificatum. *Loaf-sugar.* $C_{12}H_{22}O_{11}$.

Contained in the sap of the *Sugar-cane*. *Saccharum Officinatum* (Graminaceæ), and of the *Beet-root*, *Beta Vulgaris* (Chenopodiaceæ).

The expressed sap of the sugar-cane, being freed from albuminoids by heating with lime and drawing off from the scum, and concentrated by boiling, yields impure crystalline sugar (*raw* or *brown sugar*) and an uncrystallizable modification, *Molasses* or *Golden Syrup*.

The crystalline sugar being redissolved, the solution filtered through cotton, then through animal charcoal, and concentrated, it crystallizes out in a pure form. The mother syrup yields a less pure sugar, and an uncrystallizable modification, known as *Treacle*.

Beet-root sugar is refined at once, the "raw" sugar having so unpleasant a taste as to be uneatable.

Both beet and cane sugar occur commercially in masses or "loaves" of minute crystals. Larger crystals are exceptionally produced.

The Pharmacopœia prescribes "cane sugar", but beet sugar, which has largely supplanted it, is precisely the same compound.

Soluble in less than half its weight of water: in 100 of parts Rectified Spirit.

(**Theriaca.** The Treacle referred to above is included in Section III as a manufactured product.)

OFFICIAL PREPARATIONS.

Syrupus. A strong solution of sugar, containing 6 parts in 7, made by dissolving sugar in half its weight of water.

Mucilago Amyli. A solution (? suspension) formed by boiling 2 drms. of starch in half a pint of water.

Used as a vehicle for enemata, and for stiffening bandages.

Glycerinum Amyli. A translucent jelly, formed by heating starch with 8 parts of glycerine to 240 deg. F.

Intended as an ointment.

β. Ono Acid.

Acidum Benzoicum. *Benzoic Acid.* $C_7 H_6 O_2$.

Sublimed from **Benzoinum** (p. 51).

Silky plates or needles.

Soluble in 4 parts of Rectified Spirit, 12 parts of boiling water, 400 parts of cold water.

Sublimes at 462 deg. F.

γ. Two Neutral Principles.

Digitalinum. *Amorphous Digitalin.*

From **Digitalis Folia** (p. 25), by a method of extraction detailed below.

Yellowish scales or porous masses.

Soluble in Rectified Spirit and in acid solutions, very slightly in water.*

Santoninum. *Santonin.* $C_{15} H_{18} O_3$.

From **Santonica** (p. 21), by a method detailed below.

Flat rhombic prisms, becoming yellow in sunlight.

Freely soluble in chloroform and fixed oils, in 50 parts of Rectified Spirit, very slightly in water.

It acts as an acid in presence of alkalies.

Sodic Santonate is soluble in water.

* The Infusion of *Digitalis* contains, however, sufficient of this powerful drug to possess active properties in moderate doses.

δ. *Nine Alkaloids and Alkaloidal Salts.*

Aconitia. $C_{30}H_{47}NO_7$.

From **Aconiti Radix** (p. 28), by a method detailed below.

Amorphous powder.

Soluble in 150 parts of cold water, 50 of hot, more freely in spirit.

Fusible.

Atropia. *Atropin.* $C_{17}H_{23}NO_3$.

From **Belladonnæ Radix** (p. 28), by a method detailed below.

Acicular crystals.

Soluble in 8 parts of Rect. Spirit, 500 of water.

Atropiæ Sulphas. Produced by neutralizing a solution of Atropin with Sulphuric Acid, and evaporating to dryness.

Amorphous powder.

Soluble in 4 parts of water.

Beberia Sulphas. $C_{35}H_{40}N_2O_6 \cdot H_2SO_4$.

From **Nectandræ Cortex** (p. 24), by a method detailed below.

Translucent scales, dark brown.

Soluble in 80 parts of water, slightly in spirit.

Morphiæ Hydrochloras. $C_{17}H_{19}NO_3 \cdot HCl \cdot 3H_2O$.

From **Opium** (p. 58), by a method detailed below.

Silky acicular prisms.

Soluble in 20 parts of water, 90 of Rectified Spirit.

Morphiæ Acetas. $C_{17}H_{19}NO_3 \cdot C_2H_4O_2$.

From **Opium** (p. 58), by a method detailed below.

Amorphous Powder.

Soluble in 6 parts of water, 100 of Rectified Spirit.

Quiniæ Sulphas. *Quinine.* $(C_{20}H_{24}N_2O_2)_2 \cdot H_2SO_4 \cdot 7H_2O$.

From **Cinchona Flava** (p. 23), and from the bark of *Cinchona Lancifolia*, by a method detailed below.

Feathery crystals.

Soluble in 40 parts of Rectified Spirit, 1000 of water.

With 1 minim of Ac. Sulphuric. Dil., Ac. Phosphor. Dil., or less of Ac. Nitric. Dil. to each grain, it is soluble in 8 parts of water.

Strychnia. *Strychnine.* $C_{21}H_{22}N_2O_2$.

From **Nux Vomica** (p. 40), by a method detailed below.

Prisms or octahedra.

Soluble in 6,000 parts of water, not in cold absolute alcohol.

Veratria. $C_{32}H_{52}N_2O_8$ (*not quite pure*).

From **Sabadilla** (p. 37), by a method detailed below.

Amorphous masses; pale grey.

Soluble in 11 parts of Rectified Spirit; insoluble in cold water.

OFFICINAL PREPARATIONS.

Liquor Atropiæ.

Liquor Atropiæ Sulphatis.

Liquor Strychniæ.

Liquor Morphiæ Acetatis.

Liquor Morphiæ Hydrochloratis.

All contain $\frac{1}{60}$ of a grain in 2 minims, or $\frac{1}{2}$ grain in a drachm.

Liquor Atropiæ Sulphatis, which is intended for application to the eyes, is a solution in plain water.

The menstrua of the other liquors contain alcohol, one-eighth being rectified spirit in the case of Liq. Atropiæ, one-third in the case of the others. This is necessary to procure solution in the first case, to preserve the alkaloids from decomposition in the others.

A little dilute Hydrochloric Acid is added to procure solution of the Strychnia, which enters into a hydrochlorate.

A little dilute Acetic, and dilute Hydrochloric, Acids are added in making Liq. Morph. Acetatis and Liq. Morph. Hydrochloratis respectively, to supply the place of the acid which is sometimes lost in drying the prepared salts. Morphia by itself is insoluble in water.

Injectio Morphiæ Hypodermica. A solution of Acetate of Morphia in water, with a very slight excess of Acetic Acid. Contains 1-12th grain in each minim.

The solution is not made from Morphiæ Acetas, which, as stated above, is liable to partial decomposition. Morphia is freshly precipitated by Ammonia from Morph. Hydrochloras, carefully dissolved in acetic acid, and diluted.

Tinctura Quiniæ. A solution of Quiniæ Sulphas in Tinct. Aurantii, 1 grain in each drachm; strained from Gallate of Quinia (see p. 34).

Tinctura Quiniæ Ammoniata. A solution in proof spirit, with the same amount of Quinia as the plain tincture, and $7\frac{1}{2}$ min. of Liq. Ammoniac in the drachm.

Vinum Quiniæ. A solution in orange wine, acidulated with citric acid; 1 gr. in each fluid ounce.

Pil. Quiniæ. 2 grains in 3, with confection of Hips.

Suppositoria Morphiæ. $\frac{1}{2}$ gr. of the Hydrochlorate in each suppository.

Suppositoria Morphiæ cum Sapone. The same amount of Morphia.

Trochisci Morphiae. 1-36th gr. of the Hydrochlorate in each :
flavoured with Tolu.

Trochisci Morphiae et Ipecacuanhæ. The same, with 1-12th
gr. of Ipecacuanha.

Unguentum Aconitiæ.	} With 8 gr. of the Alkaloid in the ounce.
Unguentum Atropiæ.	
Unguentum Veratriæ.	

* * Quinia is also officinal in the form of **Ferri et Quiniæ Citras.**
See Sect. III.

The Pharmacopœia prescribes or describes the following processes for the extraction of the Alkaloids and neutral principles.

Aconitia.

- 1.—Powdered ACONITE ROOT (p. 28).
 An extract made with Rectified Spirit.
 The extract exhausted by boiling distilled water.
An infusion containing salts of Aconitia.
- 2.—The Infusion filtered.
 Liquor Ammoniaë added in slight excess, and the whole heated over a water bath.
Impure Aconitia precipitated.
- 3.—The precipitate collected, dried, and powdered.
 An extract made of it by means of ether.
Contains little but pure Aconitia.
 The extract dissolved in warm water, acidulated with sulphuric acid.
A solution of Sulphate of Aconitia.
- 4.—Liquor Ammoniaë, diluted with four times its bulk of water, added cautiously.
Aconitia precipitated.
 Washed and dried.

Atropia.

- 1.—Powdered BELLADONNA ROOT (p. 28).
 Exhausted with Rectified Spirit.
A Tincture containing salts of Atropia, with colouring and resinous matters etc.
- 2.—Lime added, and the tincture shaken.
Colouring matters precipitated.
 The solution filtered.
 The excess of Lime neutralised by Dilute Sulphuric Acid in slight excess.
The tincture now contains salts of Atropia, resins, free Sulphuric Acid, and Calcic Sulphate.
- 3.—Three-fourths of the alcohol distilled off (to avoid waste).
 Water added, and the solution gently heated, till the rest of the alcohol has passed off.
A watery solution of the above substances.
 A solution of Potassic Carbonate added nearly to neutrality.
 Allowed to stand six hours.
Resins precipitated.
 The same added to alkalinity.
Atropia precipitated.

- 4.—The liquid and precipitate shaken up with chloroform.
A chloroformic tincture of Atropia. *A solution of Calcic and Potassic Salts.*
- 5.—The Chloroformic Tincture allowed to subside, and drawn off.
 The chloroform distilled off.
 The residue of Atropia re-dissolved in Rectified Spirit.
 Digested with Purified Animal Charcoal.
Decolorized.
 The solution filtered.
 On evaporation, **Atropiæ Sulphas** crystallizes out.

Atropiæ Sulphas is prepared by diffusing Atropia in water, dissolving it in dilute sulphuric acid, and evaporating to dryness at 100 deg. F.

Beberiæ Sulphas.

- 1.—Powdered BEBEERU BARK (p. 24).
 Exhausted with diluted Sulphuric Acid (1 in 320).
A solution of Sulphate of Berberia, colouring matters, etc.
- 2.—Concentrated, and cooled.
 Milk of Lime added, but *not* to neutrality.
 Allowed to stand for two hours.
Colouring matters precipitated.
- 3.—The solution filtered, the precipitate being washed.
 Liquor Ammoniac added to alkalinity.
Berberia precipitated in an impure state.
- 4.—The precipitate collected, washed, squeezed, dried, and powdered
 Exhausted by boiling in Rectified Spirit.
A tincture of nearly pure Berberia.
- 5.—Water added and the spirit distilled off (to avoid waste).
A watery solution of nearly pure Berberia.
 Dilute Sulphuric Acid added to slight acidity.
A solution of nearly pure Sulphate of Berberia.
- 6.—Evaporated to dryness, and powdered.
 Dissolved in cold water.
 Filtered.
A solution of pure Sulphate of Berberia.
 Evaporated to the consistence of a syrup.
 Dried at 140 deg. F. on glass plates.
Beberiæ Sulphas, in garnet-coloured scales.

Morphiæ Hydrochloras.**1.—OPIUM (p. 58).**

Exhausted by prolonged maceration in cold water.

An infusion containing Morphia and Codeia as Meconates and Sulphates, with resins, extractive matter, etc.

2.—Concentrated till each pint represents a pound of Opium, and strained.

A solution of Calcic Chloride added.

Hydrochlorates of Morphia and Codeia formed, and remain dissolved.

Calcic Meconate and Sulphate formed and precipitated, with the resins.

3.—The whole evaporated till it solidifies on cooling.

The solid mass wrapped in calico and subjected to powerful pressure.

Dark liquid exudes, containing colouring matter, etc., with a small amount of alkaloid.

4.—The mass exhausted with boiling water, and the resulting solution filtered.

A solution of the Hydrochlorates of Morphia and Codeia, with colouring matter, etc.

5.—The solution evaporated till it solidifies on cooling.

Compressed as before.

More dark liquid exudes.

6.—Exhausted again.**7.—Processes Nos. 5 and 6 repeated till a nearly colourless solution is obtained.**

The solution digested with Purified Animal Charcoal for 20 minutes.

Filtered, the charcoal being washed on the filter with boiling water.

A decolorised solution.

Liquor Ammoniaë added.

Codeia remains in solution.

Morphia precipitated.

Collected.

Washed free from Codeia, *i.e.*, till the washings cease to precipitate with a solution of nitrate of silver acidulated by nitric acid.

Diffused in boiling water.

Hydrochloric Acid carefully added till all is dissolved.

A solution of pure Hydrochlorate of Morphia.

On cooling, **Morphiæ Hydrochloras** crystallizes out.

More obtained on concentrating the mother liquor.

Morphiæ Acetas.

Morphia, precipitated from **Morphiæ Hydrochloras** by Liquor Ammonia; collected, washed, and dissolved in dilute Acetic Acid. Evaporated over a water bath till it concretes on cooling. Dried at a gentle heat, and powdered.

Quiniæ Sulphas.

- 1.—YELLOW CINCHONA BARK, or the Bark of Cinch. Lancifolia (pp. 23, 77).
Exhausted with dilute Hydrochloric Acid (1 in 70).
A solution of Hydrochlorates of Quinia and other Alkaloids, with colouring matter, etc.
- 2.—Liquor Sodæ added, with agitation.
Quinia precipitated.
- 3.—The precipitate collected and washed from colouring matter.
Diffused in water.
Heated to 212 deg. Fahr.
Dilute Sulphuric Acid added till all is dissolved.
A solution of Sulphate of Quinia.
- 4.—On concentrating and cooling, **Quiniæ Sulphas** crystallizes out.
Dried.

Strychnia.

- 1.—Nux VOMICA seeds (p. 40) softened by steaming, and ground.
Exhausted by prolonged digestion with Rectified Spirit, diluted with half its bulk of water.
A tincture of Salts of Strychnia and Brucia, with colouring matter, etc.
- 2.—The spirit distilled off (to avoid waste).
The remaining watery solution concentrated, and filtered when cold.
A solution of Plumbic Acetate added, as long as a precipitate is formed.
Colouring matter, etc., precipitated.
- 3.—The solution filtered, the filter being washed.
Concentrated.
Liquor Ammonia added, with stirring.
Allowed to stand for twelve hours.
Strychnia and Brucia precipitated, with other matters.

- 4.—The precipitate collected, washed, and dried.
Exhausted by boiling with Rectified Spirit, as long as any bitter matter is yielded.
Most of the spirit distilled off.
The remainder concentrated and cooled.
Brucia remains in the mother liquor. *Strychnia, and a little Brucia, crystallize out as a white crust.*

- 5.—The crust washed on a filter with Rectified Spirit diluted with half its bulk of water, as long as Brucia is yielded, *i.e.*, as long as the washings redden with Nitric Acid.
The residue dissolved in boiling Rectified Spirit.
On cooling, **Strychnia** *crystallizes out.*
More procured by evaporating the mother liquor.

Veratria.

- 1.—CEVADILLA SEEDS (p. 37) husked and ground.
Exhausted with Rectified Spirit.
A tincture containing Gallate of Veratria, resins etc.
- 2.—Concentrated as long as no deposit forms.
Poured into cold water.
Resins precipitated.
- 3.—Filtered, the precipitate being washed.
Liquor Ammonia added in slight excess.
Gallate of Ammonia remains in solution. *Veratria precipitated, with impurities.*
- 4.—The precipitate collected and washed.
Diffused in water.
Hydrochloric acid added till all is dissolved.
A solution of impure Hydrochlorate of Veratria.
- 5.—Digested with Purified Animal Charcoal at a gentle heat.
Decolorised.
Filtered and cooled.
Liquor Ammonia added.
Purer Veratria precipitated.
- 6.—The precipitate washed till the washings cease to precipitate with a solution of Argentic Nitrate acidulated with Nitric Acid.
Dried at a gentle heat,—**Veratria.**

Digitalinum.

1.—DIGITALIS LEAVES (p. 25).

An extract made with Rectified Spirit, by macerating at 120 deg. F.

Contains Digitaline, colouring and extractive matters.

2.—The extract exhausted with water acidulated with a tenth of Acetic Acid.

A solution containing Digitaline and colouring matter, with free Acetic Acid.

3.—Digested with Animal Charcoal.

Nearly decolorised.

Filtered and diluted.

Nearly neutralised with Liquor Ammoniae.

A solution of Tannic Acid added.

Impure Digitaline precipitated.

4.—The precipitate collected and washed.

Rubbed with Litharge in Rectified Spirit.

Traces of Tannic Acid removed as an insoluble Plumbic Tannate.

Dissolved in Rectified Spirit, at 160 deg. F.

Digested with animal charcoal.

Traces of colouring matter removed.

The solution filtered off.

Evaporated to dryness at 212 deg. F.

Digitaline and a little oily matter remain.

5.—Washed with ether.

The oily matters removed.

Digitalinum *remains in an amorphous state.*

Santoninum.

1.—SANTONICA (p. 21).

Boiled with Lime; the fluid strained and expressed.

A solution containing Santonin and oily matters in combination with Lime.

2.—Concentrated.

While hot, Hydrochloric acid added, with stirring, to acidity.

Allowed to stand five days.

Oil separates on the surface of the fluid.

Santonin precipitated.

The oil skimmed off,
and the fluid decanted.

2.—The precipitate collected, and
washed from acid and colouring
matter.

Washed with diluted Liq. Am-
monia.

Washed again with water till free
from colouring matter.

Expressed and dried.

3.—Re-dissolved in boiling Rectified Spirit.

Digested with Animal Charcoal.

Filtered.

Allowed to stand two days in a cool dark place.

Santoninum *crystallizes out.*

Purified by recrystallization.

Dried in the dark and kept from light.

BOTANICAL SYNOPSIS OF THE NATURAL ORDERS AND PLANTS FURNISHING OFFICINAL DRUGS.

WITH THEIR GEOGRAPHICAL SOURCES, AND THEIR PREPARATIONS.

DICOTYLEDONES.

(THALAMIFLORÆ.)

RANUNCULACEÆ.

1. *Aconitum Napellus*. *Monkshood*. Native.
Aconiti Folia (p. 25). Ext.
Aconiti Radix (p. 28). Tr., Lin.
Aconitia (p. 62). Ung.
2. *Podophyllum Peltatum*.^{*} *American May Apple*. United States.
Podophylli Radix (p. 29).
Podophylli Resina (p. 49).

MAGNOLIACEÆ.

3. *Illicium Anisatum*. *Star Anise*. China.
Oleum Anisi (p. 45. See No. 60). Ess.

MENISPERMACEÆ.

4. *Jateorrhiza Calumba*. *Calumba*. E. Africa.
Calumbæ Radix (p. 28). Ext., Inf., Tr.
5. *Cissampelos Pareira*. *Velvet Leaf*. Brazil.
Pareiræ Radix (p. 29). Ext., Ext. Liq., Dec.

PAPAVERACEÆ.

6. *Papaver Somniferum*. *Garden Poppy*. *a.* Native. *β.* Asia Minor.
a. **Papaveris Capsulæ** (p. 37). Ext., Dec., Syr.
β. **Opium**. See p. 58.
Morphiæ Acetas (p. 62). Liq., Inj. M. Hypod.
Morphiæ Hydrochloras (p. 62). Liq., Troch., Troch. M. c. Ipec.
 Supp.
7. *Papaver Rhœas*. *Red Poppy*. Native.
Rhœados Petala (p. 20). Syr.

^{*} Placed by Bentley in Order *Berberidaceæ*.

CRUCIFERÆ.

8. *Cochlearia Armoracia*. *Horse Radish*. Native.
Armoraciæ Radix (p. 28). Sp. Co.
9. *Sinapis Nigra* and *Alba*. *Mustard*. Native.
Sinapis (p. 40). Cat., Chart. (from *S. Nigra*).
Oleum Sinapis (p. 47). Lin. Co.

POLYGALACEÆ.

10. *Krameria Triandra*. *Rhatany*. Peru and Chili.
Krameriæ Radix (p. 29). Ext., Inf., Tr.
11. *Polygala Senega*. *Senega*. United States.
Senegæ Radix (p. 30). Inf., Tr.

LINACEÆ.

12. *Linum Usitatissimum*. *Flax*. Native.
Lini Semina (p. 40). Inf.
Lini Farina (p. 40). Cat.
Oleum Lini (p. 42).

MALVACEÆ.

13. *Gossypium* :—various species. *Cotton*. N. America ; India.
Gossypium, *Cotton Wool* (p. 40). See *Pyroxylin*, in Sect. III.

BYTTNERIACEÆ.

14. *Theobroma Cacao*.
Oleum Theobromæ, *Cocoa Butter* (p. 43).

AURANTIACEÆ.

15. *Citrus Bigaradia*. *Bitter Orange*. S. Europe.
Aqua Aurantii Floris. From *Citrus Aurantium* as well (p. 20).
Syr., Inf.
Aurantii Cortex (peel. p. 34). Inf., Inf. Co., Tr., Syr.
Aurantii Fructus (p. 34). Tr. Aur. Rec.
16. *Citrus Limonum*. *Lemon*. S. Europe.
Limonis Succus (p. 34). Syr. See *Acidum Citricum* in Sect. III.
Limonis Cortex (p. 34). Tr.
Oleum Limonis (p. 45).
17. *Ægle Marmelos*. *Bael*. Malabar.
Belæ Fructus (p. 36). Ext. Liq.

CANELLACEÆ.

18. *Canella Alba*. W. Indies.
Canellæ Albæ Cortex (p. 23).

GUTTIFERÆ.

19. *Garcinia Morella*. *Gamboge*. Siam.
Cambogia (p. 53). Pil. Co.

VITACEÆ.

20. *Vitis Vinifera*. *Vine*. Spain.
Uvæ, *raisins* (p. 35).
See *Alcohol*, *Pot. Tart. Acid.*, *Ac. Tartaricum*, in Sect. III.

ZYGOPHYLLACEÆ.

- 21.
- Guaiacum Officinale*
- .
- Lignum Vitæ*
- . W. Indies.

Guaiaci Lignum (p. 22).**Guaiaci Resina** (p. 48). Mist., Tr. Amm.

RUTACEÆ.

- 22.
- Barosma Betulina*
- }
-
- „
- Crenulata*
- }
- Bucco*
- . Cape.
-
- „
- Serratifolia*
- }

Buchu Folia (p. 25). Inf., Tr.

- 23.
- Galipea Cusparia*
- .
- Angustura Bark Tree*
- . S. America.

Cuspariæ Cortex (p. 23). Inf.

- 24.
- Ruta Gravcolens*
- .
- Ruc.*
- Native.

Oleum Rutæ (p. 46).

SIMARUBACEÆ.

- 25.
- Picræna Excelsa*
- .
- Quassia*
- . Jamaica.

Quassiæ Lignum (p. 46). Ext., Inf., Tr.

(CALYCIFLORÆ.)

RHAMNACEÆ.

- 26.
- Rhamnus Catharticus*
- .
- Buckthorn*
- . Native.

Rhamni Succus (p. 37). Syr.

ANACARDIACEÆ.

- 27.
- Pistacia Lentiscus*
- .
- Mastic*
- . Levant.

Mastiche (p. 48).

AMYRIDACEÆ.

- 28.
- Canarium Commune*
- .
- Manilla*
- .

Elemi (p. 50). Ung.

- 29.
- Balsamodendron Myrrha*
- .
- Myrrh*
- . Arabia.

Myrrha (p. 53). Tr., Pil. Aloes et M.

LEGUMINOSÆ.

- 30.
- Acacia Vera*
- . E. Africa.

Acaciæ Gummi (p. 52). Muc.

- 31.
- Myroxylon Perciræ*
- . C. America.

Balsamum Peruvianum (p. 51).

- 32.
- Myroxylon Toluifera*
- . C. America.

Balsamum Tolutanum (p. 51). Syr., Tr.

- 33.
- Cassia Fistula*
- .
- Pudding-pipe Tree*
- . E. & W. Indies.

Cassiæ Pulpa (p. 36).

34. *Copaifera* ;—several species. *Copaiva Trees*. C. & S. America.
Copaiba (p. 50).
Oleum Copaibæ (p. 46).
35. *Glycyrrhiza* *Glabra*. *Liquorice*. Native.
Glycyrrhizæ Radix (p. 28). Ext., Ext. Liq., Pulv. Co.
36. *Hæmatoxylon* *Campechianum*. *Logwood*. E. & W. Indies.
Hæmatoxyli Lignum (p. 22). Ext., Dec.
37. *Pterocarpus* *Marsupium*. *Kino Tree*. Malabar.
Kino (p. 56). Pulv. Co., Tr.
38. *Physostigma* *Venenosum*. *Calabar Bean*. W. Africa.
Physostigmatis Faba (p. 40). Ext.
39. *Pterocarpus* *Santalinus*. *Red Sandal Wood Tree*. Ceylon ; Coromandel.
Pterocarpi Lignum (p. 22).
40. *Sarothamnus* *Scoparius*. *Broom*. Native.
Scoparii Cacumina (p. 18). Succ., Dec.
41. *Cassia* *Lanceolata* } *Alexandrian Senna*. Source uncertain.
Cassia *Obovata* }
Senna Alexandrina (p. 26). Inf., Tr., Mist., Syr., Conf.
Cassia *Elongata*. *Tinnevely Senna*. India.
Senna Indica (p. 26). Same preparations.
42. *Tamarindus* *Indica*. *Tamarind*. E. & W. Indies.
Tamarindus (p. 38).
43. *Astragalus* *Verus*. *Milk Vetch*. Asia Minor.
Tragacantha (p. 52). Pulv. Co., Muc.

ROSACEÆ.

44. *Amygdalus* *Communis*
var. Amara. *Bitter Almond*. Morocco.
Amygdala Amara (p. 39).
Oleum Amygdalæ (p. 42).
var. Dulcis. *Sweet or Jordan Almond*. South Europe ;
 Syria.
Amygdala Dulcis (p. 39). Pulv. Co., Mist.
Oleum Amygdalæ (p. 42).
45. *Brayera* *Anthelmintica*. *Kousso*. Abyssinia.
Cusso (p. 19). Inf.
46. *Prunus* *Lauro-cerasus*. *Cherry Laurel*. Native.
Lauro-cerasi Folia (p. 25). Aq.
47. *Prunus* *Domestica*. *Plum-tree*. Europe.
Prunum (p. 37).
48. *Rosa* *Canina*. *Dog Rose*. Native.
Rosæ Caninæ Fructus. *Hips*. (P. 35.) Conf.
49. *Rosa* *Centifolia*. *Cabbage Rose*. Native.
Rosæ Centifoliæ Petala (p. 20). Aq.
50. *Rosa* *Gallica*. *Red Rose*. Native.
Rosæ Gallicæ Petala (p. 20). Inf. Ac., Syr., Conf.

MYRTACEÆ.

51. *Melaleuca Minor*. *Cajeput*. Moluccas.
Oleum Cajeputi (p. 46). Sp.
52. *Caryophyllus Aromaticus*. *Clove*. E. Indian Islands.
Caryophyllum (p. 21). Inf.
Oleum Caryophylli (p. 45).
53. *Punica Granatum*. *Pomegranate*. Mediterranean Coast.
Punicæ Granati Cortex (p. 23). Dec.
54. *Eugenia Pimenta*. *All-spice*. W. Indies.
Pimenta (p. 35). Aq.
Oleum Pimentæ (p. 45).

CUCURBITACEÆ.

55. *Citrullus Colocynthis*. *Colocynth*. India ; Levant.
Colocynthidis Pulpa (p. 36). Ext. Co., Pil. Co., Pil. C. et Hyosc.
56. *Ecballium Officinarium*. *Squirting Cucumber*. S. Europe.
Ecballii Fructus (p. 37).
Elaterium (p. 57). Pulv. Co.

UMBELLIFERÆ.

57. *Dorema Ammoniacum*. Punjaub & Persia.
Ammoniacum (p. 53). Mist., Emp. A. c. Hydrarg.
58. *Anethum Graveolens*. *Dill*. Native.
Anethi Fructus (p. 34). Aq.
Oleum Anethi (p. 45).
59. *Pimpinella Anisum*. *Anise*. China.
Oleum Anisi (p. 45, see No. 3). Ess.
60. *Narthex Assafoetida*.* Punjaub and Persia.
Assafoetida (p. 53). Tr., Pil. Co., En., Sp. Amm. Fœtidus., Pil. Al. et Ass.
61. *Carum Carui*. *Caraway*. Native.
Carui Fructus (p. 35). Aq.
Oleum Carui (p. 45).
62. *Conium Maculatum*. *Hemlock*. Native.
Conii Folia (p. 25). Ext., Pil., Succ., Vap., Cat.
Conii Fructus (p. 36). Tr.
63. *Coriandrum Sativum*. *Coriander*. Native.
Coriandri Fructus (p. 34).
Oleum Coriandri (p. 45).
64. *Fœniculum Dulce*. *Fennel*. Europe.
Fœniculi Fructus (p. 35). Aq.
65. *Ferula Galbaniflua*. India ; Levant.
Galbanum (p. 53). Emp.
66. *Euryangium Sumbul*.† *Musk Root*. Central Asia
Sumbul Radix (p. 30). Tr.

* *Ferula Narthex* (Bentley).† *Ferula Sumbul* (Bentley).

(COROLLIFLORÆ).

CAPRIFOLIACEÆ.

67. *Sambucus Nigra*. *Elder*. Native.
Sambuci Flores (p. 20.) Aq.

RUBIACEÆ.*

68. *Uncaria Gambir*. Singapore.
Catechu Pallidum (p. 56). Pulv. Co., Inf., Tr., Troch.
69. *Cinchona Succirubra*. *Red Cinchona*. Chimborazo.
Cinchonæ Rubræ Cortex (p. 23).
70. *Cinchona Condaminea*. *Pale Cinchona*. Ecuador.
Cinchonæ Pallidæ Cortex (p. 23). Tr. Co.
71. *Cinchona Calisaya*. *Yellow Cinchona*. Bolivia ; Peru.
Cinchonæ Flavæ Cortex (p. 23). Ext. Liq., Inf., Dec., Tr.
Quiniæ Sulphas (p. 62). Pil., Tr., Tr. Amm., Vin.
72. *Cinchona Lancifolia*. New Granada.
Quiniæ Sulphas.
73. *Cephaelis Ipecacuanha*. *Ipecacuanha*. Brazil.
Ipecacuanha (p. 29). Pulv. Co., Pil. Ip. c. Sc., Vin., Troch.

VALERIANACEÆ.

74. *Valeriana Officinalis*. *Valerian*. Native.
Valerianæ Radix (p. 30). Inf., Tr., Tr. Amm.

COMPOSITÆ.

75. *Anthemis Nobilis*. *Chamomile*. Native.
Anthemidis Flores (p. 20). Ext., Inf.
Oleum Anthemidis (p. 46).
76. *Arnica Montana*. Europe.
Arnicæ Radix (p. 28). Tr.
77. *Lactuca Virosa*. *Wild Lettuce*. Native.
Lactuca (p. 18). Ext.
78. *Anacyclus Pyrethrum*. *Pellitory*. Morocco ; Spain ; Levant.
Pyrethri Radix (p. 29). Tr.
79. *Artemisia* :—species uncertain. Russia.
Santonica (p. 21).
Santoninum (p. 60).
80. *Taraxacum Dens Leonis*. *Dandelion*. Native.
Taraxaci Radix (p. 30). Ext., Succ., Dec.

LOBELIACEÆ.

81. *Lobelia Inflata*. *Indian Tobacco*. United States.
Lobelia (p. 18). Tr., Tr. Ætherea

* Nos. 68-73 fall in the separate order *Cinchonaceæ* made by some botanists.

ERICACEÆ.

82. *Arctostaphylos Uva Ursi*. *Bearberry*. Europe.
Uvæ Ursi Folia (p. 26). Inf.

SAPOTACEÆ.

83. *Isonandra Gutta-Percha*. *Gutta Pereha Tree*. Eastern Archipelago.
Gutta-Percha (p. 57). Liq.

STYRACACEÆ.

84. *Styrax Benzoin*. *Benjamin Tree*. Eastern Archipelago.
Benzoinum (p. 51). Tr. Co.
Acidum Benzoicum (p. 61).

OLEACEÆ.

85. *Olea Europæa*. *Olive*. S. Europe.
Oleum Olivæ (p. 42).
 86. *Fraxinus Ornus* and *Rotundifolia*. Sicily.
Manna (p. 57).

ASCLEPIADACEÆ.

87. *Hemidesmus Indicus*. *Indian Sarsaparilla*. India.
Hemidesmi Radix (p. 29). Syr.

LOGANIACEÆ.

88. *Strychnos Nux Vomica*. *Koochla Tree*. India.
Nux Vomica (p. 40). Ext., Tr.
Strychnia (p. 62). Liq.

GENTIANACEÆ.

89. *Ophelia Chirata*. *Chiretta*. India.
Chirata (p. 18). Inf., Tr.
 90. *Gentiana Lutea*. *Gentian*. European Mountain
Gentianæ Radix (p. 28). Ext., Inf. Co., Tr. Co., Mist.

CONVOLVULACEÆ.

91. *Exogonium Purga*. *Jalap*. Mexico.
Jalapa (p. 29). Ext., Pulv. Co., Tr.
Jalapæ Resina (p. 48).
 92. *Convolvulus Scammonia*. *Scammony*. Asia Minor.
Scammoniae Radix (p. 30).
Scammoniae Resina (p. 49). Mist., Pil. Co.
Scammonium (p. 53). Pulv. Co., Conf.

SOLANACEÆ. *Sub-order Solanææ.*

93. *Capsicum Fastigiatum*. Zanzibar.
Capsici Fructus (p. 36). Tr.
94. *Solanum Dulcamara*. *Woody Nightshade*. Native.
Dulcamara (p. 18). Inf.

Sub-order Atropeæ (ATROPACEÆ of some botanists).

95. *Atropa Belladonna*. *Deadly Nightshade*. Native.
Belladonnæ Folia (p. 25). Ext., Succ., Tr., Ung., Emp.
Belladonnæ Radix (p. 28). Lin.
Atropia (p. 62). Liq., Ung.
Atropiæ Sulphas (p. 62). Liq.
96. *Hyoscyamus Niger*. *Henbane*. Native.
Hyoscyami Folia (p. 25). Ext., Succ., Tr.
97. *Datura Stramonium*. *Thorn Apple*. Native.
Stramonii Folia (p. 26).
Stramonii Semina (p. 40). Ext., Tr.
98. *Nicotiana Tabacum*. *Tobacco*. Tropical America.
Tabaci Folia (p. 26). En.

SCROPHULARIACEÆ.

99. *Digitalis Purpurea*. *Foxglove*. Native.
Digitalis Folia (p. 25). Inf., Tr.
Digitalinum (p. 61).

LABIATÆ.

100. *Lavandula Vera*. *Lavender*. Native.
Oleum Lavandulæ (p. 45). Sp., Tr. Co.
101. *Mentha Piperita*. *Peppermint*. Native.
Oleum Menthæ Piperitæ (p. 45). Aq., Ess., Sp.
102. *Mentha Viridis*. *Spearmint*. Native.
Oleum Menthæ Viridis (p. 45). Aq.
103. *Rosmarinus Officinalis*. *Rosemary*. S. Europe.
Oleum Rosmarini (p. 45). Sp.

(MONOCHLAMYDEÆ.)

POLYGONACEÆ.

104. *Rheum*:—species uncertain. *Rhubarb*. Central Asia.
Rhei Radix (p. 29). Ext., Inf., Tr., Vin., Pil. Co., Pulv. Co., Syr

THYMELACEÆ.

105. *Daphne Mezereum*, or *Laureola*. *Spurge Laurel*. Native.
Mezerei Cortex (p. 23). Extr. Ether.

MYRISTICACEÆ.

106. *Myristica Officinalis*. *Nutmeg*. Eastern Archipelago.
Myristica (p. 40).
Oleum Myristicæ (p. 45). Sp.
Oleum Myristicæ Expressum (p. 42).

LAURACEÆ.

107. *Camphora Officinarum*. *Camphor Plant*. China ; Japan.
Camphora (p. 46). Aq., Sp., Tr. Co., Lin., Lin. Co.
 108. *Cinnamomum Zeylanicum*. *Cinnamon*. Ceylon.
Cinnamomi Cortex (p. 23). Pulv. Co., Aq., Tr.
Oleum Cinnamomi (p. 45).
 109. *Nectandra Rodiaei*. *Bebeeru, or Green-heart Tree*. Guiana.
Nectandræ Cortex (p. 24).
Bebericæ Sulphas (p. 62).
 110. *Sassafras Officinale*. N. America.
Sassafras Radix (p. 29).

ARISTOLOCHIACEÆ.

111. *Aristolochia Serpentaria*. *Serpentary*. Virginia.
Serpentariæ Radix (p. 30). Inf., Tr.

EUPHORBIACEÆ.

112. *Croton Eluteria*. *Cascarilla*. Bahamas.
Cascarillæ Cortex (p. 23). Inf., Tr.
 113. *Croton Tiglium*. *Croton-oil Plant*. India.
Oleum Crotonis (p. 42). Lin.
 114. *Rottleria Tinctoria*. India.
Kamala (p. 37).
 115. *Ricinus Communis*. *Castor-oil Plant*. India.
Oleum Ricini (p. 42).

PIPERACEÆ.

116. *Cubeba Officinalis*. *Cubebæ*. Java.
Cubeba (p. 36). Tr.
Oleum Cubebæ (p. 46).
 117. *Artanthe Elongata*. *Matiao*. Peru.
Matica Foliæ (p. 25). Inf.
 118. *Piper Nigrum*. *Black Pepper*. India.
Piper Nigrum (p. 37). Conf.

MORACEÆ.

119. *Morus Nigra*. *Mulberry*. Native.
Mori Succus (p. 34). Syr.
 120. *Ficus Carica*. *Fig Tree*. Asia Minor.
Ficus (p. 37).

CANNABINACEÆ.

121. *Cannabis Sativa*. *Indian Hemp*; *Bhang*. India.
Cannabis Indica (p. 19). Ext., Tr.
122. *Humulus Lupulus*. *Hop*. Native.
Lupulus (p. 19). Ext., Inf., Tr.

ULMACEÆ.

123. *Ulmus Campestris*. *Elm*. Native.
Ulmi Cortex (p. 24). Dec.

CUPULIFERÆ.

124. *Quercus Pedunculata*. *Oak*. Native.
Quercus Cortex (p. 24). Dec.
125. *Quercus Tinctoria*. *Gall Oak*. Asia Minor.
Galla (p. 56). Tr., Ung., Ung. G. c. Opio.

LIQUIDAMBARACEÆ.

126. *Liquidambar Orientale*. Asia Minor.
Styrax Præparatus (p. 51).

(GYMNOSPERMIA.)

CONIFERÆ.

127. *Juniperus Communis*. *Juniper*. N. Europe.
Oleum Juniperi (p. 46). Sp.
128. *Juniperus Sabina*. *Savin*. Native.
Sabinæ Cacumina (p. 18). Tr., Ung.
Oleum Sabinæ (p. 47).
129. *Larix Europæa*. *Larch*. Europe.
Laricis Cortex (p. 23). Tr.
130. *Abies Excelsa*. Switzerland.
Pix Burgundica (p. 48). Empl. Picis.
131. *Abies Balsamea*. *Balm of Gilead Fir*. Canada.
Terebinthina Canadensis (p. 50).
132. *Pinus Palustris*, *Tæda*, *Pinaster*. America; France.
Thus Americanum (p. 50).
Oleum Terebinthinæ (p. 47). Conf., En., Ung., Lin., Lin. Acet.,
 Emp.
Resina (p. 49). Ung., Emp.

MONOCOTYLEDONES.

SMILACEÆ.

133. *Smilax Officinalis*. *Sarsaparilla*. Central America.
Sarsæ Radix (p. 29). Ext. Liq., Dec., Dec. Co.

ZINGIBERACEÆ.

134. *Elettaria Cardamomum*. *Cardamom*. Malabar.
Cardamomum (p. 39). Tr. Co.
135. *Zingiber Officinale*. *Ginger*. E. & W. Indies.
Zingiber (p. 31). Tr., Tr. Fort., Syr.

IRIDACEÆ.

136. *Crocus Sativus*. *Saffron*. S. Europe.
Crocus (p. 20). Tr.

LILIACEÆ.

137. *Aloe Vulgaris*. *Barbadoes Aloes*. Barbadoes.
Aloe Barbadensis (p. 55). Ext., Pil., Pil. A. et Fer., En.
138. *Aloe Perryi*. Socotra.
Aloe Socotrina (p. 55). Ext., Dec. Co., Tr., Vin., Pil., Pil. A. et Ass.,
 Pil. A. et Myr., En.
139. *Urginea Scilla*. *Squill*. S. Europe.
Scilla (p. 30). Tr., Acet., Oxy., Syr., Pil. Co., Pil. Ipec. c. Sc.

MELANTHACEÆ.

140. *Colchicum Autumnale*. *Meadow Saffron*. Native.
Colchici Cormus (p. 28). Ext., Ext. Acet., Vin.
Colchici Semina (p. 39). Tr.
141. *Asagrea Officinalis*. *Cevadilla*. Mexico.
Sabadilla (p. 37).
Veratria (p. 62). Ung.
142. *Veratrum Viride*. *Green Hellebore*. N. America.
Veratri Viridis Radix (p. 30). Tr.

PALMACEÆ.

143. *Areca Catechu*. *Betel Nut*. India.
Areca (p. 39).

GRAMINACEÆ.

144. *Triticum Vulgare*. *Wheat*.
Farina Tritici (p. 39). Cat. Fermenti.
Amylum (p. 60). Muc., Glys.
145. *Hordeum Distichon*. *Barley*. Native.
Hordeum Decorticatum (p. 39). Dec.
146. *Saccharum Officinarum*. *Sugar Cane*. W. Indies.
Saccharum Purificatum (p. 60). Syrupus.
(Theriaca.)

FILICES.

147. *Aspidium Filix Mas*.^{*} *Male Fern*. Native.
Filix Mas (p. 28). Ext. Liq.

FUNGI.

148. *Claviceps Purpurea*. *Ergot of Rye*. Native.
Ergota (p. 17). Ext. Liq., Inf., Tr.
149. *Torula Cerevisiæ*. *Yeast Plant*. Native.
Cerevisiæ Fermentum (p. 17). Cat.

LICHENES.

150. *Cetraria Islandica*. *Iceland Moss*. Iceland.
Cetraria (p. 17). Dec.

* *Lastrea Filix Mas*. (*Bentley*).

NON-OFFICINAL DRUGS FROM THE VEGETABLE KINGDOM.

The following are those that chiefly demand notice.

PLANTS OR PARTS OF PLANTS.

- ACTÆA RACEMOSA.** The root of *Actæa*, or *Cimicifuga*, *Racemosa*, *Black Snake Root* (Ranunculaceæ). N. America.
An active principle, *Cimicifugin*, has been isolated.
- CHIMAPHILA.** The dried leaves of *Chimaphila Umbellata*, *Pipsissewa* or *Winter Green* (Pyrolaceæ). N. America.
Contains a crystalline neutral principle, *Chimaphilin*.
- CONVALLARIA.** The plant *Convallaria Majalis*, *Lily of the Valley* (Liliaceæ).
The active principle is a glucoside, *Convallamarin*.
- DATURA TATULA.** The leaves of *Datura Tatula* (Solanaceæ). Native.
Probably the same constituents as *Stramony* (p. 26).
- DUBOISIA.** The plant *Duboisia Myoporoides* (Solanaceæ). New S. Wales and Queensland.
Contains an alkaloid, *Duboisin*, probably identical with *Atropin* (p. 62).
- EUCALYPTUS.** The leaves of *Eucalyptus Globulus*, *Blue Gum Tree* (Myrtaceæ) Tasmania, Australia, Algeria.
Studded with glands containing *Eucalyptol* (p. 87).
- GELSEMIUM.** The root of *Gelsemium Sempervirens*, *Yellow Jasmine* (Loganiaceæ). Southern States.
The active principle is an alkaloid, *Gelsemin*.
- GUARANA.** The powdered seeds of *Paullinia Sorbilis* (Sapindaceæ). Brazil.
Contains *Caffein* (p. 89).
- HAMAMELIS.** The bark of *Hamamelis Virginica*, *Witch Hazel* (Hamamelidaceæ). N. America.
Composition not yet satisfactorily investigated.
- HYDRASTIS.** The rhizome and rootlets of *Hydrastis Canadensis*, *Yellow Root* or *Golden Seal* (Ranunculaceæ). N. America.
Contains two alkaloids, *Berberia* and *Hydrastin*, the latter said to be the active principle.
- IGNATIA.** *St. Ignatius' bean*, the seeds of *Strychnos Ignatia* (Loganiaceæ). Philippine Isles.
Contains *Strychnia*, in larger amount than *Nux Vomica* (p. 40) and *Brucia*.

RHAMNUS FRANGULA. The bark of *Rhamnus Frangula*, *Black Alder* (Rhamnaceæ). Native.

SPIGELIA. The root of *Spigelia Marilandica*, *Carolina Pink* (Loganiaceæ). Southern States.

Contains *Tannic Acid*, *fixed and volatile oils*, *resin*, and a *bitter principle*.

PREPARATIONS.

The United States Pharmacopœia prescribes—

Extractum Ignatiæ. Dose, $\frac{1}{4}$ — $\frac{1}{2}$ grain.

Extractum Cimicifugæ Fluidum. Dose, 20—60 min.

„ *Chimaphilæ Fluidum.* Dose, 1—2 drms.

„ *Gelsemii* „ „ 5—10 min.

„ *Spigeliæ* „ „ } Dose, 2—4 drms.

„ *Spigeliæ et Sennæ Fluidum.* }

Decoctum Chimaphilæ. Dose, 1—2 oz.

Infusum Spigeliæ. „ 2—4 oz.

A *Tincture*, 1 part in 8, of *Gelsemium* is usually prescribed in this country. Dose, 10—30 min.

A saturated *Tincture* of *Eucalyptus* is sometimes made. It is simply a spirit of Eucalyptol. Dose, $\frac{1}{2}$ —2 drms.

Of *Convallaria*, *extracts* and *tinctures* are made; *Hamamelis*, *Hydrastis*, and *Rhamnus Frangula* are usually prescribed in *tinctures*; which are advertised, with their scales of dose, by the principal manufacturing chemists.

Guarana is administered in the form of powder, Dose, 1—2 drms. *Datura Tatula* in that of cigarettes.

Ignatia is used mainly as a source of *Strychnia*.

Duboisia, as a source of *Duboisin* (? *Atropin*).

EXUDATIONS AND CONCRETE JUICES.

GUMMI RUBRUM. *Australian Red Gum*. An exudation from the bark of *Eucalyptus Rostrata* (Myrtaceæ). Australia.

Contains an astringent principle, probably *Tannin*, and *gummy matter*.
90 p.e. soluble in cold water.

CATECHU NIGRUM. *Black Catechu*. A native extract from the heart-wood of *Acacia Catechu* (Leguminosæ). Pegu.

Similar composition to Pale Catechu (p. 56).

LACTUCARIUM. The juice exuding from incisions in the stem of the *Lactuca Virosa* (p. 18), and *Lactuca Sativa*, *Garden Lettuce*, dried.

Contains *Lactucin*, a substance allied to Mannite (p. 57), and 40 p.e. of a crystalline substance, *Lactucione*.

* * * The doses of the above are given in the Posological Tables.

PREPARATIONS.

Catechu Nigrum is used for *Catechu Pallidum* (p. 56).

Lactucarium is prescribed in pills.

Squire (*Companion*, 12th ed., p. 157) gives formulæ for
Extractum Gummi Rubri Liquidum. Dose, $\frac{1}{2}$ —1 drm.

Decoctum " " Dose, 1—2 oz.

Tinctura " " " 20—40 mins.

Syrupus " " " $\frac{1}{2}$ —1 drm.

Suppositoria " "

* * * As Squire's *Companion* is in the possession of most dispensing Chemists, it is sufficient in prescribing to append "(Squire)" to the name of the preparation to ensure the required strength.

OLEO-RESIN.

BALSAMUM DIPTEROCARPI. *Gurjun Oil*. From *Dipterocarpus Lævis*, *Turbinatus*, and other species (Dipteraceæ). India.

Closely resembles Copaiba.

* * * Used only for external application.

FIXED OIL.

OLEUM GYNOCARDIÆ. *Chaulmoogra Oil*. From the seeds of *Gynocardia Odorata* (Bixaceæ). India.

* * Used only for external application.

VOLATILE OILS.

OL. CHENOPODII. From the fruit ("seeds") of *Chenopodium Anthelminticum* (Chenopodiaceæ). United States.

OL. GUALTHERIÆ. *Oil of Winter Green*. From *Gualtheria Procumbens*, *Partridge-berry* (Ericaceæ). N. America.

Composed of *Methylic Salicylate*, $\text{CH}_3 \cdot \text{C}_7 \text{H}_5 \text{O}_3$. The source of the purest *Salicylic Acid* (see Section III).

OL. EUCALYPTI. *Eucalyptol*. From the leaves of *Eucalyptus Globulus* (p. 84).

Possesses strong antiseptic properties.

OL. SANTALI FLAVI. *Sandalwood Oil*. From the wood of *Santalum Flavum* (Santalaceæ). India.

Very fragrant.

OL. STAPHISAGRIÆ. From the seeds of *Delphinium Staphisagria* (Ranunculaceæ). S. Europe.

* * Ol. *Staphisagria* is used in an ointment (1 in 8) as a *pediculicide* only.

The doses of the rest are given in the Posological Tables.

ACTIVE PRINCIPLES OF DEFINITE COMPOSITION.

CHRY SOPHANIC ACID. $C_{10}H_8O_3$.

Obtained from Goa Powder (*Araroba*), a product found in the clefts of the wood of a Leguminous tree in the province of Bahia; probably produced by the decomposition of a resinous exudation. 80 p. c. of the powder is Chrysophanic Acid.

Acicular crystals, bright yellow.

Slightly soluble in cold water or spirit.

Forms soluble reddish-brown Chrysophanates with alkaline solutions, and insoluble brownish Chrysophanates with Magnesia and some other metallic oxides (see **Rheum**, p. 29).

MENTHOL. $C_{10}H_{20}O$, or Peppermint Camphor.

Deposited from Chinese Oil of Peppermint, which is the essential oil of *Mentha Javanica* (Labiatae), and consists mainly of Menthol and a liquid isomeride (see p. 44).

Acicular crystals, or a crystalline mass, colourless.

Soluble in spirit, ether, and fixed oils, only slightly in water.

Volatilo, with a strong odour of peppermint.

THYMOL. $C_{10}H_{14}O$. A Phenol.

Deposited from Oil of Thyme (*Thymus Vulgaris*—Labiatae) and Oil of Ajowan (*Ptychotis Ajowan*,—Umbelliferae). Egypt and Persia.

Large prisms, colourless.

Fuses at 111 deg. F.

Volatile.

Soluble in $\frac{1}{2}$ part of Rectified Spirit, 200 of Glycerine; very slightly in water. Arrests fermentation.

SALICIN. $C_{13}H_{18}O_7$. A Glucoside.

From the bark of *Salix Caprea*, *Willow*, (Salicaceae). Native.

White scales and silky needles.

Soluble in 28 parts of water; also in alcohol.

Boiled with dilute Sulphuric Acid, or treated with Emulsin, it yields glucose and *Saligenol* $C_7H_8O_2$.* Oxidized by Potassic Bichromate and Sulphuric Acid, it yields *Salicylol* $C_7H_6O_2$,† a compound which forms the essential oil of *Spiraea Ulmaria* (*Meadow Sweet*). (See Salicylic Acid, in Section III.)†

* * The name of *Ergotine* is given to various extracts of Ergot, freed from albumen and gum.

The "*Jalapine*" of the shops is Jalap resin (see p. 48) decolorized by animal charcoal, and precipitated in a finely-divided state.

* $C_{13}H_{18}O_7 + H_2O = C_6H_{12}O_6 + C_7H_8O_2$.

† Salicylol is found in the urine of patients taking Salicin. It strikes a deep purple with Ferric Salts.

(ALKALOIDS.)

CAFFEIN.

From Coffee-berries, the seeds of *Coffea Arabica* (Rubiaceæ). Ceylon, Java, India, &c. Also from the Kola Nut, the seed of *Cola Acuminata* (Sterculiaceæ). Africa.* Used either as the alkaloid, or as a citrate.

The alkaloid in silky acicular or feathery crystals soluble in 100 parts of water, 150 of Rect. Spirit.

The citrate, which is commonly used, occurs in similar crystals, similarly soluble.

CINCHONA ALKALOIDS.

CINCHONIN. $C_{20}H_{24}N_2O$.

CINCHONIDIN. ditto.

QUINIDIN. $C_{20}H_{24}N_2O_2 \cdot 2H_2O$.

From the **Cinchona Barks** (p. 23).

Cinchonin is the most important. It is used in the form of a Sulphate :—

Short oblique prisms.

Soluble in 54 parts of water, freely in spirit.

The others are seldom prescribed alone, but preparations consisting of a mixture of Quinine and Cinchonin with the other Cinchona Alkaloids, are sold.

Cinchonin is distinguished from Quinine by the want of fluorescence of its solutions, and the absence of the green colour which the latter gives with Chlorine and Ammonia.

CODEIA. $C_{18}H_{21}NO_3 \cdot H_2O$.

From **Opium** (p. 58). Crystallizes out on evaporating the liquid left after precipitating Morphia (p. 67). Redissolved in ether and crystallized again.

Octahedra.

Soluble in water and alcohol.

Gives *no* red colour with Nitric Acid, or blue colour with Ferric Salts, as Morphia does.

ESERIN. $C_{15}H_{21}N_2O_2$.

From **Physostigma** (p. 40).

Amorphous, yellowish.

The alkaloid itself is soluble in alcohol, sparingly in water.

Its sulphate, which is commonly used, is of reddish colour, and is freely soluble in water.

* Found also in Tea, in Guarana (p. 84), and in Matè or Paraguay Tea (the leaves of *Ilex Paraguayensis*).

PILOCARPIN.

From the leaves and bark of *Pilocarpus Pinnatifolius*, *Jaborandi*, (Rutaceæ). S. America.

Amorphous, yellow, deliquescent.

The Nitrate is commonly used; it crystallizes in white lamellæ, and is permanent in air.

It is soluble in 8 parts of water.

* * APOMORPHIA. $C_{17}H_{19}NO_3$ (*i.e.* Morphia, less one molecule of H_2O .)
An artificial alkaloid prepared by heating Morphia to $300^\circ F.$, with excess of Hydrochloric Acid, in a closed tube, for several hours.

White powder, turning green on exposure to air and moisture.

Soluble in water.

* * Chrysophanic Acid and Menthol are used only as external applications; the former as an *ointment* with Vaseline or Lard (1 in 8—64); the latter in a saturated *solution*, or else rubbed on in the crystalline form. Thymol is used in various *solutions* as an antiseptic, and has been given internally in the form of pill.

* * The doses of the other active principles are given in the Posologica Tables.

SECTION III.

MINERALS AND PRODUCTS OF MANUFACTURE.

SIX GROUPS.

First Group	Non-metallic Elements and their compounds.
Second	„	Mineral Acids.
Third	„	Alcohol and its derivatives ; Amylic Alcohol and its derivatives.
Fourth	„	Tars and their derivatives ; Organic Acids.
Fifth	„	Pyroxylin, Glycerine, their derivatives, and Miscellanea.
Sixth	„	Metals and their compounds.

The Official Preparations are described with each group.

* * * Non-official drugs in general use are described with the official. The names of the latter are printed in **Clarendon** type, those of the former in ROMAN CAPITALS.

* * * An asterisk (*) prefixed to the description of a process of manufacture signifies that it is one prescribed by the Pharmacopœia (see p. 1, second paragraph).

FIRST GROUP.

<i>a.</i> Aqua.	Sulphuris Iodidum.
Aqua Destillata.	<i>δ.</i> (Os Ustum.)
<i>β.</i> Carbo Ligni.	Phosphorus.
Carbo Animalis.	<i>ε.</i> Liquor Chlori.
Carbo Animalis Purificatus.	Bromum.
<i>γ.</i> Sulphur Sublimatum.	Iodum.
Sulphur Præcipitatum.	<i>ζ.</i> NITROUS OXIDE.

OFFICIAL PREPARATIONS.

a.—**Aqua.** The purest natural water that can be obtained, cleared if necessary by filtration.

* Ten gallons of **Aqua** being distilled from a copper still with a block-tin worm; the first half-gallon that passes over may contain carbonic acid and ammonia. This is rejected. The next eight gallons are almost absolutely pure, and form **Aqua Destillata**. Distillation cannot be carried further without risk of contamination from charred organic matter.

Colourless, odourless, tasteless.

β.—**Billets of Wood** being heated to dull redness in cast iron cylinders;—water, gaseous, and volatile matters, are driven off, and **Carbo Ligni**, (*Wood-Charcoal*), *Carbon* with about 2 p.c. of alkaline and metallic salts, remains.

Brittle porous masses, black.

Insoluble in any known liquid.

Capable of absorbing and condensing gases within its pores (90 times its own volume of Ammonia, 55 of Sulphuretted Hydrogen).

Also of absorbing organic colouring, odorous, and septic matters, and oxidizing them by means of the condensed atmospheric oxygen contained in its pores.

BONES being similarly treated, yield **Carbo Animalis** (*Bone-black*), a mixture of Calcic Phosphate and Carbonate, with about 20 p. c. of *Carbon*, and a little Iron Carbide. (See *Phosphorus*, and *Calcis Phosphas*.)

Grey powder. Used as a filter in coarse processes.



* On digesting **Carbo Animalis** with dilute Hydrochloric Acid, a solution of Calcic Chloride and Acid Calcic Phosphate, $H_4Ca(PO_4)_2$, is substituted for the insoluble carbonate and neutral phosphate, $Ca_3(PO_4)_2$. On filtering off the solution and washing the residue, **Carbo Animalis Purificatus**, almost pure *Carbon*, remains.

Dried by heating to redness in a covered vessel. Black Powder.

Similar properties to those of wood-charcoal.

Power of decolorizing more marked.

Chiefly used as a filtering agent.

Carbo Animalis Purificatus.

γ.—By distilling native Sulphur from its earthy impurities, or by burning the native sulphides, Iron and Copper Pyrites, is obtained *crude, or rough Sulphur*.

Purified by sublimation, it forms **Sulphur Sublimatum** (*Flowers of Sulphur*).

Gritty powder, pale yellow.

Soluble in Benzene, Carbon Disulphide, and Sulphur Chloride; with heat, in Turpentine and Oils.

Fuses at 236 deg. F., volatilizes at 836 deg. F.

Burns in air, forming *Sulphurous Anhydride*, SO_2 .

From **Sulphur Sublimatum** are obtained,—

- 1.* By boiling in water with slaked lime,—a solution of *Calcic Pentasulphide*, CaS_5 , and *Calcic Sulphosulphate*,* CaS_2O_3 . On filtering, and adding Hydrochloric Acid,—a deposit of **Sulphur Præcipitatum** (*Milk of Sulphur*), Calcic Chloride remaining in the solution.

Collected, washed, dried.

Pale yellow powder, not gritty.

Same properties as those of Sulphur Sublimatum.

- 2.* By mixing with four times its weight of Iodine and heating gently, finally fusing—**Sulphuris Iodidum** S_2I_2 .

Allowed to solidify. Steel grey mass, crystalline.

Smells of Iodine.

Soluble in 60 parts of Glycerine.

Decomposed by boiling water.

δ.—BONES, or more commonly, Bone Black (p. 92) which has served its purpose as a filtering agent, is burned in the open fire. Carbon is burnt off, and **Os Ustum** (*Bone-ash*), chiefly *neutral Calcic Phosphate*, $\text{Ca}_3(\text{PO}_4)_2$, with some Calcic Carbonate, remains.

On mixing with Sulphuric Acid,—Calcic Sulphate, and *Acid Calcic Phosphate*, $\text{H}_4\text{Ca}(\text{PO}_4)_2$, are formed. By washing out with water, the latter, which is soluble, is separated from the insoluble sulphate. On evaporating to dryness the solution of *Acid Calcic Phosphate*, and then heating the salt to redness, hydrogen and oxygen are driven off as water, and *Calcic Metaphosphate*, $\text{Ca}(\text{PO}_3)_2$, remains. On distilling with charcoal,† Carbonic Oxide and **Phosphorus** pass over; and neutral Calcic Phosphate, again formed, remains. The **Phosphorus** is condensed under water.



Wax-like, semi-transparent.

Becomes hard and brittle at low temperatures




* Or "Hyposulphite".

† In practice, the charcoal is added before evaporation to dryness is quite complete, to ensure thorough admixture.

Sulphur Præcipitatum.



Phosphorus.



Soluble in Carbon Disulphide, Benzene, Oil of Turpentine, fixed and essential oils. slightly in Ether.

Fuses at 112 deg. F. (may, therefore, be fused under water).

Burns in air, forming phosphoric anhydride, P_2O_5 . Burns slowly in air at all temperatures above freezing-point (must, therefore, be preserved under water).

- ε.* On heating Hydrochloric Acid with Black Oxide of Manganese, *Chlorine Gas* is evolved. Washed and passed into water till a solution is formed containing 2·66 grains in the ounce, it forms **Liquor Chlori**.

Yellowish green, evolving chlorine gas.

Chlorine decomposes many organic substances containing hydrogen by replacing that element; hence is destructive of vegetable colours and of organic odorous or infectious matters. (See *Calx Chlorata*, *Liquor Sodæ Chloratæ*.)

Liq. Chlori decomposes in sunlight, yielding Hydrochloric Acid and Oxygen.

The mother liquor of Sea Water, left after evaporation and crystallization of all crystallizable salts, contains *Magnesian Bromide*. On passing Chlorine into it, **Bromine** separates, colouring the fluid yellow. Ether, shaken with the fluid, takes up the **Bromine** alone, and floats on the surface. The ethereal solution being drawn off and shaken with solution of Caustic Potash, the **Bromine** combines with the latter, forming *Potassic Bromide* and *Bromate*. The watery solution being separated from the ether, evaporated to dryness, and ignited so as to decompose the Bromate, *Potassic Bromide* remains. On distillation with Sulphuric Acid and Black Oxide of Manganese, **Bromine (Bromum)** passes over.

Condensed in a receiver. Dark brown liquid, evolving red fumes. Strong odour.

Volatile. Boils at 139 deg. F. Freezes at -12 deg. F.


Soluble in 30 parts of water. Miscible with Alcohol and Ether in all proportions.

Chemical action similar to that of Chlorine, but less strongly developed.


KELP, the ashes of burnt seaweed, yields on lixiviation, *Sodic Iodide*, Bromide, Chloride, Sulphide, Sulphosulphate and Carbonate, Potassic Chloride, and Potassic-Sodic Sulphate. On evaporating, the sulphate and chlorides, with most of the carbonate, crystallize out, leaving the other salts in the mother liquor, or *ley*. Mixed with Sulphuric Acid, the *ley* evolves Carbonic and Sulphurous Anhydrides and Sulphuretted Hydrogen, Sulphur is precipitated, and Sodic Sulphate crystallizes out. The *ley*, which now contains *Sodic Iodide*, Bromide, and (a little) Chloride, with excess of Sulphuric Acid, being decanted, and distilled with successive small amounts of Black Oxide of Manganese, **Iodum (Iodine)** is liberated and passes over.*

* *Bromine* passes over *after* the Iodine. The temperature is not allowed to rise high enough to liberate *Chlorine*.

Chlorum.



Bromum.



Condensed in receivers ; purified by sublimation.
Bluish-black scales, lustrous.

Soluble in 5,500 parts of water.
 „ 12 parts of Rect. Sp.
 „ 4 parts of Ether.
 „ 4 parts of Chloroform.

Freely soluble in solutions of the Iodides.

Fuses at 236 deg. F. ; volatilizes at 347 deg. F. : volatilizes in the steam of hot watery solutions.

Chemical action similar to that of Chlorine and Bromine, but more feebly developed.

Unites readily with many metals.

Strikes a deep blue colour with starch.

§.—Ammonic Nitrate, $\text{NH}_4 \text{NO}_3$, (see Sixth Group) when heated decomposes into watery vapour and NITROUS OXIDE GAS, N_2O .

Condensed in iron bottles, under a pressure of 50 atmospheres, and firmly secured.

Colourless liquid, becoming gaseous on being liberated.

OFFICIAL PREPARATIONS.

Liquor Chlorig. (See above.)

„ **Iodid.** $2\frac{1}{2}$ gr. in a drachm of solution of Potassic Iodide.

Oleum Phosphoratum. 1-16th gr. in 10 minims of Almond Oil.

Tinctura Iodid. $1\frac{1}{3}$ gr. in a drachm of Rectified Spirit, with half the amount of Potassic Iodide.

Confectio Sulphuris. Sublimed Sulphur, Cream of Tartar, and Syrup of Orange, as 4, 1, 4 parts.

Pilula Phosphori. 1-30th gr. in 3 gr. of Tolu and Beeswax.

Linimentum Iodid. 7 gr. in a drachm of Rectified Spirit, with Camphor and a little Potassic Iodide.

Unguentum Sulphuris. 12 gr. in a drachm.

„ **Sulphuris Iodidid.** 4 gr. in a drachm.

„ **Iodid.** 2 gr. in a drachm.

Vapor Iodid. A drachm of Tinct. Iodid inhaled from an ounce of warm water.

Vapor Chlorig. (See **Calx Chlorata.**)

Cataplasma Carbonis. A quarter of an ounce of *Carbo Ligni* in a poultice made of bread 2 oz., Linseed meal $1\frac{1}{2}$ oz. A similar amount of charcoal sprinkled on the surface.

SECOND GROUP. (MINERAL ACIDS.)

- | | |
|--------------------------------------|---|
| <i>a.</i> Acidum Nitricum. | <i>γ.</i> Acidum Phosphoricum |
| Acidum Hydrochloricum. | Dilutum. |
| <i>β.</i> Acidum Sulphuricum. | <i>δ.</i> Acidum Nitrohydrochloricum |
| Acidum Sulphurosum. | Dilutum. |

PROPERTIES.

OFFICINAL PREPARATIONS.

- a.** On distilling from a mixture of Potassic (or Sodie) Nitrate with diluted Sulphuric Acid,—Acid Potassic (or Sodie) Nitrate, K (or Na)HSO₄, is formed; and *Nitric Acid* passes over, with water and some Nitrous Acid, forming a liquid of sp. gr. 1.42. On heating to expel Nitrous Acid,—**Acidum Nitricum**, containing 70 p. c. of HNO₃, results.

Colourless liquid, emitting acid fumes.

- ** On distilling from a mixture of Sodie Chloride (common salt) with diluted Sulphuric Acid,—Sodie Sulphate is formed, and gaseous *Hydrochloric Acid*, HCl, passes over. Conducted through a wash-bottle into distilled water till the latter has increased in volume by 32 p. c., it forms **Acidum Hydrochloricum**, containing nearly 32 p. c. by weight of H Cl.

Colourless fuming liquid.

- β.*—The fumes of burning Sulphur or Iron Pyrites, FeS₂, containing *Sulphurous Anhydride*, SO₂, and atmospheric air, are passed into a leaden chamber, where water and watery vapour are introduced. By placing a pot of Sodie Nitrate mixed with Sulphuric Acid in the furnace, vapours of Nitric Acid are passed in also.

The Sulphurous Anhydride, SO₂, is oxidized at the expense of the Nitric Acid to *Sulphuric Anhydride*, SO₃, which unites with water to form *Sulphuric Acid*, H₂SO₄; and Nitric Oxide, NO, is liberated.

The following changes then take place, and continue in rotation :—(1.) Nitric Oxide unites with Atmospheric Oxygen to form Nitric Peroxide, NO₂; (2.) Nitric Peroxide, Sulphurous Anhydride, and a little watery vapour unite to form a compound of uncertain nature; which (3.) in presence of more water, is resolved into *Sulphuric Acid* and Nitric Oxide, NO.

On evaporating the dilute acid thus obtained as long as water alone passes off,—**Acidum Sulphuricum**, containing 96.8 p. c. of H₂ SO₄, results.

Oily, colourless liquid.

Acidum Sulphuricum.

- 1.— $3 \text{ SO}_2 + 2 \text{ HNO}_3 + 2 \text{ H}_2\text{O} = 3 \text{ H}_2\text{SO}_4 + 2 \text{ NO}$.
- 2.— $2 \text{ NO} + \text{O}_2 = 2 \text{ NO}_2$.
- 3.— $\text{NO}_2 + \text{SO}_2 + \text{H}_2\text{O} = \text{H}_2\text{SO}_4 + \text{NO}$.

- * On distilling from a mixture of Wood Charcoal with Sulphuric Acid,—Carbonic Anhydride, CO_2 , and *Sulphurous Anhydride*, SO_2 , pass over. The gases being conducted into water,—the former passes through; the latter is absorbed, uniting with water to form *Sulphurous Acid*, H_2SO_3 . On continuing the process till the bubbles of gas pass through the water *undiminished in size*,—**Acidum Sulphurosum**, containing 11.8 p. c. of H_2SO_3 , results.

Colourless liquid, evolving Sulphurous Anhydride.

- γ. * On heating Phosphorus in diluted Nitric Acid,—*Phosphoric Acid*, H_3PO_4 , Phosphorous Acid H_3PO_3 , and Nitric Oxide, are formed. On distilling, *Phosphoric*, Phosphorous, and Nitric Acids, Nitric Oxide and water, pass over. Returning the distillate from time to time, and proceeding till the Phosphorus has all disappeared,—a *dilute Phosphoric Acid*, contaminated with Nitric Acid and Nitric oxides, is obtained. On evaporating till water and Nitric fumes cease to pass off,—*glacial Phosphoric Acid*, H_3PO_4 , is left. On diluting, while hot, to a stated degree,—**Acidum Phosphoricum Dilutum**, containing 13.8 p. c. of H_3PO_4 .

Colourless liquid.

- δ. * On mixing 3 parts of **Acidum Nitricum** with 4 of **Acidum Hydrochloricum**, and standing for 24 hours in a partially closed bottle, *Chlorine* and *Nitrosyl Chloride*, NOCl , are formed, and retained in the bottle. On diluting with 25 parts of water, shaking up the bottle as it is added,—the so-called **Acidum Nitrohydrochloricum Dilutum** is produced.

Colourless liquid, evolving Chlorine.

PROPERTIES.


Specific Gravity of the Official Acida.

Ac. Sulphuricum	. 1.843.
Ac. Nitricum	. 1.420.
Ac. Hydrochloricum	. 1.160.
Ac. Sulphurosum	. 1.040.


Relative Neutralizing Power.

40 gr. of Sod. Hydrate, Na HO ,	are neutralized by
50.6 gr. of Ac. Sulphuricum,	
90 " Ac. Nitricum,	
114.8 " Ac. Hydrochloricum.	
347.4 " Ac. Sulphurosum.	


Acidum Sulphurosum.



Acidum Phosphoricum.



Acidum Nitrohydrochloricum.



Corrosive Action.

Nitric Acid is a powerful oxidizing agent, 4HNO_3 being readily decomposed into $4\text{NO} + 2\text{H}_2\text{O} + 3\text{O}_2$ (Nitric Oxide, water, and nascent Oxygen).

Sulphuric Acid has a powerful affinity for water, abstracting not only water from moist bodies and hydrated substances, but the elements of water from combination in organic compounds.

Hydrochloric Acid Gas possesses a similar but less powerful property.

Hence the *corrosive* action of the corresponding Pharmacopœial **Acida** upon organic substances, and the *desiccating* and *charring* actions of **Acidum Sulphuricum**, which is nearly pure Sulphuric Acid.

In the action on metals of Sulphuric or Hydrochloric Acid, the metallic element simply replaces and sets free the Hydrogen of the Acid, or a portion of it. In the action of Nitric Acid on metals, part of the Acid is decomposed, water and Nitrous Acid, or Nitric Oxide, or Nitric Peroxide, or Nitrogen, or several of these, being set free instead of Hydrogen.

Sulphurous Anhydride or Sulphurous Acid, in presence of certain organic substances, (vegetable colours, ferments, odorous and septic matters,) forms fresh combinations. Hence its *bleaching*, *deodorizing*, and *disinfecting* properties, and its power of *arresting fermentation and putrefaction*. From its (colourless) combinations with vegetable colours the latter may be set free by the action of an alkali.

Acidum Nitricum undergoes slow spontaneous decomposition in sunlight, and becomes coloured by Nitrous Acid.

Acidum Sulphurosum undergoes slow spontaneous oxidation, and becomes converted into a solution of Sulphuric Acid.

Acidum Sulphuricum and water, mixed together, undergo contraction in bulk, heat being disengaged in the process.

OFFICIAL PREPARATIONS OF THE ACIDA.

(DILUTE ACIDS.)

Acidum Nitricum Dilutum. 20 p. c. of the **Acidum** in water.

Acidum Hydrochloricum Dilutum. 30 p. c. of the **Acidum** in water.

Acidum Sulphuricum Dilutum. $8\frac{1}{3}$ p. c. of the **Acidum** in water.

Acidum Sulphuricum Aromaticum. 7 p. c. of the **Acidum** in Rectified Spirit, flavoured by maceration on Cinnamon and Ginger.

Acidum Phosphoricum Dilutum	} Official only in the diluted form. (See above.)
Acidum Nitrohydrochloricum Dilutum	

THIRD GROUP.

ALCOHOLS AND THEIR DERIVATIVES.


- | | |
|---|--|
| <p>α. Spiritus Rectificatus.
 Spiritus Tenuior.
 ABSOLUTE ALCOHOL.
 Acetum.</p> <p>β. Ether. Ether Purus.
 Ether Aceticus.
 Spiritus Etheris Nitrosi.</p> | <p>γ. Chloral Hydras.
 CROTON-CHLORAL HYDRATE.</p> <p>δ. Chloroformum.
 IODOFORM.</p> <p>ε. Alcohol Amylicum.
 Amyl Nitris.</p> |
|---|--|

PROPERTIES.

OFFICIAL PREPARATIONS.

- α.**—By the action of Malt-extract on STARCH, $C_6 H_{10} O_5$,—*Sugar*, $C_{12} H_{22} O_{11}$, is formed, the elements of water being taken into combination.
 By the action of Yeast on SUGAR,—*Glucose*, $C_6 H_{12} O_6$; the elements of water being again combined.
 By a similar fermentation of GLUCOSE,—*Alcohol*, $C_2 H_6 O$; Carbonic Anhydride, CO_2 , being liberated.
 By these means weak alcoholic fluids (*Worts*) are obtained from POTATOES or GRAIN, which contain Starch; from infusions of MALT, which contains a crystallizable form of sugar; from MOLASSES, or the residue of the Beet Sugar manufacture, which contain uncrystallizable forms; or from GRAPE-JUICE, which contains Glucose.
 On distilling such *Worts*, the Alcohol passes over in the first portions of the distillate. By stopping the distillation at an early stage, *Spirits*, liquids containing a large proportion of Alcohol, are obtained.
 Repetitions of the process (rectification) furnish *Spirits* containing progressively larger proportions of *Alcohol*.
 Spirit rectified till only 16 p. c. of water remains (84 p. c. of Alcohol) forms **Spiritus Rectificatus**. It is free from flavouring or colouring matter.
 * Diluted with three-fifths of its bulk of water, it forms **Spiritus Tenuior** (*Proof Spirit**), containing 49 p. c. of Alcohol.

* The old "proof", from which this apparently arbitrary standard was fixed, was to saturate gunpowder with the spirit and ignite it. If "above proof", the alcohol fired the gunpowder; if "below proof", it burned off, and left the powder damp.

Alcohol.

Simple distillation fails to separate Alcohol from a final 10 p. c. of water. By digesting strong rectified spirit on Quicklime, CaO , this residual water is separated to form Calcic Hydrate, CaH_2O_2 ; and ABSOLUTE ALCOHOL, $\text{C}_2\text{H}_6\text{O}$, can be distilled off.

On leaving the Alcoholic *Wort* obtained by fermenting infusion of Malt or unmalted grain exposed to the air, the Alcohol is oxidized (through the agency of a microscopic organism, the *Mycoderma Aceti*) to *Acetic Acid*, $\text{C}_2\text{H}_4\text{O}_2$;—and **Acetum** (*Malt Vinegar*), containing about 5 p. c. of Acetic Acid, results.

Pale brown liquid, evolving Acetic Acid.

β .—From **Spiritus Rectificatus** are prepared: 1. **Æther**. 2. **Æther Aceticus**. 3. **Spiritus Ætheris Nitrosi**.

1. *On distilling **Spiritus Rectificatus** with Sulphuric Acid, the following changes take place. α . *Ethylic Hydric Sulphate*, $\text{C}_2\text{H}_5\cdot\text{H}\cdot\text{SO}_4$, and water are formed. β . By reaction of the former with a fresh portion of the Alcohol, *Ethylic Ether*, $(\text{C}_2\text{H}_5)_2\text{O}$, is formed, and Sulphuric Acid set free again.

γ . The *Ether* passes over, with water, alcohol, Sulphurous Acid, Acetic Acid, *Ethylic Sulphate*, and other substances, which are formed in small quantities at the same time.

δ . The Sulphuric Acid remains, ready to act on a fresh portion of the Spirit, which in practice is supplied continuously to the still, until the Sulphuric Acid is wasted in producing the impurities alluded to above.

On agitating the distillate with slaked Lime and a solution of Calcic Chloride, the impurities are mostly dissolved and separated from the *Ether*, which floats on the surface. On separating the latter, and redistilling,—**Æther**, containing 92 p. c. of *Ethylic Ether*, the rest being chiefly Alcohol,—is obtained.

On agitating **Æther** with water to remove the Alcohol, then digesting with recently burnt lime and Calcic Chloride, to remove all traces of water, and finally distilling again, **Æther Purus**, pure *Ethylic Ether*, $(\text{C}_2\text{H}_5)_2\text{O}$.


2* On mixing with Sulphuric Acid, *Ethylic Hydric Sulphate* is formed as before. On adding fused (anhydrous) Sodid Acetate and distilling, **Æther Aceticus** (*Ethylic Acetate*) $\text{C}_2\text{H}_5\cdot\text{C}_2\text{H}_3\text{O}_2$, is formed, and passes over, with water and other impurities.

Purified by redistillation from Calcic Chloride.


3* On mixing with Sulphuric Acid,—*Ethylic Hydric Sulphate*, as before. On distilling with Nitric Acid and Copper, *Ethylic Nitrite*, $\text{C}_2\text{H}_5\cdot\text{NO}_2$, is formed, and passes over, with alcohol and water. On dilution with Rectified Spirit to a stated degree,—**Spiritus Ætheris Nitrosi**, containing 2 p. c. of *Ethylic Nitrite*, or *Nitrous Ether*.

All three colourless liquids, with peculiar odours.

Acetic Acid.



Æther.



γ.—On passing dry Chlorine gas through absolute Alcohol,—after a long series of substitution changes,—*Chloral Alcoholate*, $C_2HCl_3O.C_2H_5O$, is formed. Other products of the action of Chlorine on the alcohol are mixed with it. On shaking with Sulphuric Acid,—*Chloral* (Trichloraldehyde), C_2HCl_3O , separates as an oily layer, and is distilled off. By redistillation with quicklime, it is obtained in a still purer state. On adding a small quantity of water,—**Chloral Hydras**, $C_2HCl_3O.H_2O$, is formed; with evolution of heat.

Crystallizes on cooling. Rhomboidal plates.

On passing Chlorine gas into Ethylic Aldehyde (C_2H_4O), *Crotonic Aldehyde*, C_4H_6O , is first formed by the abstraction of the elements of water; and, by a further substitution change,—the so called CROTON-CHLORAL HYDRATE (*Trichlorocrotonic Aldehyde*), $C_4H_3Cl_3O$.

Minute shining scales.

δ. * On mixing Bleaching Powder and Slaked Lime with freely-diluted Alcohol, and exposing the mixture to a heat between 100 deg. and 180 deg. F.,—among a variety of products, some of which combine with the lime,—*Chloral Hydrate* is formed; but, in presence of the lime, is immediately decomposed to form Calcic Formate $Ca(CHO_2)_2$ and **Chloroformum** (Trichloromethane) $CHCl_3$. On distillation, **Chloroformum**, water, and a little alcohol pass over.

Washed from alcohol by water, from water with Sulphuric Acid. Finally redistilled from Calcic Chloride and Slaked Lime.

Colourless liquid, ethereal odour.


A solution of alkaline Iodide and Iodate, formed by adding Iodine to one of Caustic Alkali, is mixed with Alcohol and free Alkali. On passing Chlorine into the mixture, *Iodine* is liberated. By a reaction with the Alcohol and Alkali similar to that concerned in the production of Chloroform,—**IODOFORM** (*Tri-iodomethane*), CHI_3 , is produced, and deposited from the solution.

Filtered and washed. Yellow scales.


ε.—The WORTS produced by fermenting saccharine solutions with yeast (see above, p. 99) contain two forms of *Amylic Alcohol*, $C_5H_{12}O$, boiling at 268 deg. and 270 deg. F. respectively. In distilling crude spirit (p. 99), they pass over. In rectifying the spirit, they are left behind with the water, the Acetic, and other ethers. With the ethers, they separate spontaneously from the water, forming *Oil of Grain*. On fractional distillation, the ethers pass over first, the *Amylic Alcohols*, nearly pure, last; forming **Alcohol Amylicum** (*Fusel Oil*).

Colourless liquid, oppressive odour.



Æther Aceticus.




Spiritus Ætheris Nitrosi.



Croton Chloral.



Chloroformum.



From **Alcohol Amylicum**, further purified, is formed,—

- a. By passing Nitrous Anhydride into it, **Amyl Nitris**, $C_5H_{11}NO_2$.
Purified from adhering Acid, by mixing with Caustic Alkali, and redistilling.

Or, β . By heating with Nitric Acid;

- Or, γ . By a process similar to that employed in making Nitrous Ether (see above),—**Amyl Nitris**, with various other products, from which it is almost entirely separated by distilling at a temperature below 212 deg. F.

Purified from acid by mixing with Caustic Alkali; from the remains of the other impurities by distilling at 205 deg. F.
Yellowish fluid, peculiar odour.

* * * The preparation of Valerianic Acid from Amylic Alcohol is described under **Sodæ Valerianas**, in the Sixth Group.

PROPERTIES.

(SOLIDS.)

Chloral Hydrate is soluble in $\frac{1}{3}$ part by weight of water (3 oz. in 1 oz. water gives 2 oz. $5\frac{1}{2}$ drachms of solution.—*Squire*), in less than 1 part of Rectified Spirit.

Iodoform is soluble in 120 parts of Rectified Spirit, in 20 parts of Ether, or Chloroform, freely in Olive Oil (not in water).

Croton-Chloral is soluble in 50 parts of water, in equal parts of Rectified Spirit or glycerine.

Chloral Hydrate is fusible at a gentle heat. Chloral Hydrate, in presence of alkalies, is decomposed, giving Chloroform and a Formate. (See the process for making chloroform.)

A mixture of equal parts of Chloral Hydrate and Camphor liquefies spontaneously, and may be used as a liniment.

(LIQUIDS.)

Specific gravity.

Æther Purus720
Æther735
Absolute Alcohol794
Alcohol Amylicum818
Spiritus Rectificatus838
Spiritus Ætheris Nitrosi845
Amyl Nitris877
Æther Aceticus900
Spiritus Tenuior920
Acetum	1.018
Chloroformum	1.490

Boiling points.

Æther Purus	,	.	.	96 deg. F.
Æther	.	.	.	105 "
Chloroformum	.	.	.	142 "
Æther Aceticus	.	.	.	166 "
Absolute Alcohol	.	.	.	172 "
Spiritus Rectificatus	.	.	.	180 "
Amyl Nitris	.	.	.	205 "
Alcohol Amylicum	.	.	.	270 "

Ether and Chloroform, and in a less degree Alcohol and Spirits, evaporate rapidly at ordinary temperatures, abstracting heat powerfully from contiguous bodies.

Rectified Spirit mixes with Water, Ether, Acetic Ether, Amylic Alcohol, and Amyl Nitrite, in all proportions.

Ether mixes with Acetic Ether and Amylic Alcohol in all proportions.

Æther purus is soluble in 10 parts by measure of water. It also takes up 1-10th part of water.

Acetic Ether is soluble in 12 parts of water.

Chloroform in 200 parts of water, $1\frac{1}{2}$ part of Ether, 7-10ths part of Rectified Spirit (not in Glycerine).

Amyl Nitrite is insoluble, Fusel Oil hardly soluble, in water.

Eighteen parts of Rectified Spirit mixed with 18 parts of water are contracted to 34 parts. (*Squire*).

Rectified Spirit dissolves balsams, Camphor and volatile oils, Castor Oil, Sugar, Tannic and Gallic Acids, vegetable alkaloids and colouring matters, Iodine and Bromine, Caustic Potash and Soda, Lithia, Ammonia, and many alkaline and metallic salts, chiefly those that are deliquescent.

It partially dissolves wax with heat.

Ether and Chloroform dissolve resins, balsams, Gutta Percha, Camphor, fixed and volatile oils, fats, most vegetable alkaloids, Iodine and Bromine.

They separate the last three from watery solutions.

They dissolve wax partially.

Ether dissolves Corrosive Sublimate and Gun-Cotton.

Ether exposed to air and light is partially decomposed, Acetic Acid being formed.

The vapour of Ether makes an explosive mixture with atmospheric air.

OFFICINAL PREPARATIONS OF THE ALCOHOLS AND THEIR DERIVATIVES.

Aq. Chloroformi. Water saturated with *Chloroform*.

Spiritus Chloroformi. *Chloroform*, diluted with 19 parts of Rectified Spirit.

Spiritus Ætheris. *Ether*, diluted with 2 parts of Rectified Spirit.

Spiritus Ætheris Nitrosi (sec above). Contains 2 p.c. of Nitrous Ether.

Tinct. Chloroformi Co. *Chloroform*, diluted with 4 parts of Rect. Spirit and 5 parts of Tr. Cardamom. Co. (twice as strong as the *Spiritus*).

Syrupus Chloral. 10 grains in a drachm.

Vin. Aurantii. An alcoholic fluid made by fermenting saccharine solutions flavoured with bitter orange peel ("Orange Wine" of the grocers). Contains about 12 p. c. of Alcohol.

Vin Xericum. Sherry Wine. Contains 17 or 18 p. c. of Alcohol.

Spiritus Vini Gallici. French Brandy. Contains from 48 to 56 p. c. of Alcohol.

Used as vehicles.

Mistura Spiritus Vini Gallici. 4 oz. of French Brandy and the same amount of *Aq. Cinnamomi*, mixed with the yolks of two eggs, and sweetened with half an ounce of sugar.

Linimentum Chloroformi. *Chloroform*, mixed with equal parts of *Lin. Camphoræ*. (Oil is soluble in *Chloroform*.)

* * **Amyl Nitrite**, which is used as an inhalation, is usually dispensed in thin glass *capsules*, holding three minims, which are readily broken in a handkerchief at the time of use.

FOURTH GROUP.

a. Pix Liquida.

OIL OF CADE.

Creasotum.

OFFICINAL PREPARATIONS.

β. Acidum Aceticum.**Ac. Aceticum Glaciale.****γ. Ac. Carbolicum.**

SALICYLIC ACID.

δ. Ac. Tartaricum.**Ac. Citricum.****ε. Ac. Tannicum.****Ac. Gallicum.****ζ. LACTIC ACID.****η. Ac. Hydrocyan. Dil.**

NEUTRALIZING POWER.

OFFICINAL PREPARATIONS.

* * For **Ac. Benzoicum**, see pp. 51-61.

For CHRYSOPHANIC ACID, p. 88.

a.—On destructive distillation of the wood of *Pinus Sylvestris* and other species of Pines, three classes of products pass over.

- 1.—Gaseous bodies, chiefly Carbonic Anhydride, Carbonic Oxide, and Hydrogen.
- 2.—Water, and liquids soluble in water, chiefly Acetic Acid, Methyl, Ethyl, and Allyl alcohols, Methylic Acetate, and Acetone.
- 3.—Oily liquids insoluble in water, holding in solution solid substances also insoluble in water.

The liquids and solids of Class 3, which include a number of hydrocarbons, paraffins, benzene, etc.; a number of alcohols, chiefly phenols; resinous substances; and various other bodies such as naphthalene and pyrene;—form *Wood-tar* or **Pix Liquida**.

Thick, viscid, brownish-black, with an aromatic odour.

OLEUM CADINUM, *Oil of Cade*, consists of the oily matters furnished by the tar distilled from the wood of *Juniperus Oxycedrus* (*Coniferæ*) in France and Germany.

Composition similar to other Tar-oils.

The heavier portions of the *Oil* obtained from *Wood tar* contain a variety of alcohols, chiefly *phenols*, with a number of hydrocarbons, some acids, and other compounds. By washing these heavier portions with a solution of Sodid Carbonate, they are freed from acid. On distilling, two oils are obtained, a heavier and a lighter, which separate on mixing with water. The lighter oil, which floats on the water, contains the *alcohols* and some of the hydrocarbons.

On adding a solution of caustic potash to the *lighter oil* the *alcohols* unite with the potash to form soluble compounds. The solution is decanted. On boiling in the open air, some impurities separate. On adding dilute Sulphuric Acid, the *alcohols* are deposited, but not quite in a pure state. By repeating the treatment with alkali and acid alternately, the

impurities (hydrocarbons) are gradually eliminated. By digesting with Calcic Chloride, the adherent water is abstracted; and on distilling, the *Alcohols* pass over;—forming **Creasotum**.

Oily liquid, colourless, strongly refracting, pungent odour, burning taste.

Sp. gr., 1.071. Boiling point about 400 deg. F.

Soluble in 400 parts of water, in equal parts of glacial Acetic Acid.

Insoluble in Glycerine.

Separates on diluting its solution in Acetic Acid with water.

Coagulates albumen; preserves animal substances from decay.

Ignites spontaneously with Argentic Oxide.

OFFICIAL PREPARATIONS.

Mistura Creasoti. Creasote dissolved in water faintly acidulated with Acetic Acid, sweetened, and flavoured with Juniper. One minim of Creasote in the fluid ounce.

Vapor Creasoti:—12 minims of creasote in 8 oz. of boiling water. Air to be passed *through* the solution.

Unguentum Creasoti. Creasote in Simple Ointment,—1 part in 9.

Unguentum Picis Liquidæ. 5 parts of Tar stirred into 2 of melted Beeswax.

β.—On distilling the *liquid products* of the destructive distillation of wood, Methylic Alcohol and Acetone pass over first. They are separated. On further distillation, "Pyroligneous Acid", *i.e.*, a dilute and impure *Acetic Acid*, passes over. By mixing "Pyroligneous Acid" with milk of Lime, a solution of impure *Acetate* ("pyrolignite") of Lime is obtained, from which Acetic Acid for manufacturing purposes is prepared.

On mixing the solution of "pyrolignite" with one of Sodie Sulphate, Calcic Sulphate is precipitated, and impure *Sodie Acetate* remains in the solution, which is decanted and evaporated to dryness. On fusing the *Acetate*, tarry matters are burned off; by redissolving and repeatedly crystallizing it, it is completely purified.

On mixing the purified and crystallized *Sodie Acetate*, $\text{Na C}_2\text{H}_3\text{O}_2 \cdot 3\text{H}_2\text{O}$, with Sulphuric Acid, and distilling,—**Acidum Aceticum**, containing 33 p.c. of Acetic Acid, $\text{C}_2\text{H}_4\text{O}_2$, passes over.

Colourless liquid, pungent odour.

By using fused (*i.e.*, anhydrous) Sodie Acetate, and concentrated **Acidum Sulphuricum**,—**Acidum Aceticum Glaciale**, containing at least 99 p.c. of $\text{C}_2\text{H}_4\text{O}_2$, is obtained.

Colourless liquid, with a more powerful odour than the **Acidum**; or flat pearly plates.

The strongest Acetic Acid obtainable.

Specific Gravity of

Acidum Aceticum . . . 1.044
 „ „ Glaciale . 1.065, increased by adding 10 p. c. of
 „ water.

Acidum Aceticum Glaciale crystallizes at 34 deg. F., and remains
 crystalline till 48 deg. F.

Acetic Acid volatilizes at ordinary temperatures.

Acetic Acid mixes with water, Rectified Spirit, and ether in all pro-
 portions.

The Glacial Acid dissolves Resins, Gum-resins, Camphor, volatile oils,
 and Creasote. They are precipitated on dilution with water.

The Glacial Acid is highly corrosive to organic substances.

OFFICIAL PREPARATION.

Acidum Aceticum Dilutum. 12½ p.c. of the Acidum in
 water (*i.e.* it contains about 4 p.c. of Acetic Acid).

γ.—On distilling COAL-TAR, Ammonia and some permanent gases pass over
 first, then water and Ammoniacal salts. With the water come *oily*
matters, at first lighter than water, but gradually increasing in specific
 gravity. The *lighter oils* that float on the water contain *Phenol*
 (**Acidum Carbolicum**), C₆H₅O, with Cresol and other homologues ;
 also Naphthalene and other hydrocarbons, and many bodies of undeter-
 mined constitution.

On agitating these *lighter oils* with Sulphuric Acid, some of the impurities
 are deposited as a tarry mass. On treating the remaining oil with
 solution of Soda, or milk of Lime, *Phenol* and its homologues
 enter into combination with the base ; and a solution of *Phenates*, etc.,
 is obtained, which holds Naphthalene and other substances dissolved.
 On dilution, much of the Naphthalene is precipitated ; on exposure to
 air, other impurities also. By the cautious addition of Sulphuric Acid,
 the homologues, which are weaker acids than Phenol, are precipitated,
 leaving a comparatively pure solution of *Phenate* and Sulphate. On
 saturating the filtered solution with Sulphuric Acid, *Phenol*, compara-
 tively pure, is precipitated. By repeating the processes of solution in
 alkali and neutralization with acid, the *Phenol* is further purified.
 Finally, on fractional distillation, pure **Acidum Carbolicum** passes
 over at and about 360° F.

Colourless liquid, pungent odour.

Crystallizes on cooling. Long needles, which become pink on keeping.

Soluble in 15 parts of water.

1½ part of olive oil.
 ½ „ chloroform.
 2 „ glycerine.
 ¼ „ ether.
 ½ „ alcohol. (*Squire.*)

Also in volatile oils.

Fuses at 95 deg. F. When fused it will unite with from 1-8th to 1-4th
 part of water to form an oily fluid ("*Carbolic Oil*"), which re-
 mains liquid at ordinary temperatures.

At and above 95 deg. F., "*Carbolic Oil*" is miscible with water in all proportions; but below that temperature the acid dissolves in water only in the normal proportion, and any excess of the "*oil*" remains separate, in globules which sink to the bottom of the vessel.

Deliquescent in air above 60 deg.

Volatilizes slowly at ordinary temperatures.

Does not redden litmus.

Coagulates albumen.

"Carbolic Oil" destroys animal tissues, coagulating albumen and abstracting water.

Carbolic Acid destroys septic and odorous particles, even in a diluted form.

Carbolic Acid being mixed with Caustic Soda,—*Sodic Phcnate*, $\text{NaC}_6\text{H}_5\text{O}$, is formed. On passing Carbonic Anhydride through *Sodic Phcnate*, dried and heated to 220—250 deg. F.,—*Sodic Salicylate*, $\text{NaC}_7\text{H}_5\text{O}_3$, is formed. On decomposing a solution of *Sodic Salicylate* by an acid, SALICYLIC ACID, $\text{HC}_7\text{H}_5\text{O}_3$, is precipitated.

Washed, and dissolved in boiling water. Crystallizes out on cooling. Shining white needles.

Always contaminated with some other substances when produced by this method.

Or:—

OIL OF WINTER GREEN (p. 87) is mainly composed of *Methyl Salicylate*, $\text{CH}_3\cdot\text{C}_7\text{H}_5\text{O}_3$. On boiling the oil with a solution of Caustic Potash, *Potassic Salicylate* is formed. By distillation, it is separated from other substances. On adding Hydrochloric Acid to a solution of the distilled salt, **Salicylic Acid** is precipitated.

Redissolved and purified by crystallization, as above.

Obtained nearly pure by this method, which is, however, a costly one.

Soluble in 800 parts of cold water.

10 " boiling water.

5 " Rectified Spirit.

Also in 8 " Lard at 180 deg. F. (*Squire*).

Soluble in 24 parts of cold water with 1 part of Potassæ Acetas, or 2 of Potassæ Citras.

Destroys septic particles. Preserves organic matters from putrefaction.

* * Salicylic Acid is usually prescribed for internal administration in the form of its Sodic or Ammonic Salt.

SODIO SALICYLATE occurs in minute pearly scales.

It is soluble in equal parts of water; in 4 parts of Rectified Spirit.

AMMONIO SALICYLATE occurs in powder.

It is soluble in equal parts of water.

δ. * On adding Chalk to a solution of Acid Potassic Tartrate (p. 116) in boiling water,—Carbonic Anhydride is given off, *Calcic Tartrate* is precipitated, and neutral *Potassic Tartrate* remains in solution. On adding, further, a solution of Calcic Chloride, the whole of the Tartaric Acid is precipitated as *Calcic Tartrate*; Potassic Chloride remaining in solution. On digesting the *Calcic Tartrate* (collected and washed) with diluted Sulphuric Acid, a solution of **Acidum Tartaricum**, $\text{H}_2\text{C}_4\text{H}_4\text{O}_6$, with a little Calcic Sulphate, is obtained; the greater part of the Calcic

Sulphate remaining undissolved. On evaporating the filtered solution to a sp. gr. of 1.21, the sulphate crystallizes out. On further evaporating the mother liquor, **Acidum Tartaricum** crystallizes out.

Purified by recrystallization. Oblique rhombic prisms.

* LEMON JUICE contains *Citric Acid*, $\text{H}_3\text{C}_6\text{H}_5\text{O}_7$. On neutralizing with **Creta Præparata**, *Calcic Citrate* is precipitated. From the *Calcic Citrate*, **Acidum Citricum**, $\text{H}_3\text{C}_6\text{H}_5\text{O}_7 \cdot \text{H}_2\text{O}$, is prepared precisely as **Acidum Tartaricum** from *Calcic Tartrate*.

Rhombic prisms.

Acidum Tartaricum dissolves in—

$1\frac{1}{2}$ part of water; 8 parts of Rectified Spirit.

Acidum Citricum in—

$\frac{2}{3}$ part of water; $1\frac{1}{2}$ part of Rectified Spirit; 2 parts of Glycerine.

ε.—In powdered **Galls** (p. 56), which contain a glucoside, *Gallo-tannin*, are formed—

1.* On exposure to air and moisture for three days,—**Acidum Tannicum**, $\text{C}_{27}\text{H}_{22}\text{O}_{17}$.

Extracted by ether and water. The former allowed to evaporate spontaneously, the latter at 212 deg. F.

Scales or amorphous masses; light brown.

2.* On similar treatment for six weeks,—*Gallic Acid*, $\text{H}_3\text{C}_7\text{H}_3\text{O}_5$. Extracted by boiling water. On cooling, **Acidum Gallicum**, $\text{H}_3\text{C}_7\text{H}_3\text{O}_5 \cdot \text{H}_2\text{O}$, crystallizes out.

Needles; fawn-coloured.

Tannic Acid is soluble in—

Equal parts of water or Rectified Spirit; 3 parts of Glycerine.

Gallic Acid in—

100 parts of cold water; 3 parts of boiling water; 8 parts of Rectified Spirit;* 20 parts of Glycerine.

Both **Gallic** and **Tannic Acids** give a black precipitate with **Ferric Salts**.

Tannic Acid (not **Gallic**) precipitates albumen, gelatine, and vegetable alkaloids.

Tannic Acid, on boiling with **Sulphuric Acid**, yields glucose and **Gallic Acid**.

ζ.—On standing a mixture of 8 parts of Sugar and 1 part of Cheese in 50 parts of water, at a temperature of 70—80 deg. F., for two or three weeks,—*Lactic Acid*, $\text{C}_3\text{H}_6\text{O}_3$, is slowly formed by a process of fermentation. Chalk being added from time to time, crystals of *Calcic Lactate* are formed. They are separated, washed, and purified by dissolving in boiling water, and recrystallizing. On careful addition of **Sulphuric** or **Oxalic Acid**, **LACTIC ACID** separates; **Calcic Sulphate** or **Oxalate** remaining.


A syrupy liquid, pale brown.

Miscible in all proportions with water, Rectified Spirit, or Ether.

η.—On heating **REFUSE ANIMAL MATTER** to dull redness with **Potassic Carbonate** and **Iron filings**, the **Carbonate** loses oxygen, and the **Potassium com-**

* From this solution the acid is not precipitated by dilution.

Acidum Tartaricum.



bins with carbon and nitrogen to form *Potassic Cyanide*, KCN , (KCy). On lixiviating the mass in the open air, the Cyanide dissolves, and combines with the iron and with oxygen, forming *Potassic Ferrocyanide* (**Potassæ Prussias Flava**), K_4FeCy_6 , and Potassic Hydrate, KHO , both of which remain in solution. On evaporating, the *Ferrocyanide* crystallizes out.

(Truncated pyramids, lemon-yellow.)

- * On distilling from a mixture of Sulphuric Acid with solution of Potassic Ferrocyanide,—*Hydrocyanic Acid*, HCN , passes over; Acid Potassic Sulphate and Ferrous Dipotassic Ferrocyanide, $\text{FeK}_2\text{FeCy}_6$, remaining. Passed into water, the acid is dissolved. Diluted to a stated degree, it forms **Acidum Hydrocyanicum Dilutum**, containing 2 p. c. of HCN .

Colourless liquid, evolving Hydrocyanic Acid.

RELATIVE NEUTRALIZING POWER OF THE ORGANIC "ACIDA."

40 grains of Caustic Soda are neutralized by—

3 drachms of Acidum Aceticum.

1 drachm of Acid. Acet. Glaciale.

70 grains of Acidum Citricum.

75 grains of Acidum Tartaricum.

OFFICINAL PREPARATIONS OF THE ORGANIC "ACIDA."

Acidum Aceticum Dilutum. See above, under Acidum Aceticum.

Glycerinum Acidi Carbolici. Nearly 1 gr. in 5 min.; $11\frac{1}{2}$ gr. in 1 drachm.

,, **Acidi Tannici.** 1 gr. in 5 min.; 12 gr. in 1 drachm.

,, **Acidi Gallici.** ditto.

Suppositoria Acidi Tannici. 3 gr. in each, with Cocoa-butter, lard, and wax.

,, **Acidi Tannici cum Sapone.** 3 gr. in each, with curd soap and starch.



,, **Acidi Carbolici cum Sapone.** 1 gr. in each, with curd soap and starch.

Trochisci Acidi Tannici. $\frac{1}{2}$ gr. in each, flavoured with Tolu.

Vapor Acidi Hydrocyanici. 10 to 15 min. of Ac. Hydrocy. Dil., inhaled from a drachm of water.

* * * Lactic Acid is usually prescribed diluted with water.

Ac. Hydrocyanicum Dilutum.



FIFTH GROUP.

Pyroxylin.**Collodium.****Collodium Flexile.****Glycerinum,**

NITROGLYCERIN.

Mica Panis.**Theriaca.**

COTTON WOOL (*Gossypium*, p. 40) consists almost entirely of pure *Cellulose*, $C_6H_{10}O_5$. (*) On soaking for three minutes in a mixture of equal parts of **Acidum Sulphuricum** and **Ac. Nitricum**, two Nitrates, $C_{18}H_{29}O_7(NO_3)_8$ and $C_{18}H_{23}O_8(NO_3)_7$, are produced, which form **Pyroxylin** (*Gun-cotton**).

Washed from acid, dried at 212 deg. F.

Indistinguishable in appearance from the original Cotton-wool.

Explosive.

Soluble partly in ether (the second nitrate), and partly in a mixture of ether and alcohol (the first nitrate).

Collodium (*Collodion*) is a solution of Pyroxylin in 36 parts of Ether with 12 of Rectified Spirit.

Colourless.

Dries rapidly, leaving a film impermeable to air, which contracts in drying.

Highly inflammable.

Collodium Flexile is Collodion with 20 grains of Canada Balsam and 10 minims of Castor Oil added to the fluid ounce.

Leaves a film which does not contract.

FIXED OILS and FATS are composed of Glyceric salts of acids of the Acetic or Oleic Series, and yield *Glycerin*, $C_3H_8O_3$, on decomposition with alkalies (see Soaps, pp. 116, 120; also *Emplastrum Plumbi*, under Lead Compounds; and p. 42).

On placing fat and water in a gently heated still, and passing steam heated to 500 deg. F. into the mixture, the Glyceric Salts are decomposed; and water, *Glycerin*, and fatty acids pass over. The aqueous solution of *Glycerin* is separated from the insoluble acids. On evaporating till water ceases to be driven off, **Glycerinum** (*Glycerine*), containing 95 p. c. of Glycerin, is left.

A syrupy, colourless fluid; sweet taste.

Sp. gr., 1.250.

Miscible with water, Rectified Spirit, or chloroform in all proportions.

Not miscible with ether or oils.

Absorbs water from the air.

Distils with superheated steam at 500 deg. F.; cannot be distilled alone at ordinary pressures without decomposition.

* In preparing gun-cotton for explosive purposes, more concentrated acids are used, and a tri-nitrate $C_6H_7O_2(NO_3)_3$ is produced, which is more explosive than **Pyroxylin**, but is soluble only in Acetic Ether.

Dissolves, amongst other substances, Tannic and Gallic Acids, Carbolie Acid, Borax, Iodine, Bromine, Iodide of Sulphur, Perchloride of Iron, Salicylic Acid, Nitrate of Bismuth, Salts of Morphia, Quinia, Strychnia, Veratria, and Atropia.

Pure, or diluted with 10 parts of water, it preserves animal matters from putrefaction.

On adding Glycerine to concentrated Nitric Acid, cooled by a mixture of ice and salt, and afterwards mixing with concentrated Sulphuric Acid, NITROGLYCERIN, or GLONOLIN, (*Glyceric tri-nitrate*), $C_2H_5(NO_3)_3$, separates.

A yellow oil.

Slightly soluble in water, more so in Rectified Spirit.

Highly explosive.

Mica Panis is the *crumb* of ordinary wheaten bread.

Theriaca, *Treacle*, is an uncrystallizable modification of Cane-sugar, formed during the process of refining. (See p. 60.)

Very tenacious.

* * * Glycerine is used as a solvent (p. 7); Mica Panis in making Charcoal Poultice (pp. 10, 95); Treacle in making pills.

Nitroglycerin is supplied as a 1 p. c. solution in Rectified Spirit; and is usually administered in the form of a pill.

SIXTH GROUP.

METALS AND METALLIC COMPOUNDS.

Described in the following order.

POTASSIUM COMPOUNDS.		BISMUTH COMPOUNDS
SODIUM	„	ANTIMONIAL „
AMMONIUM	„	ARSENICAL „
IRON	„	MAGNESIUM „
MERCURIAL	„	ALUM.
CALCIUM	„	LITHIUM COMPOUNDS
ZINC	„	SILVER „
LEAD	„	COPPER, CADMIUM, and CERIUM COMPOUNDS

OFFICINAL PREPARATIONS.

* * * *The Properties are described at the end of each division.*

POTASSIUM COMPOUNDS. 17 *Officinal.*

NITRE.	{ <i>Potassæ Nitras.</i> <i>Acid Potassic Sulphate.</i>	}	Potassæ Sulphas.
				Potassæ. { <i>Potassii Iodidum.</i> <i>Potassii Bromidum.</i> <i>Potassæ Permanganas.</i> <i>Sapo Mollis.</i>
WOOD ASHES.—	Potassæ Carbonas.	{	}	Potassæ Bicarbonas.
				Potassæ Acetas.
				Potassæ Citras.
				Potassæ Chloras.
				Potassa Sulphurata.
				Potassæ Bichromas.
				Potassæ Prussias Flava.
				Potassæ Tartras.
ARGOL.—	Potassæ Tartras Acida. ..			Potassæ Tartras.

NITRE, or SALTPETRE. Impure *Potassic Nitrate*, formed spontaneously in soils containing the base *Potassa*, by decomposition of organic matters and oxidization of the resulting Ammonia to form Nitric Acid; found as an efflorescence in the E. Indies, cultivated artificially in Europe. Separated by lixiviation and crystallization.

a. From NITRE are obtained,

1. By solution and recrystallization,—pure **Potassæ Nitras**, KNO_3 .

Six-sided prisms, striated.

2. By heating with diluted Sulphuric Acid,—a solution of *Acid Potassic Sulphate*, KHSO_4 (see *Acidum Nitricum*, p. 96). On neutralizing with Potassic Carbonate,—one of *neutral Potassæ Sulphas*, K_2SO_4 .

Crystallizes out on evaporation.

Six-sided prisms, with pyramidal ends.

WOOD ASHES, obtained by burning wood in heaps or pits, sheltered from the wind, contain *Potassic Carbonate*, the result of ignition of organic salts of Potassium, mixed with other salts, soluble and insoluble, and with organic matter.

- β. From WOOD ASHES are obtained, (1) by lixiviation and evaporation,—*Potashes*, i.e., Potassic Carbonate contaminated with other soluble salts, and with organic matter; (2) on igniting Potashes in a reverberatory furnace so as to burn off the organic impurities,—*Pearlash*, i.e., Potassic Carbonate contaminated only with other salts, soluble, but less so than itself; (3) on lixiviating Pearlash with *its own weight only* of distilled water, and decanting,—a nearly pure solution of **Potassæ Carbonas**, K_2CO_3 .

Evaporated to dryness,—crystalline powder.

7. From **Potassæ Carbonas** are prepared,

- 1.* By boiling in distilled water with slaked Lime, CaH_2O_2 , decanting, and diluting to a stated degree,—**Liquor Potassæ**; which, (*)on evaporation, leaves **Potassa Caustica**, KHO , in a fused state.

Cooled in moulds,—solid sticks.

- 2.* By passing Carbonic Anhydride, CO_2 , through a strong solution for about a week,—**Potassæ Bicarbonas**, KHCO_3 .

Deposited in crystals. More crystals separate on evaporating the mother liquor,—rhombic prisms.

- 3.* By neutralizing with Acetic Acid,—**Potassæ Acetas**, $\text{KC}_2\text{H}_3\text{O}_2$.

Evaporated to dryness, fused and solidified in mass,—foliaceous satiny masses.

- 4.* By neutralizing with Citric Acid,—**Potassæ Citras**, $\text{K}_3\text{C}_6\text{H}_5\text{O}_7$.

Evaporated to dryness, and granulated by stirring during the process; then powdered.

- 5.* By mixing with slaked Lime, moistening with water, and passing Chlorine gas through the mixture,—Calcic Chloride, Calcic Carbonate, and **Potassæ Chloras**, KClO_3 .

The Chloride and *Chlorate* are dissolved out by boiling water; the *latter* crystallizes out on cooling.

Purified by recrystallization,—rhomboidal plates.

- 6.* By treating with half its weight of Sulphur till effervescence (evolution of water and Carbonic Anhydride) ceases, and then fusing,—**Potassa Sulphurata**, a mixture of Potassic Sulphate and Sulphide, with some Sulphite, Sulphosulphate, and other Sulphur compounds.

Solidified in mass,—amorphous; grey.

7. By heating in a reverberatory furnace with Chrome Ironstone, $\text{FeO.Cr}_2\text{O}_3$, and a little chalk to keep the mass porous and prevent complete fusion,—*Potassic Chromate*, K_2CrO_4 . On dissolving out, and supersaturating the solution with Nitric Acid,—**Potassæ Bichromas**, $\text{K}_2\text{Cr}_2\text{O}_7$.

Crystallizes out on evaporation.

Purified by recrystallization. Four-sided tables, orange red.

8. As described on page 110,—**Potassæ Prussias Flava**.

8. From **Liquor Potassæ** are prepared,


- 1.* By the addition of Iodine,—a solution containing *Potassic Iodide*, KI , and *Iodate*, KIO_3 . On evaporating to dryness and fusing the dried salts with charcoal,—Carbonic Oxide is given off, and **Potassii Iodidum**, KI , remains.

Dissolved out, filtered, and crystallized. Opaque cubes.

Potassæ Chloras.



Potassæ Bichromas.



- 2.* By a similar process, using Bromine for Iodine,—**Potassii Bromidum**, KBr.

Cubes.

From **Potassa Caustica** are prepared,

- 1.* By semifusing with Black Oxide of Manganese, MnO_2 , and Potassic Chlorate (as an oxidizing agent),—*Potassic Manganate*, K_2MnO_4 , and Chloride. On dissolving out and adding Sulphuric Acid to the solution,—**Potassæ Permanganas**, $\text{K}_2\text{Mn}_2\text{O}_8$, Potassic Sulphate, and Manganic Peroxide, which is precipitated.

The *Permanganate* crystallizes out on evaporation.

Purified by dissolving, filtering through asbestos, and recrystallizing. Dark red needles.

2. By boiling its solution with Olive Oil (p. 42), which consists chiefly of *Olein*, i.e., Glyceric Oleate, $\text{C}_3\text{H}_5(\text{C}_{18}\text{H}_{33}\text{O}_2)_3$,—Glycerin, and **Sapo Mollis**, (*Soft Soap*), i.e., impure Potassic Oleate.

Washed from glycerine and water by a strong solution of Caustic Potash.

Soft gelatinous mass. Yellow.

ARGOL, or **CRUDE TARTAR**, the crust deposited on the interior of wine casks, consists of *Acid Potassic Tartrate* and Calcic Tartrate, with colouring and extractive matter.

- ε. From **ARGOL** is separated by lixiviation, and decolorized by filtration through charcoal and clay,—a solution of **Potassæ Tartras Acida** (*Acid Potassic Tartrate*, "*Cream of Tartar*,") $\text{KHC}_4\text{H}_4\text{O}_6$.

Crystallizes out on evaporation,—gritty powder, or cakes crystallized on the surface.

From a solution of **Potassæ Tartras Acida** in boiling water, is prepared, (*) by neutralizing with Potassic Carbonate,—one of **Potassæ Tartras**, $\text{K}_2\text{C}_4\text{H}_4\text{O}_6$.

Crystallizes out on evaporation,—four or six-sided prisms.



PROPERTIES.

*Solubility.**



	Parts by weight of water, at 60 deg. F.	Of water, at 212 deg. F.	Corresponding parts of Rect. Spirit.
P. Acetas is soluble in . . .	less than one	3
Potassa	$\frac{1}{2}$	$\frac{1}{2}$
P. Citras	less than one	1
P. Iodidum	" " "	16
P. Carbonas	" " "	—
P. Tartras	" " "	—

* The solubility of these, and of other metallic compounds, is given, for the most part, according to Squire, when his statements differ from those of other chemists.

Potassii Iodidum.



Potassæ Permanganas.



	Parts of water, at 60 deg. F.	Of water at 212 deg. F.	Of Rectified Spirit.
P. Bromidum	2	..	90
P. Bicarbonas	3	..	—
P. Nitras	4	2½	—
P. Sulphas	10	4	—
P. Chloras	16	2	—
P. Permanganas	16	(Decomposes)	(Decomposes)
P. Tartras Acida.	200	18	—

Diffusion Power.

All have a high rate of diffusion power, except

P. Sulphas, P. Tartras, P. Tartras Acida.

The following are *deliquescent*.

Potassa, P. Acetas, P. Carbonas, P. Citras.

Potassa Caustica, and in a less degree P. Carbonas, *corrode* organic matters by reason of their affinity for water or its elements, and their power of dissolving nitrogenous substances.

P. Permanganas *corrodes* by parting with some of its oxygen, and decomposing into Manganate and Manganic Peroxide.*

P. Permanganas is also deoxidized by Sulphurous Acid, by solutions of Sulphides, and by acidulated solutions of Sulphites, Sulpho-sulphates (Hyposulphites), Nitrites, Arsenites, Ferrous and Mercurous Salts.

Potassæ Chloras parts with Oxygen readily at 752 deg. F.; leaving Potassic Chloride and Perchlorate.† It parts with Oxygen to Sulphur or Phosphorus on being rubbed forcibly with it.

Treated with Hydrochloric Acid it evolves Chlorine.‡

P. Sulphas is a very hard substance, and is used to aid in triturating other drugs finely.

The following are *fusible*.

Potassa, P. Nitras, P. Chloras, P. Acetas, P. Iodidum, P. Bromidum.

The following are also *volatilizable*.

Potassa, P. Iodidum, P. Bromidum.

By *heat*,

- P. Bicarbonas is reduced to Carbonate.
- P. Carbonas „ Oxide (K_2O).
- P. Nitras „ Nitrite, then to Oxide and Peroxide (K_2O_2).
- P. Chloras „ Chloride and Perchlorate (see above), then to Chloride.
- P. Permanganas „ Manganate and Manganic Peroxide, giving off Oxygen.

P. Acetas, Citras, Tartras, and Tartras Acida, are reduced to Carbonate, blackened by products of incomplete combustion. In the case of the last named salt, the residue is known as "*black flux*".

P. Nitras and P. Chloras *deflagrate* on heating.

* * * Potassæ Bichromas is introduced only as an agent in making Sodæ Valerianas.


Potassæ Prussias Flava, as the source of Hydrocyanic Acid.

* $K_2Mn_2O_8 = K_2MnO_4 + MnO_2 + O_2$

† $2 KClO_3 = KCl + KClO_4 + O_2$. At a red heat, $-KClO_4$ gives $KCl + 2 O_2$.

‡ $KClO_3 + 6 HCl = KCl + 3 H_2O + Cl_6$.

Sapo Mollis.



SODIUM COMPOUNDS. 18 *Officinal*.

COMMON SALT (<i>Sodii Chloridum</i>). . . . <i>Crude Sodæ Sulphate</i> .	{	Sodæ Sulphas Sodæ Carbonas.	{	Sodæ Carbonas.—Sodæ Arsenias. Exsiccata.
				S. Bicarbonas.—S. Citro-Tartras Effervesens.
SODA NITRE. — Sodæ Nitras.	{		{	Soda. { Sapo Durus. Sapo Animalis. Sodæ Valerianas.
				Sodæ Acetas. Soda Tartarata. Sodæ Phosphas. HYPOSULPHITE OF SODA. Sodæ Hypophosphis. Soda Chlorata.
Borax.				

COMMON SALT (*Sodii Chloridum*), NaCl , found in solution in sea-water and brine-springs, and in solid deposits as rock-salt; separated from other substances contained in the first two by simple evaporation, from those contained in the last by solution and evaporation.

Crystallizes out in minute cubes.

- a. From COMMON SALT is prepared, by heating with sulphuric acid,—*Salt cake*, *i.e.*, crude Sodæ Sulphate; Hydrochloric Acid gas (p. 96) being given off.

On dissolving *Salt cake* in water and evaporating,—pure Sodæ Sulphas, $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$, crystallizes out.

Oblique rhombic prisms.

On heating *Salt cake* in a reverberatory furnace with chalk and carbon (coal dust),—(1) the sulphate is decoxidised to sulphide, and carbonic oxide is given off; (2) part of the chalk is reduced to quicklime; (3) the remainder undergoes mutual decomposition with the Sodæ Sulphide, and *Ball Soda* is formed; *i.e.*, a mixture of Sodæ Carbonate, Quicklime and Calcic Sulphide. On lixiviation, a solution of Sodæ Carbonate is obtained, but impure, as some Caustic Soda, NaHO , is formed by decomposition of the Carbonate with the Lime. On evaporation,—Sodæ Carbonas, $\text{Na}_2\text{CO}_3 \cdot 10\text{H}_2\text{O}$, separates.


Rhombic crystals.

- β. From Sodæ Carbonas are prepared—

- 1.* By heating till the water of crystallization (63 p. c. by weight) is driven off,—Sodæ Carbonas Exsiccata, Na_2CO_3 .

Powdered.

Sodæ Carbonas.



- 2.* By mixing with $1\frac{1}{2}$ part of **Sodæ Carbonas Exsiccata**, and passing Carbonic Anhydride into the mixture,—**Sodæ Bicarbonas**, NaHCO_3 .

Washed from carbonate by half its weight of distilled water, drained and dried.

Powder, or small scales.

- 3.* By a similar process to that of making *Liquor Potassæ* and *Potassa Caustica*,—**Liquor Sodæ** and **Soda Caustica**.

4. By a similar process to that of making *Potassæ Acetas*,—a solution of *Sodic Acetate*, $\text{NaC}_2\text{H}_3\text{O}_2$. On evaporation,—**Sodæ Acetas**, $\text{NaC}_2\text{H}_3\text{O}_2 \cdot 3\text{H}_2\text{O}$, crystallizes out.

Oblique rhombic prisms.

- 5.* By neutralizing with a solution of *Potassæ Tartras Acida*,—a *double Tartrate*, of Potassium and Sodium. On evaporation, **Soda Tartarata**, $\text{NaK C}_4\text{H}_4\text{O}_6 \cdot 4\text{H}_2\text{O}$, crystallizes out.

Right rhombic prisms.

- 6.* By neutralizing with a solution of *Acid Calcic Phosphate*, prepared as in the making of Phosphorus (p. 93), a solution of *Hydric Disodic Phosphate*, Na_2HPO_4 . On evaporation,—**Sodæ Phosphas**, $\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$, crystallizes out.

Rhombic prisms.

7. By passing Sulphurous Anhydride, SO_2 , into its solution, a solution of *Sodic Sulphite*, Na_2SO_3 ; Carbonic Anhydride being evolved. On digesting the Sodic Sulphite with Sulphur,—*Sodic Sulpho-sulphate* ("Hyposulphite"), $\text{Na}_2\text{S}_2\text{O}_3$, is formed. On evaporation, **HYPOSULPHITE OF SODA**, $\text{Na}_2\text{S}_2\text{O}_3 \cdot 5\text{H}_2\text{O}$, crystallizes out.

Striated rhombic prisms.

8. By adding it to a solution of *Calcic Hypophosphite*,—a solution of **Sodæ Hypophosphis**, NaPH_2O_2 ; and a precipitate of Calcic Carbonate.


The solution filtered, and evaporated to dryness. Granules.

- 9.* By passing Chlorine into solution of a stated strength,—**Liquor Sodæ Chloratæ**, containing Sodic Chloride, *Sodic Hypochlorite* (NaClO), and Bicarbonate.


Colourless. Odour of chlorine.

- γ. From **Sodæ Carbonas Exsiccata** is prepared, (*)by fusing with Arsenious Acid, As_2O_3 , and Sodic Nitrate,—*Sodic Pyro-arsenate*, $\text{Na}_4\text{As}_2\text{O}_7$, Carbonic and Nitrous Anhydrides being driven off. On


Sodæ Bicarbonas.



Soda Tartarata.



Liquor Sodæ Chloratæ.



cooling partially, and dissolving in boiling water,—*Hydric Disodic Arseniate*, Na_2HAsO_4 , is formed. On filtering and cooling,—**Sodæ Arsenias**, $\text{Na}_2\text{HAsO}_4 \cdot 7\text{H}_2\text{O}$. crystallizes out.

Prisms.

- δ. From **Sodæ Bicarbonas** is prepared, (*) by heating 17 parts with 8 of powdered Acidum Tartaricum and 6 of powdered Acidum Citricum, at a temperature of 200 deg. F., until the particles begin to aggregate, then stirring till granules are formed, and finally sifting,—**Sodæ Citro-Tartras Effervescens**, an intimate mixture of the drugs named.

Effervesces when placed in water, forming Sodic Citrate and Tartrate, and Carbonic Anhydride.

- ε. From **Soda** are prepared—

1. By a similar process to that of making Sapo Mollis (*Potassic Oleate*),—**Sapo Durus** (Castile Soap), chiefly *Sodic Oleate*.

Washed from glycerine by a solution of Sodic Chloride.

Solid mass, greyish white.

2. By a similar process, employing purified tallow, which contains 60 p. c. of Glyceric Stearate,—**Sapo Animalis** (Curd Soap), chiefly *Sodic Stearate*, $\text{NaC}_{18}\text{H}_{35}\text{O}_2$.

White.

- 3.* On distilling from a mixture of Amylic Alcohol, $\text{C}_5\text{H}_{12}\text{O}$, (p. 101) with Potassic Bichromate and diluted Sulphuric Acid,—*Valeric Acid*, $\text{C}_5\text{H}_{10}\text{O}_2$, *Amyl Valerate*, $\text{C}_5\text{H}_{11} \cdot \text{C}_5\text{H}_9\text{O}_2$, and water, pass over. On saturating the distillate with **Liquor Sodæ**, the alkali combines with the Valeric Acid and also decomposes the ethereal salt, forming **Sodæ Valerianas**, $\text{NaC}_5\text{H}_9\text{O}_2$, Amylic Alcohol, and water.

Cleared from the Amylic Alcohol, which floats on the surface, evaporated to dryness, fused, and cooled.

Amorphous masses. Odour of Valerian.

SODA-NITRE, a crude Sodic Nitrate, found native in Peru. Purified by solution and crystallization, twice repeated, it forms **Sodæ Nitras**, NaNO_3 .



Rhombohedral.

Borax (*Sodic Biborate*), $\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O}$, found native in Thibet and Persia.




Purified by solution and crystallization.

Six sided prisms.

Sodæ Arsenias.



Sodæ Valerianas.



PROPERTIES.

Solubility.

	Parts of water at 60° F.	Parts of water at 212° F.
Soda Caustica, soluble in . . .	1	..
S. Carbonas " . . .	2	..
S. Tartarata " . . .	2	1
S. Arsenias " . . .	2	..
Hyposulphite " . . .	2	..
S. Hypophosphis " . . .	2	..
S. Chloridum " . . .	2 $\frac{3}{4}$..
S. Sulphas " . . .	3	4
S. Phosphas " . . .	5	2
S. Valerianas " . . .	5	..
S. Bicarbonas " . . .	10	..
Borax " . . .	12	2

The following are soluble in Rectified Spirit—

Soda Caustica, S. Valerianas, S. Hypophosphis, Hyposulphite.
Sapo Durus. Sapo Animalis.

Borax is soluble in 1 part of Glycerine.

S. Hypophosphis is soluble in 2 parts of Glycerine.

Two only are *deliquescent*.

Soda Caustica. Sodæ Nitras.

Six are *efflorescent*.

S. Sulphas, S. Carbonas, S. Bicarbonas, S. Phosphas, Borax, S. Tartarata (slightly).

Solutions of S. Phosphas, S. Arsenias, and Borax have an *alkaline reaction*.

Effects of *heat*.

The following lose their water of crystallization—

Borax, S. Carbonas, S. Acetas, S. Phosphas, S. Tartarata, S. Sulphas.
S. Arsenias, Hyposulphite of Soda.

The first five undergo "*aqueous fusion*" in so doing, *i.e.*, the salt dissolves in the separated water.

Dehydrated Borax is light and porous, and is known as "*Calcined Borax*".

The following are *fusible*—

Borax, Soda Caustica, S. Acetas, S. Valerianas, S. Chloridum.
The Soaps are softened.

Strongly heated—

S. Nitras deflagrates.

S. Bicarbonas is reduced to (anhydrous) Carbonate*.

S. Acetas, S. Tartarata, S. Valerianas, and the Soaps are reduced to Carbonate, blackened by products of imperfect combustion.

S. Phosphas is reduced to Sodæ Pyrophosphate, $\text{Na}_4\text{P}_2\text{O}_7$.

S. Arsenias " " " Pyro-arsenate, $\text{Na}_4\text{As}_2\text{O}_7$.

The Sodæ Hypochlorite contained in Liq. Sodæ Chloratæ is decomposed into Chloride and Chlorate†.

* A solution of S. Bicarbonas loses Carbonic Acid on boiling.

† $3 \text{ NaClO} = 2 \text{ NaCl} + \text{NaClO}_2$.

8 parts of S. Sulphas with 5 of Acidum Hydrochloricum form a freezing mixture.

The properties of Liq. Sodæ Chloratæ are due to the Sodæ Hypochlorite, NaClO , which it contains.

In contact with organic matter, the Hypochlorite parts with its oxygen, and becomes a Chloride.

Exposed to air which contains Carbonic Anhydride, Sodæ Carbonate is slowly formed and Hypochlorous Acid liberated, which immediately decomposes, giving off free Chlorine.

In presence of a stronger acid, these changes take place rapidly.

S. Caustica, S. Carbonas, and their solutions, act on organic matters in the same way as the corresponding Potassium Compounds.

* * Sodæ Nitras is used only in making Sodæ Arsenias.

Sodæ Acetas ,, ,, ,, Ferri Arsenias and Ferri Phosphas

AMMONIUM COMPOUNDS. 9 *Officinal.*

AMMONIACAL LIQUOR.—	Ammonii Chloridum.	{	Ammoniæ Carbonas. — Ammoniæ Acetas.
	<i>or Ammonic Sulphate.</i>		
		{	Ammoniæ Citras.
			Ammoniæ Phosphas.
			Ammoniæ Nitras.
			Ammonii Bromidum.
			Ammoniæ Benzoas.

AMMONIACAL LIQUOR, a waste product of the manufacture of gas, contains Ammonia in combination with Carbonic and Hydrosulphuric acids.

- a. From AMMONIACAL LIQUOR is obtained, by neutralizing with Hydrochloric Acid,—a solution of **Ammonii Chloridum** (*Sal Ammoniac*), NH_4Cl , contaminated with tarry products; Carbonic Anhydride and Sulphuretted Hydrogen being given off.

Crystallizes out on evaporation.

Purified by sublimation,—tough fibrous masses.

(By a similar process, using Sulphuric Acid,—*Ammonic Sulphate*, $(\text{NH}_4)_2\text{SO}_4$).

- β. From **Ammonii Chloridum**, or *Ammonic Sulphate*, are obtained.

1. On subliming from a mixture of either salt with chalk (Calcic Carbonate),—*Ammonic Carbonate* $(\text{NH}_4)_2\text{CO}_3$; which is immediately decomposed by the heat necessary for sublimation, forming **Ammoniæ Carbonas**, a mixture of Acid Ammonic Carbonate with Ammonic Carbamate, $2\text{NH}_4\text{HCO}_3$. $(\text{NH}_3)_2\text{CO}_2$; and gaseous Ammonia.

Condenses in tough masses.

- 2*. On heating in a retort with slaked Lime,—gaseous *Ammonia*, NH_3 , which, carried over and dissolved in water to the point of saturation, forms **Liquor Ammoniæ Fortior**.

Liq. *Ammoniæ Fortior*,(*) diluted with twice its bulk of water, forms **Liquor Ammoniæ**.

Colourless liquids, evolving gaseous Ammonia abundantly.

- γ. Solutions of *Ammonic Acetate*, *Citrate*, *Phosphate*, *Nitrate*, *Bromide*,* and *Benzoate* are prepared by neutralizing

1. *Acetic Acid with *Ammoniæ Carbonas*.
2. *A solution of Citric Acid,
65 gr. to the oz. ... ,, Liq. *Ammoniæ Fortior*.
3. *Ac. Phosphoric Dil. ... ,, " *Liquor Ammoniæ*."
4. Ac. Nitric Dil. ... ,, " *Liquor Ammoniæ*."
5. Hydrobromic Acid ... ,, " " "
6. *Benzoic Acid ... ,, " " " (diluted).

* Solution of the Bromide is also prepared by adding *Ammoniæ Carbonas* to a solution of Ferrous Bromide, and filtering from the precipitate of Ferrous Carbonate. The solution of Ferrous Bromide is made by digesting Iron and Bromine in water.

The first solution, diluted to a stated degree, forms

Liq. Ammoniae Acetatis* ($\text{NH}_4\text{C}_2\text{H}_3\text{O}_2$).

The second forms

Liquor Ammoniae Citratis (NH_4)₃. $\text{C}_6\text{H}_5\text{O}_7$.)

Colourless liquids.

The fourth, evaporated to dryness, and the residue fused, yields *anhydrous Ammoniae Nitras*, NH_4NO_3 .

Confused crystalline masses.

From the third, fifth, and sixth, on evaporation, with a slight excess of Ammonia added from time to time—

Ammoniae Phosphas (NH_4)₂ HPO_4 ,—

Prisms.

Ammonii Bromidum NH_4Br ,—

Small crystals.

and **Ammoniae Benzoas** $\text{NH}_4\text{C}_7\text{H}_5\text{O}_2$,—

Laminae.

—crystallize out.

PROPERTIES.

Solubility.

	Parts of Cold Water.	Parts of Rect. Spirit.
Amm. Bromidum, soluble in	1½	13
Amm. Phosphas „	2	—
Amm. Chloridum „	4	55
Amm. Carbonas „	4	200
Amm. Benzoas „	5	18

Amm. Chloridum, Bromidum, Carbonas, Benzoas, *sublime* unchanged.

Amm. Phosphas is decomposed by heat.

Amm. Carbonas is decomposed by boiling water or alcohol.

Exposed to the air it loses its carbamate, leaving an opaque residue of *Bicarbonate* (“mild carbonate of Ammonia”) NH_4HCO_3 .

The evolved Carbamate decomposes into Ammonia and Carbonic Anhydride.

Ammonia in solution exercises an action on organic matters similar to that of Potassa and Soda.

* * * Ammoniae Nitras is used only as a source of Nitrous Oxide. For this reason it is required in the anhydrous form.

* * * Ammoniae Phosphas is a solvent for Urates of Soda and Lime.

* Contains a little Carbonic Acid in solution.

- 4.* By dissolving in diluted Hydrochloric Acid with the aid of heat,—a solution of *Ferrous Chloride*, FeCl_2 . On further heating with Nitric Acid and additional Hydrochloric,—a solution of *Ferric Chloride*, Fe_2Cl_6 , Nitric Oxide being given off.

a. On evaporation to a stated strength,—**Liquor Ferri Perchloridi Fortior.**

β . On diluting Liq. Ferr. Perchl. Fortior, with three times its bulk of distilled water, it forms **Liquor Ferri Perchloridi.**

γ . On diluting similarly with Rectified Spirit,—**Tinctura Ferri Perchloridi.**

All orange brown liquids.

- 5.* By macerating in Sherry for thirty days, the Iron being not wholly immersed, and the vessel frequently opened and shaken,—**Vinum Ferri**, containing a little *Ferric Tartrate*, *Malate*, etc.

Filtered. Brown.

β . From Ferri Sulphas are prepared—

- 1.* By heating until all the separable water of crystallization is driven off,—**Ferri Sulphas Exsiccata**, $\text{FeSO}_4 \cdot \text{H}_2\text{O}$.

Powdered. Whitey brown.

- 2.* By mixing its solution with one of Ammoniae Carbonas,—a precipitate of *Ferrous Carbonate*, FeCO_3 , with some *Ferric Oxide*. Collected, mixed with sugar, and dried,—**Ferri Carbonas Saccharata**, containing 37 p. c. of the carbonate.

Small lumps, brownish grey.

- 3.* By mixing its solution with one of Sodæ Phosphas and Sodæ Acetas,—a precipitate of **Ferri Phosphas**, $\text{Fe}_3(\text{PO}_4)_2$, some Acetic Acid being set free.



Collected, washed, dried at 120 deg. F.

Amorphous powder, slate blue.


- 4.* By a similar process, using Sodæ Arsenias instead of Phosphas,—**Ferri Arsenias**, $\text{Fe}_3(\text{AsO}_4)_2$.

Dried at 100 deg. F. Amorphous powder, pale green.

Ferri Perchloridi.




Ferri Phosphas.



Sodæ Phosphas being an acid salt, and Ferri Phosphas a neutral one, some free acid is produced by the decomposition. The presence of the Sodæ Acetate provides that the free acid should be Acetic, and not Sulphuric, in which Ferrous Phosphate is more soluble.

Ferri Arsenias.



The same applies here.

- 5.* By mixing its solution with $\frac{1}{11}$ part by weight of Sulphuric, and the same of Nitric Acid,—a solution of *Ferric Sulphate*, $\text{Fe}_2(\text{SO}_4)_3$, blackened by Nitric Oxide. On concentration, the Nitric Oxide is suddenly disengaged, and the liquid turns from black to red. Reduced to a stated strength, it forms **Liquor Ferri Persulphatis**.

Brownish red.

γ. From **Liquor Ferri Persulphatis** are prepared—

- 1.* By mixing with a solution of Potassic Acetate in Rectified Spirit, and shaking together,—a solution of *Ferric Acetate*, $\text{Fe}_2(\text{C}_2\text{H}_3\text{O}_2)_6$, Potassic Sulphate being precipitated. Filtered, and reduced to a stated strength, it forms **Tinctura Ferri Acetatis**.

Rose-red.

- 2.* By mixing with a solution of Potassic Sulphate,—a liquor from which, on evaporation, IRON ALUM ("Ferri Aluminas"), *Ferric Potassic Sulphate*, $\text{K}_2\text{SO}_4 \cdot \text{Fe}_2(\text{SO}_4)_3 \cdot 24\text{H}_2\text{O}$, crystallizes out.

Octahedra, pale violet.

- 3.* By pouring it into Liquor Sodæ, a precipitate of *Hydrated Ferric Oxide*, $\text{Fe}_2\text{O}_3 \cdot 3\text{H}_2\text{O}$. Collected, washed, and drained, it forms **Ferri Peroxidum Humidum**, about 46 p. c. of which is water mechanically engaged.

Brownish red.

*Dried at 212 deg. F., and powdered,—**Ferri Peroxidum Hydratum**, $\text{Fe}_2\text{O}_3 \cdot \text{H}_2\text{O}$.

Dark brown.

δ. From moist Ferric Oxide, precipitated, when required, by Liquor Ammoniaë instead of Liquor Sodæ, are prepared—

- 1.* By mixing with Potassic Acid Tartrate, standing for twenty-four hours, and heating to 140 deg. F.,—**Ferrum Tartaratum** (*Ferric Potassic Tartrate*), $\text{K}_3\text{Fe}(\text{C}_4\text{H}_4\text{O}_6)_3$.

Dissolved out, filtered, evaporated at 140 deg. F. to a syrupy consistence, spread on glass and dried at 100 deg.

Thin transparent scales, garnet colour.


- 2.* By dissolving in a solution of Citric Acid in excess, and neutralizing by Liquor Ammoniaë,—a solution of **Ferri et Ammoniaë Citras**.

Evaporated and dried as above. Similar aspect.

- 3.* By dissolving in a solution of Citric Acid in excess, then adding Quinia (precipitated from Quinia Disulphas by Ammonia), which also dissolves in the acid, and neutralizing by Liquor Ammoniaë,—a solution of **Ferri et Quinia Citras**.

Similarly treated. Similar scales, greenish yellow.

Ferri Persulphas.



- ε. From **Ferri Peroxidum Hydratum** is prepared, (*)by heating to strong redness in a gun barrel, and passing over it dried Hydrogen gas,—**Ferrum Redactum**, *i.e.*, reduced metallic Iron, with a variable amount of Magnetic Oxide.

Dark grey powder, strongly magnetic.

- ζ. A precipitate of **Ferri Oxidum Magneticum**, $\text{Fe}_3\text{O}_4 \cdot 4\text{H}_2\text{O}$, is obtained by (*)adding Liquor Sodæ to a mixture of Liq. Ferri Persulphatis with a solution of Ferrous Sulphate.

Washed, dried at 120 deg. F. Brownish black powder, strongly magnetic.

PROPERTIES.

F. et Amm. Citras is *soluble* in $\frac{1}{2}$ part by weight of water.

F. Quiniæ	“	“	$\frac{1}{2}$	“	“	“
F. Iodidum	“	“	1	“	“	“
Iron Alum	“	“	1	“	“	“
F. Sulphas	“	“	$1\frac{1}{2}$	“	“	“
F. Tartaratum	“	“	4	“	“	“

F. Tartaratum is sparingly soluble in Rectified Spirit; the other salts above named are practically insoluble in Rectified Spirit. Ferri

Sulphas is insoluble, even in Proof Spirit. (See “F. Sulphas Granulata”.)

Ferri Phosphas is soluble in Acids; Ferri Arsenias in Hydrochloric Acid. Reduced Iron, the Oxides, and the Carbonate are insoluble.

Liq. Ferri Persulphatis, L. F. Pernitratis, L. F. Perchloridi Fortior, are miscible in all proportions with Rectified Spirit.

Tinct. F. Acetatis in all proportions with water.

Exposed to air—

F. Sulphas becomes coated with a yellow insoluble oxysulphate.

F. Iodidum deliquesces.

In solution it slowly decomposes into Iodine and Ferric Oxide; depositing a rusty sediment and colouring the solution brown. A coil of iron wire extending throughout the vessel prevents this coloration, fresh iodide being continually formed.

Tinct. F. Acetatis deposits a gelatinous basic acetate.


By heat—

Ferri Sulphas is dehydrated.

Ferri Iodidum is decomposed, giving off Iodine and leaving Ferric Oxide.

* * Liquor Ferri Persulphatis is introduced only to prepare the Acetate and Peroxide.

Ferrum Redactum.



Ferri Oxidum Magneticum.



MERCURIAL COMPOUNDS. 11 *Officinal.*

Hydrargyrum.	{	Hydrargyri Iodidum Viride.*	{	Hydrargyri Subchloridum.* — <i>Mercurous Oxide.</i>	{	Hydrargyri Oxidum
		Hydrargyri Nitrates.				Flavum.
		Hydrargyri Oxidum Rubrum.				Hydrargyri Iodidum Rubrum.
	{	Hydrargyri Sulphas.		Hydrargyri Perchloridum.		Hydrargyrum Ammoniatum.

* *Mercurous* Salts, the rest being *Mercuric*.

Hydrargyrum,—Metallic Mercury, found native; or extracted from Cinnabar, a native Mercuric Sulphide, by simple roasting, or by distilling with Caustic Lime, which gives Mercury, Calcic Sulphide, and Calcic Sulphate.

Purified, if necessary, by redistillation.

a. From **Hydrargyrum** are prepared,

- 1.* By trituration with Iodine (moistening with a few drops of spirit to keep the mass cool), — **Hydrargyri Iodidum Viride** (*Mercurous Iodide*), Hg_2I_2 .

Dried in the dark. Greenish yellow powder.


- 2.* By dissolving in excess of diluted Nitric Acid, in stated proportions,—a solution containing *Mercuric Nitrate*, $\text{Hg}(\text{NO}_3)_2$, and Nitric Acid; Nitric Oxide being given off. On boiling to expel Nitric Oxide, it forms **Liquor Hydrargyri Nitratis Acidus**.

Colourless.


- 3.* By triturating with dry Mercuric Nitrate (prepared by dissolving Mercury as above and evaporating to dryness), and subsequently heating,—**Hydrargyri Oxidum Rubrum** (crystalline *Mercuric Oxide*), HgO ; Nitric Peroxide being evolved.

Crystalline powder, orange red.

Hydrargyri Nitrates.



Hydrarg. Oxidum Rubrum.



- 4.* By dissolving in Sulphuric Acid, with the aid of heat,—**Hydrargyri Sulphas**, HgSO_4 .

Evaporated to dryness. Crystalline powder, nearly white. Turns yellow on adding water.

- β. From **Hydrargyri Sulphas** are prepared,

- 1.* By mixing with Chloride of Sodium and metallic Mercury, and subliming into a large chamber,—**Hydrargyri Subchloridum** (*Mercurous Chloride*, or *Calomel*), Hg_2Cl_2 .

Washed from Mercuric Chloride (a little may be formed in the process), and dried at 212 deg. F.

Heavy powder, dull white.

- 2.* By mixing with Chloride of Sodium and Black Oxide of Manganese,* and subliming,—**Hydrargyri Perchloridum** (*Mercuric Chloride*, or *Corrosive Sublimate*), HgCl_2 .

Masses of prismatic crystals.

- γ. From **Hydrargyri Subchloridum** is prepared, (*)by adding 60 gr. to a pint of Liquor Calcis,—**Lotio Hydrargyri Nigra** (*Black Wash*), i.e., *Mercurous Oxide*, Hg_2O , suspended in a weak solution of Calcic Chloride.

- δ. From **Hydrargyri Perchloridum** are prepared,

- 1.* By adding 36 gr. to a pint of Liquor Calcis,—**Lotio Hydrargyri Flava** (*Yellow Wash*), a similar suspension of *Mercuric Oxide*.

- 2.* By pouring its solution into Liquor Sodæ,—a precipitate of **Hydrargyri Oxidum Flavum** (*Mercuric Oxide* in fine powder), HgO .

Separated, washed, dried at 212 deg. F. Fine yellow powder.

- 3.* By adding its solution to one of Iodide of Potassium,—a precipitate of **Hydrargyri Iodidum Rubrum** (*Mercuric Iodide*), HgI_2 .


Washed and dried, as above. Fine powder, brilliant red.

- 4.* By mixing its solution with Liquor Ammoniaë,—a precipitate of **Hydrargyrum Ammoniatum** (*Mercur-Ammonic Chloride*, or *White Precipitate*), NH_2HgCl ; Ammonic Chloride remaining in solution.


Washed and dried as above. White powder.

* The Black Oxide sets free a little chlorine, which prevents the accidental formation of Calomel.


Hydrarg. Subchloridum.



Hydrarg. Perchloridum.



Hydrarg. Ammoniatum.



PROPERTIES.

Hydrarg.-Perchloridum (Corrosive Sublimate) is *soluble* in 20 parts of water or 7 parts of Rectified Spirit. Also in ether.

Hydrarg. Iodidum Rubrum is soluble in ether, and slightly in Rectified Spirit. It is soluble in solutions of Corrosive Sublimate, of Iodide of Potassium, or of Chloride of Sodium.

The Oxides, the Green Iodide, the Subchloride (calomel) and White Precipitate, are insoluble in water, alcohol, or ether.

The Sulphate, in presence of water, forms a yellow basic sulphate.

Exposed to *light*,

The Green Iodide blackens.

When *heated* gently,

The Red Iodide turns yellow ;

The Green Iodide is decomposed into metallic Mercury and yellow Mercuric Iodide, which becomes red on cooling.

Metallic Mercury and its compounds are volatilized at a heat below redness. The Oxides and the Green Iodide are decomposed at the same time.

Metallic Mercury volatilizes slowly at ordinary temperatures. It boils at 662° F., and freezes at —39.

Sp. gr. of Metallic Mercury, 13.5.

Solutions of Corrosive Sublimate coagulate albumen, harden and contract animal tissues, and preserve them from putrefaction.

* * Hydrarg. Sulphas is introduced only to prepare the Chlorides.

CALCIUM COMPOUNDS. 11 *Officinal.*

BONE ASH

(Os Ustum). .. **Calcis Phosphas.**

CHALK

(Creta). .. { **Creta Præparata.**
Calcii Chloridum.—**Calcis Carbonas Præcipitata.**LIMESTONE. .. { **Calx.**—**Calcis Hydras.** { **Calcis Hypophosphis.**
Calx Chlorata.

MARBLE

(Marmor Album).

BONE ASH (*Os Ustum*), the residue left after burning bones in air, or igniting used animal charcoal, consists of *Tricalcic Phosphate* (**Calcis Phosphas**), $\text{Ca}_3(\text{PO}_4)_2$, with about 10 per cent. of Calcic Carbonate, both insoluble salts.

- a. * On digesting BONE ASH in dilute Hydrochloric Acid, a solution of Calcic Chloride and (soluble) *Calcic Superphosphate*, $\text{CaH}_4(\text{PO}_4)_2$, is obtained, Carbonic Anhydride being given off. On diluting, and adding Liquor Ammoniaë,—**Calcis Phosphas**, $\text{Ca}_3(\text{PO}_4)_2$, is precipitated; and Ammonic Chloride with some Calcic Chloride remains in the solution.

Collected, washed, dried at 212 deg. F. Amorphous powder.

- β. From CHALK (*Creta*), native friable Calcic Carbonate, contaminated with Ferrous Carbonate, are prepared,

1. By washing out from impurities (Elutriation), and drying,—**Creta Præparata** (Prepared Chalk).

Amorphous powder, aggregated into cones.

2. By adding it to Hydrochloric Acid, to neutrality,—a solution of **Calcii Chloridum**, CaCl_2 , contaminated with Ferrous Chloride. On adding a little Chlorinated Lime (*Calx Chlorata*, see below) and slaked Lime, the Ferrous is converted into a Ferric Salt and precipitated as Ferric Oxide, leaving the solution pure.




Filtered. Evaporated to solidity, the residue dried at 400 deg. F. Agglutinated masses.

- γ. From **Calcii Chloridum** is obtained, (*) on mixing its solution with one of Sodic Carbonate, a precipitate of *Calcic Carbonate*, CaCO_3 , which, when washed, and dried at 212 deg. F., forms **Calcis Carbonas Præcipitata**.

Crystalline powder.

- δ. From LIMESTONE, a native concrete Calcic Carbonate; is prepared, by ignition in a kiln,—**Calx** (Caustic Lime), CaO ; Carbonic Anhydride being driven off.

Compact masses.

Calcis Phosphas.

From **Calx**, (*) by addition of water ($\frac{1}{2}$ pint to the lb.),—**Calcis Hydras** (Slaked Lime), CaH_2O_2 , with evolution of heat.

Sifted. Amorphous powder.

ε. From **Calcis Hydras**,

- 1.* By heating with water and Phosphorus, a solution of **Calcis Hypophosphis**, $\text{Ca}(\text{H}_2\text{PO}_2)_2$; Phosphuretted Hydrogen being evolved.

Filtered. On passing Carbonic Anhydride, uncombined Lime separates as Carbonate.

Crystallizes out on evaporation. Pearly crystals.

2. By exposure to Chlorine Gas as long as it is absorbed,—**Calx Chlorata** (Bleaching Powder), a mixture of *Calcic Chloride* and *Hypochlorite*, $\text{Ca}(\text{ClO})_2$, with some unchanged lime.

Dull white powder.

* * Os Ustum, and Creta, are introduced merely as materials for preparing other drugs. Marmor Album (Marble) is introduced for the purpose of evolving Carbonic Anhydride.

PROPERTIES.

Calcii Chloridum is soluble in 2 parts of water, also in Rectified Spirit.

Calcis Hypophosphis in 8 parts of water; not in Rectified Spirit.

Calcis Hydras in about 900 parts of water; much more freely in syrup; not in Rectified Spirit.

The Carbonate and Phosphate are insoluble; the latter, with acids, yields a soluble Superphosphate (see above).


Exposed to air, **Calx** absorbs water, forming the Hydrate; **Calx** and **Calcis Hydras** and their solutions, absorb Carbonic Anhydride, forming the Carbonate.

Calcii Chloridum is fusible.


Calcis Hypophosphis is decomposed at a red heat.

The properties of **Calx Chlorata** are similar to those of **Soda Chlorata**, (see p. 119).

Calcis Hypophosphis.



Calx Chlorata.



ZINC COMPOUNDS. 8 *Officinal*.

Zincum.—Z.
Granulatum. . { **Zinci Chloridum.**
 { **Zinci Sulphas.** { **Z. Valerianas.**
 { **Z. Carbonas.** { **Z. Oxidum.**
 { **Z. Acetas.**

a. **Zincum** (commercial zinc) fused and poured into cold water, forms **Zincum Granulatum**, Zinc still contaminated with Iron and Tin.

β. From **Zincum Granulatum** are prepared,

1.* By dissolving in diluted Hydrochloric Acid, with the aid of heat,—a solution of *Zincic Chloride*, Zn Cl_2 , with a little Ferrous and Stannous Chlorides. On adding Chlorine Water, the impurities become Ferric and Stannic Chlorides. On stirring in a little insoluble Zincic Carbonate, they are precipitated as Ferric and Stannic Oxides, and a solution of *Zincic Chloride*, contaminated only by Chlorine, remains.

Filtered. On concentration to a stated strength (the Chlorine driven off),—**Liquor Zinci Chloridi**. Evaporated till only a fused salt remains,—**Zinci Chloridum**, Zn Cl_2 .

Solidified in moulds. Sticks.

2.* By similar processes, using Sulphuric Acid,—a pure solution of *Zincic Sulphate* Zn SO_4 . On evaporation, **Zinci Sulphas**, $\text{Zn SO}_4 \cdot 7\text{H}_2\text{O}$ crystallises out.

Small slender prisms.

γ. From boiling solutions of **Zinci Sulphas** are prepared,

1.* By mixing with a boiling solution of Sodæ Valerianas,—a solution of **Zinci Valerianas**, $\text{Zn (C}_5\text{H}_9\text{O}_2)_2$.

Crystals separate on cooling: more on evaporation at 200 deg. F. Pearly tables, odour of valerian.

2.* By mixing with a boiling solution of Sodæ Carbonate,—a precipitate of **Zinci Carbonas** (*Hydrated basic Zincic Carbonate*), $\text{ZnCO}_3 \cdot 2\text{ZnO} \cdot 3\text{H}_2\text{O}$; Carbonic Anhydride being evolved.

Separated by decantation, washed and dried. White powder.

δ. From **Zinci Carbonas** are prepared,

1.* By exposing to a dull red heat,—**Zinci Oxidum**, ZnO ; Carbonic Anhydride and water being driven off.



White powder; becomes yellow on heating.

2.* By adding to diluted Acetic Acid gently heated,—a solution of *Zincic Acetate*. $\text{Zn (C}_2\text{H}_3\text{O}_2)_2$. Boiled to expel Carbonic Anhydride, and filtered. On cooling,—**Zinci Acetas**, $\text{Zn(C}_2\text{H}_3\text{O}_2)_2 \cdot 2\text{H}_2\text{O}$, crystallizes out.

More crystals on evaporation of the mother liquor.

Thin crystalline plates.

CALAMINE, a native impure Zincic Carbonate, is not officinal.

Zinci Carbonas.

PROPERTIES.

Solubility.

		Parts of cold water.	Parts of boiling water.	Parts of Rect. Sp.
Zinci Chloridum	soluble in	less than one		—
Zinci Sulphas	„ „	2½	less than one	—
Zinci Acetas	„ „	2½	„ „ „	—
Zinci Valerianas	„ „	120	„ „ „	60

The Oxide and Carbonate are insoluble.

Zinci Chloridum is *deliquescent*.

Zinci Sulphas *effloresces* in dry air.

Heated gently,

Zinci Oxidum turns yellow.

Heated more strongly,

Zinci Sulphas and Acetas are dehydrated; the former undergoing
“watery fusion”.

Heated strongly,

Zinci Sulphas, Acetas, Valerianas, and Carbonas are decomposed, leaving
the Oxide.

Zinci Chloridum forms insoluble compounds with albumen and gelatine.

It accordingly preserves organic substances from putrefaction; and
applied to living tissues, produces an eschar.

LEAD COMPOUNDS. 6 *Officinal*.

LEAD. { **Plumbi Oxidum.**—**Plumbi Acetas.**—**Plumbi Subacetatis.**—**Plumbi**
[**Carbonas.**]
{ **Plumbi Nitras.**—**Plumbi Iodidum.**

a. From Metallic LEAD are obtained,

1. On heating in a current of air at a temperature sufficient to fuse the product,—**Plumbi Oxidum** (*Litharge*), PbO .

Heavy scales, brick-red.

2. On dissolving in diluted Nitric Acid,*—a solution of **Plumbi Nitras**, $\text{Pb}(\text{NO}_3)_2$.

Crystallizes out on evaporation. Octahedra.

β . From **Plumbi Oxidum** is prepared, (*)by dissolving in diluted Acetic Acid with the aid of heat,—a solution of *Plumbic Acetate*, $\text{Pb}(\text{C}_2\text{H}_3\text{O}_2)_2$. On evaporation (keeping slightly acid with Acetic Acid), **Plumbi Acetas** (*Sugar of Lead*), $\text{Pb}(\text{C}_2\text{H}_3\text{O}_2)_2 \cdot 3\text{H}_2\text{O}$, crystallizes out.

Masses of acicular crystals.

γ . From **Plumbi Acetas**, (*)by boiling in water with *Litharge*,—a solution of *Basic Plumbic Acetate*, $\text{Pb}(\text{C}_2\text{H}_3\text{O}_2)_2 \cdot 2\text{PbO}$. Diluted to a stated strength it forms **Liquor Plumbi Subacetatis**† (*Goulard's Extract*).

Colourless.

δ . From a solution of *Basic Plumbic Acetate* is obtained, on passing Carbonic Anhydride,‡ a precipitate of **Plumbi Carbonas**, $2\text{PbCO}_3 \cdot \text{PbH}_2\text{O}_2$ (*White Lead*) leaving Plumbic Acetate in the solution.

Heavy white powder.

ϵ . From **Plumbi Nitras** is obtained, (*)on mixing its solution with one of Iodide of Potassium, a precipitate of **Plumbi Iodidum**, PbI_2 .


Washed, dried at a gentle heat. Yellow powder.

* *Litharge* may be used instead of Lead.

† *Liq. Pb. Subacet. Dilutus* (*Goulard water*) contains 2 drachms of *Liq. Pb. Subacetatis* in the pint, with an equal amount of Rectified Spirit.

‡ In the "*Dutch Process*", preferred for making white lead to be used as a pigment, coils of metallic lead are placed in open jars containing weak vinegar, and the jars embedded in spent tan. The tan, putrefying, becomes heated, the acetic acid rises in vapour, the lead becomes oxidised by the air, and the oxide unites with the acetic vapour to form *basic acetate*. Finally, the carbonic anhydride from the putrifying tan precipitates a compact layer of *Carbonate* on the surface of the Lead, leaving Plumbic Acetate free to unite with oxide to form fresh *basic acetate*.

Plumbi Carbonas.



PROPERTIES.

Plumbi Acetas is soluble in $2\frac{1}{2}$ parts of water, also in Rectified Spirit.

Plumbi Iodidum is soluble in solutions of the Alkaline Iodides, or of Caustic Potash, in Acetic Acid, and slightly in Alcohol.

The Oxide and Carbonate are insoluble.

Plumbi Acetas, in dry air, effloresces, and is superficially decomposed by Carbonic Anhydride.

In solution, the Acetate is only partially decomposed by Carbonic Acid, the Acetic Acid liberated protecting the remainder.

When *heated*, Plumbi Acetas is first dehydrated and then decomposed.

„ „ Plumbi Iodidum is decomposed.

Solution of Plumbic Acetate has an acid reaction, of the Basic Acetate an alkaline one

Plumbic Acetate and Basic Acetate form insoluble compounds with albumen.

The latter forms an opaque jelly with mucilage.

* * Plumbi Nitras is used only to prepare the Iodide.

BISMUTH COMPOUNDS. 6 *Officinal.*

Bismuthum. — Bismuthum Purificatum. — <i>Bismuthous Nitrate.</i>	{	Bismuthi Subnitras. — Bismuthi Oxidum.
		Bismuthi Carbonas. Bismuthi et Ammoniaë Citras.

- a. **Bismuthum**, commercial BISMUTH, which contains Sulphur and Arsenic, being fused with Nitre, which removes the impurities, forms **Bismuthum Purificatum**.
- β. From **Bismuthum Purificatum** is prepared, (*)by dissolving in diluted Nitric Acid,—a solution of *Bismuthous Nitrate*, $\text{Bi}(\text{NO}_3)_3$.

The solution is filtered and concentrated.

- 1.* On pouring it into cold water, **Bismuthi Subnitras** (*Basic Bismuthous Nitrate*, “Magistery of Bismuth”), $\text{Bi}(\text{NO}_3)_3 \cdot \text{Bi}_2\text{O}_3 \cdot 3\text{H}_2\text{O}$,* is precipitated; an acid salt being left in the solution.

Washed; dried at 150 deg. F. White powder, minute scales.

- 2.* On pouring it into a cold solution of *Ammoniaë Carbonas*, —**Bismuthi Carbonas** (*Basic Bismuthous Carbonate*), $2(\text{Bi}_2(\text{CO}_3)_3 \cdot 2\text{Bi}_2\text{O}_3) \cdot 3\text{H}_2\text{O}$,† is precipitated.

Washed; dried at 150 deg. F. White powder.

- 3.* On adding a solution of Citric Acid (twice the weight of the Bismuth present), and neutralizing with *Liquor Ammoniaë*, a solution is obtained, containing a *double Citrate of Bismuth and Ammonia*, and an equivalent amount of Ammonic Nitrate. Diluted to a stated strength, it forms **Liquor Bismuthi et Ammoniaë Citratis**.

- γ.* On boiling **Bismuthi Subnitras** with *Liquor Sodæ*, a residue of **Bismuthi Oxidum**, Bi_2O_3 , is left.

Washed; dried at 212 deg. F. Lemon-yellow powder.

PROPERTIES.

Liq. Bism. et Amm. Citratis mixes with water in all proportions.

The other compounds are insoluble in water or alcohol.

They are blackened in air containing Sulphuretted Hydrogen.

The weight of Bism. Subnitras must be remembered in prescribing. It contains about $\frac{2}{3}$ by weight of Bismuth, the “Carbonas” only about $\frac{1}{3}$.

* Or $\text{Bi}_2\text{O}_3 \cdot 2\text{HNO}_3 \cdot \text{H}_2\text{O}$.

† Or $2(\text{Bi}_2\text{O}_3 \cdot \text{CO}_2) \cdot \text{H}_2\text{O}$.

ANTIMONIAL COMPOUNDS. 5 *Officinal*.

Antimonium Nigrum. ... { **Antimonium Sulphuratum.**
Antimonii Chloridum.—**Antimonii Oxidum.**—**Antimonium Tartaratum.**

Antimonium Nigrum (Black Antimony);—a native Antimonious Sulphide, Sb_2S_3 , purified from siliceous matter by fusion, and powdered. Greyish black, crystalline.

a. From **Antimonium Nigrum** are prepared—

1.* By boiling in Liquor Sodæ, a solution containing a soluble *double Sulphide*, and a soluble *double Oxide*, of Sodium and Antimony; Na_3SbS_3 , and Na_3SbO_3 . On gradual addition of diluted Sulphuric Acid,—*Antimonious Sulphide* in another form is precipitated, mixed with about 12 p.c. of *Antimonious Oxide*, Sb_2O_3 , forming **Antimonium Sulphuratum**; and Sodice Sulphate remains in the solution.

Washed; dried at 212 deg. F. Orange red powder.

2.* By heating in excess of Hydrochloric Acid,—an acid solution of *Antimonious Chloride* SbCl_3 . Filtered and concentrated, it forms **Liquor Antimonii Chloridi**.


Heavy liquid, yellowish red.

β.* On pouring **Liquor Antimonii Chloridi** into water,—*Antimonious Oxychloride*, $\text{SbCl}_3 \cdot \text{Sb}_2\text{O}_3$, is precipitated. The precipitate is collected and washed. On treating it with a solution of Sodice Carbonate for half an hour, **Antimonii Oxidum** (*Antimonious Oxide*), Sb_2O_3 , remains; Carbonic Anhydride having been evolved and Sodice Chloride retained in the solution.



Washed; dried at 212 deg. F. Greyish-white powder.

γ.* On making **Antimonii Oxidum** into a paste with Acid Potassic Tartrate, $\text{KHC}_4\text{H}_4\text{O}_6$, and leaving it for twenty-four hours,—*Potass-Antimonious Tartrate*, $\text{K}(\text{SbO})\text{C}_4\text{H}_4\text{O}_6$, is formed. On dissolving in boiling water, filtering, and evaporating the solution,—**Antimonium Tartaratum** ("Tartar Emetic"), $\text{K}(\text{SbO})\text{C}_4\text{H}_4\text{O}_6 \cdot \text{H}_2\text{O}$, crystallizes out.

More crystals on evaporating the mother liquor.
 Rhombic octahedra.

Antimonium Sulphuratum.

The equations in the brackets account for the formation of *Antimonious Oxide*. A little Sulphuretted Hydrogen is at the same time given off.

Antimonii Oxidum.**Antimonium Tartaratum.**

The compound (SbO) here plays the part of a monad. Sb_2O_3 may be considered as $(\text{SbO})_2\text{O}$.

PROPERTIES.

Tartar Emetic is *soluble* in 15 parts of water, insoluble in Rectified Spirit.

Antimonious Chloride is soluble in a small quantity of water, but is precipitated as an Oxychloride on dilution.

It is soluble in Hydrochloric Acid.

The Oxide and Sulphide are insoluble in water or spirit.

The Oxide is *fusible*.

The Sulphide, heated in air, forms Sulphurous Anhydride and Antimonious Oxide.

Tartar Emetic is dehydrated at 212 deg. F.; heated strongly is decomposed.

Liq. Antim. Chloridi has a similar action on organic tissues to that of Zinci Chloridum (p. 134).

ARSENICAL COMPOUNDS. 5 *Officinal*.

MISPICKEL.—**Acidum Arseniosum**. .. $\left\{ \begin{array}{l} \text{Potassic Arsenite.} \\ \text{Arsenic Trichloride.} \end{array} \right.$
Sodæ Arsenias.—**Ferri Arsenias**.

- a. In roasting MISPICKEL, a native Arseni-sulphide of Iron, FeAsS , as a preliminary to the extraction of the iron, Sulphurous Anhydride and *Arsenious Anhydride*, As_2O_3 , are formed. The latter is condensed in chambers through which the flues pass. Purified by sublimation, it forms commercial *Arsenious Acid* ("White Arsenic"). Resublimed,—it forms **Acidum Arseniosum**.

Stratified masses, or heavy white powder.

β. From **Acidum Arseniosum** are prepared—

- 1.* By dissolving in hot water with an equal weight of Potassæ Carbonas,—a solution of *Potassic Arsenite*; which, diluted and flavoured with Tinct. Lavandulæ Co., forms **Liquor Arsenicalis**.

Pink.

- 2.* By boiling in water with Hydrochloric Acid, a solution of *Arsenic Trichloride*, AsCl_3 ; which, diluted, forms **Liquor Arsenici Hydrochloricus**.
3. **Sodæ Arsenias** (see p. 120).
4. **Ferri Arsenias** (see p. 126).

PROPERTIES.

Acidum Arseniosum is soluble in 100 parts of cold water, in 20 parts of boiling water.

It is freely soluble in glycerine.

It sublimes at 400 deg. F.

Nitric Acid, and Iodine in presence of water, oxidize it to Arsenic Acid ($\text{As}_2\text{O}_3 + \text{I}_4 + 5 \text{H}_2\text{O} = 4 \text{HI} + 2 \text{H}_3\text{AsO}_4$).

Its solutions preserve organic matters from putrefaction.

MAGNESIUM COMPOUNDS. 5 *Officinal*.

DOLOMITE.	} Magnesiæ Sulphas.	{ Magnesiæ Carbonas.—Magnesia.
SEA WATER.		
ALUM-ORE.		
		{ Magnesiæ Carbonas Levis.—Magnesia Levis.

- a. DOLOMITE, or Magnesian Limestone, consists chiefly of Magnesian and Calcic Carbonates. On treatment with Sulphuric Acid,—*Magnesian* and Calcic *Sulphates* are formed. On dissolving,—a solution of *Magnesian Sulphate*, MgSO_4 , with a little Calcic Sulphate. On evaporation,—**Magnesiæ Sulphas** ("Epsom Salts"), $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, crystallizes out.

On evaporating the mother liquor of SEA-WATER, after separation of the Sodid Chloride, **Magnesiæ Sulphas** crystallizes out.

On evaporating the mother liquor of ALUM manufacture (p. 143), **Magnesiæ Sulphas** crystallizes out.

Minute rhombic prisms.

- β. From **Magnesiæ Sulphas** are prepared—

- 1.* By mixing a strong boiling solution with a similar solution of Sodid Carbonate, and evaporating to dryness,—**Magnesiæ Carbonas** (*Basic Magnesian Carbonate*), $3\text{MgCO}_3 \cdot \text{MgO} \cdot 5\text{H}_2\text{O}$.

Washed; dried at 212 deg. F. White granular powder.

- 2.* By mixing weak cold solutions of the same, and boiling for fifteen minutes,—**Magnesiæ Carbonas Levis** (*same composition*).

Washed; dried at 212 deg. F.

White powder, very light, partly amorphous, partly crystalline.

- γ. **Magnesiæ Carbonas** and **Magnesiæ Carbonas Levis**,(*) on calcination, yield **Magnesia** and **Magnesia Levis**, MgO , respectively.

White powders, the latter very light.

PROPERTIES.

Magnesiæ Sulphas is *soluble* in equal parts of water, insoluble in Rectified Spirit, precipitated by Rectified Spirit from strong aqueous solutions.

The Carbonate and Oxide are insoluble in water or spirit. The former is *soluble* in solutions of Carbonic Acid.

Magnesiæ Sulphas is *efflorescent*.

By *heat*,

Magnesiæ Sulphas is first dehydrated, undergoing aqueous fusion; then fused.

The Carbonate is reduced to **Magnesia** (see above).

ALUM.

ALUM ORE contains Iron Pyrites (Ferric Sulphide), FeS_2 , and *Alumina*, Al_2O_3 . On roasting the ore, or exposing it to the air, the Pyrites yields Ferrous Sulphide, FeS , and *Sulphuric Anhydride*, SO_3 . The latter forms, with the Alumina,—*Aluminic Sulphate*, $\text{Al}_2(\text{SO}_4)_3$; the former is oxidized to form Ferrous Sulphate.

On lixiviating, the two Sulphates dissolve out. On evaporating the solution, the Ferrons salt separates in crystals. On mixing the mother liquor, containing Aluminic Sulphate, with a concentrated solution of Ammonic Sulphate, **Alumen** (*Ammonic Aluminic Sulphate*), $\text{NH}_4\text{Al}(\text{SO}_4)_2 \cdot 12\text{H}_2\text{O}$, crystallizes out.

Re-dissolved and re-crystallized. Octahedra.

* On heating **Alumen** till all its water of crystallization is expelled, **Alumen Exsiccatum**, $\text{NH}_4\text{Al}(\text{SO}_4)_2$, remains.

White powder.

PROPERTIES.

Alumen is *soluble* in 18 parts of cold water, in less than its own weight of boiling water. It is insoluble in Rectified or even in Proof Spirit.

Its solutions have a strongly acid reaction, and dissolve zinc or iron with evolution of hydrogen.

Alumen Exsiccatum is insoluble, but on being moistened, it resumes its water of crystallization slowly, with evolution of heat.

Alumen *effloresces* slowly in air.

It undergoes "*aqueous fusion*" when heated. Strongly heated, it loses its acid.

Alum forms insoluble compounds with albumen, fibrin, casein, and gelatin.

Alumen Exsiccatum is used as an escharotic owing to this property, and also to its capacity for abstracting water.

LITHIUM COMPOUNDS. 2 *Officinal.*

- α . On adding a solution of Ammonia Carbonas to one of Lithic Sulphate or Chloride, a precipitate of **Lithiæ Carbonas**, L_2CO_3 , is obtained.
Washed, dried. White powder.
- β . * By dissolving **Lithiæ Carbonas** in a solution of Citric Acid, a solution of **Lithiæ Citras**, $\text{L}_3\text{C}_6\text{H}_5\text{O}_7$, is prepared.
Evaporated to visciduity. Dried at 240 deg. F., and powdered.
Amorphous powder, white.

PROPERTIES.

Lithiæ Citras is soluble in $2\frac{1}{2}$ parts of water ; Lithiæ Carbonas in 100 parts ; more readily in solutions of Carbonic Acid.

Lithiæ Citras is very deliquescent.

SILVER COMPOUNDS. 3 *Officinal*.

By dissolving **Argentum Purificatum**, pure metallic Silver, in diluted Nitric Acid with the aid of heat, a solution of **Argenti Nitras** ("lunar caustic"), AgNO_3 , is prepared.

Decanted. Crystallises out on evaporation.

Rhombic tabular prisms; or fused, and moulded into sticks.

On pouring a solution of **Argenti Nitras** into Liquor Calcis, a precipitate of **Argenti Oxidum**, Ag_2O , is obtained.

Washed; dried at 212 deg. F. Olive-brown powder.

PROPERTIES.

Argenti Nitras is soluble in half its weight of water, in 15 parts of Rectified Spirit.

Argenti Oxidum is very slightly soluble in water; readily soluble in Liq. Ammonia.

Argenti Nitras blackens when exposed to light, or when placed in contact with organic matter. Its solutions blacken under similar circumstances. It is reduced to metallic Silver in presence of Copper or Phosphorus.

Argenti Oxidum is liable to explode in contact with Creasote.

By *heat*, both are reduced to metallic Silver, the Nitrate fusing before reduction.

Argenti Nitras forms insoluble compounds with albumen and albuminoid substances. These compounds, at first white, blacken if exposed to light.

COPPER, CADMIUM, CERIUM.

On roasting **Cuprum**, pure Copper, *Cupric Oxide*, CuO , is formed. On adding Sulphuric Acid,—*Cupric Sulphate*, CuSO_4 . Dissolved in hot water. On evaporating, **Cupri Sulphas**, $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, crystallizes out.

Oblique Prisms, bright blue.

Soluble in 4 parts of water, insoluble in pure alcohol, slightly in Rectified Spirit.

Dehydrated by heat.

With albumen and albuminoids, its action is similar to that of Arg. Nitras.

Cadmii Iodidum, CdI_2 , is prepared by direct union of Cadmium and Iodine in presence of water.

Flat pearly crystals.

Insoluble in water or Rectified Spirit; fused by heat.

CERITE contains Iron, Cerium, and some other metals. On powdering, and mixing with Sulphuric Acid, Ferrous and *Cerous Sulphates* are formed. On roasting in a reverberatory furnace, the former is decomposed. By lixiviation, a solution of *Cerous Sulphate*, $\text{Ce}_2(\text{SO}_4)_3$, is obtained. On adding a solution of Ammonic Oxalate, **Cerii Oxalas** (*Cerous Oxalate*), $\text{Ce}_2(\text{C}_2\text{O}_4)_3 \cdot 9\text{H}_2\text{O}$, is precipitated.

Washed; dried at 212° deg. White Powder.

Insoluble in water or alcohol.

Dehydrated by heat.

Native **BLACK OXIDE OF MANGANESE** is introduced into the Pharmacopœia, for the purpose of evolving Chlorine from Hydrochloric Acid, as **Manganesii Oxidum Nigrum**.

Heavy black powder.

OFFICINAL PREPARATIONS OF METALS AND METALLIC COMPOUNDS.

Hydrargyrum cum Creta. Mercury triturated with twice its weight of
Light grey. Chalk till globules cease to be visible.

Pulvis Antimonialis. Antimonii Oxidum with twice its weight of Calcis
White. Phosphas.

Pulvis Cretæ Aromaticus. Powdered Cinnamon, Nutmeg, Saffron,
Yellow. Cloves, and Cardamoms, with Sugar and Chalk,—1 part of
Chalk in 4.

Pulvis Cretæ Aromat. cum Opio. The same, with one-fortieth part
of powdered Opium.

Liquor Ammoniae Fortior (see p. 123) :—16 grains of gaseous Ammonia
in the drachm.

„ **Ammoniae** :—The above, diluted to one-third the strength.

„ **Ammoniae Acetatis** (see p. 123). Contains Carbonic Acid in
solution, which improves the flavour.

„ **Ammoniae Citratis** (see p. 123).

Liquor Antimonii Chloridi (see p. 139) :—The terchloride dissolved in
Hydrochloric Acid.

Liquor Arsenicalis :—Contains Potassic Arsenite (see p. 141); coloured
Pink. with Tinct. Lavandulae Co. (*Fowler's Solution*).

„ **Arsenici Hydrochloricus** :—Contains the terchloride (see p. 141),
dissolved in Hydrochloric Acid.

Both the above are made with half a grain of Acid. Arseniosum
to the drachm.

Liquor Bismuthi et Ammoniae Citratis (see p. 138) :—3 grains of
Oxide of Bismuth in the drachm.

Liquor Calcis :—Saturated;—about $\frac{1}{18}$ grain of Calcis Hydras in the
drachm.

„ **Calcis Saccharatus** :—About 1 grain of Calcis Hydras in the
drachm, dissolved by the aid of sugar.

„ **Calcis Chloratae** :—Made with $5\frac{1}{4}$ grains of Calx Chlorata to the
drachm. Contains the soluble ingredients, the hypochlorite
and chloride.

Liquor Ferri Perchloridi Fortior (see p. 126) :—About 32 grains in the
Dark brown. drachm, with some free Hydrochloric Acid.

„ **Ferri Perchloridi** :—The same, diluted to one-fourth the strength,
Brown.

„ **Ferri Pernitratis** (see p. 125).
Reddish brown.

„ **Ferri Persulphatis** (see p. 127).
Dark brown.

Liquor Hydrargyri Nitratis Acidus (see p. 129):—Strongly acid.

„ **Hydrargyri Perchloridi**:— $\frac{1}{16}$ grain in the drachm, with the same amount of Chloride of Ammonium.

Liquor Magnesiae Carbonatis (Fluid Magnesia):—Freshly precipitated Magn. Carbonas dissolved in water impregnated with Carbonic Acid under pressure; 13 grains in the ounce. (To be kept corked.)

Liquor Plumbi Subacetatis (see p. 136):—16 grains in the drachm.

„ **Plumbi Subacet. Dilutus**:—The same diluted to one-eightieth the strength, with an equal part of Rectified Spirit:— $\frac{1}{8}$ grain (and $\frac{3}{4}$ min. of Rectified Spirit) in the drachm.

Liquor Potassæ (see p. 115):—Nearly $3\frac{1}{2}$ grains of Potassic Hydrate in the drachm.

„ **Potassæ Permanganatis**:— $\frac{1}{2}$ grain in the drachm.

Liquor Sodæ (see p. 119):—About $2\frac{1}{3}$ grains in the drachm. About the same saturating power as Liq. Potassæ.

„ **Sodæ Arseniatis**:— $\frac{1}{2}$ grain of *dehydrated* S. Arsenias in the drachm.

„ **Sodæ Chloratæ** (see p. 119).

Liquor Zinci Chloridi (see p. 134):—About 40 grains in the drachm.

(EFFERVESCENT.)

Liquor Potassæ Effervescens:—30 grains of Potassæ Bicarbonas in a pint of water, saturated with Carbonic Acid under a pressure of 7 atmospheres.

Liquor Sodæ Effervescens:—A similar preparation of Sodæ Bicarbonas.

Liquor Lithiæ Effervescens:—A similar preparation of Lithiæ Carbonas, 10 grains in the pint.

Liquor Magnesiae Citratis:—100 grains of Magnes. Carbonas dissolved in a solution of 200 grains of Citric Acid, and flavoured with Syrup of Lemons; made up to half a pint with water; 40 grains of Pot. Bicarb. added, and the bottle secured to retain the Carbonic Acid.

Tinctura Ferri Perchloridi:—Liquor F. Perchl. Fortior diluted with Brown. 3 parts of Rectified Spirit. Same strength as the Liquor. Contains free acid.

„ **Ferri Acetatis** (see p. 127).
Deep brown.

Spiritus Ammoniae Aromaticus:—Distilled from Ammoniae Carbonas, Liq. Ammon. Fortior, Volatile Oil of Nutmeg, and Oil of Lemon, with Rectified Spirit and Water.

Vinum Antimoniale :—Tartar Emetic in Sherry, $\frac{1}{4}$ grain in the drachm.
Brown.

Vinum Ferri (see p. 126):—Contains Ferric Tartrate, Malate, etc., in Sherry.
Brown.

,, **Ferri Citratis** :—F. et Amm. Citras in Orange Wine,—1 grain in the drachm.
Brown.

Syrupus Ferri Iodidi (see p. 125) :— $4\frac{1}{2}$ grains in the drachm.

Becomes decolorised on keeping, unless a coil of Iron wire is kept in it (*Squire*).

,, **Ferri Phosphatis** (see p. 126):—1 grain in the drachm.

Glycerinum Boracis :— $11\frac{1}{2}$ gr. in the drachm.

(1 oz. with 4 fl. oz. of Glycerine.)

Mel Boracis :—8 gr. in the drachm.

Yellow.

Pilula Aloes et Ferri :—Ferri Sulphas and Barbadoes Aloes, with Pulv. Cinnamomi Co. and Conf. of Roses; $\frac{3}{4}$ gr. of the Sulphate and 1 of the Aloes in 5 grains.

Pilula Ferri Carbonatis :—F. Carbonas Saccharata with Confection of Roses; 4 grains in 5.

,, **Ferri Iodidi** (see p. 125):—With sugar and Liquorice root; nearly $1\frac{1}{2}$ grain in 5.

Pilula Hydrargyri :—Metallic Mercury, with Confection of Roses and Liquorice Root; 1 grain in 3.

,, **Hydrargyri Subchloridi Co.** :—Calomel and Antim. Sulphuratum (1 grain of each in 5), with Guaiac Resin and Castor Oil.

Pilula Plumbi cum Opio :—Plumbi Acetas and Opium, with Confection of Roses; 3 grains of the Acetate, $\frac{1}{2}$ grain of the Opium in 4.

Pilula Saponis Co. :—Opium with Castile Soap and water; nearly 1 grain of the Opium in 5.

Trochisci Bismuthi :—2 grains of Bism. Subnitras in each.

Trochisci Ferri Redacti :—1 grain in each.

Trochisci Potassæ Chloratis :—5 grains in each.

Trochisci Sodæ Bicarbonatis :—5 grains in each.

Unguentum Antimonii Tartarati :—1 in 5.

Unguentum Cadmii Iodidi :—1 in 8.

Unguentum Hydrargyri :—Nearly 1 in 2 (of metallic Mercury).

Bluish grey.

,, **Hydrargyri Co.** :—About 1 in $4\frac{1}{2}$ (of metallic Mercury) with Camphor.

Bluish grey.

,, **Hydrargyri Ammoniati** :—1 in 8.

Unguentum Hydrargyri Iodidi Rubri:—1 in 28.

Bright scarlet.

„ **Hydrargyri Nitratis:**—1 in 15.

Lemon yellow.

„ **Hydrargyri Oxidi Rubri:**—1 in 8.

Dull red.

„ **Hydrargyri Subchloridi:**—1 in 6½.

Cream colour.

Unguentum Plumbi Acetatis:—1 in 37½.

„ **Plumbi Carbonatis:**—1 in 8.

„ **Plumbi Iodidi:**—1 in 8.

Orange.

„ **Plumbi Subacetatis Co.:**—1 in 20 (of Plumbic “Sub-acetate”), with Camphor.

Unguentum Potassii Iodidi:—Nearly 1 in 9.

„ **Potassæ Sulphuratæ:**—1 in 15½.

Greenish grey.

Unguentum Zinci:—1 in 6½ (of Zinci Oxidum).

Linimentum Ammoniaë:—Liquor Ammoniaë, with 3 parts of Olive-oil.
Creamy.

A species of *Ammonia Soap* is produced, which forms an emulsion with the rest of the oil.

Linimentum Calcis:—Equal parts of Liquor Calcis and Olive Oil.
Creamy.

A similar preparation.

Linimentum Hydrargyri:—Equal parts of Unguent. Hydrargyri, Liq. Creamy, Ammoniaë, and Lin. Camphoræ.
bluish grey. 1 part of Mercury in 6.

Linimentum Potassii Iodidi cum Sapone:—Potass. Iodidum, with Soft jelly. Castile Soap, Glycerine, and water; scented with Oil of Lemon.
1 of Iodide in 9.

Linimentum Saponis:—Castile Soap and Camphor dissolved in spirit; Pale yellow. scented with Rosemary.

Emplastrum Ferri:—Ferri Peroxid. Hydrat., with Burgundy Pitch and Red. Lead Plaster.

Emplastrum Hydrargyri:—Mercury triturated with oil in which Sulphur is dissolved, added to Lead Plaster.

„ **Ammoniaci cum Hydrargyro:**—The same, Ammoniacum being substituted for Lead Plaster.
Brownish grey.

Emplastrum Plumbi:—Lead-Soap; made by boiling Litharge (Pl. Oxidum) with Olive Oil and water.

„ **Plumbi Iodidi:**—Iodide of Lead, with Soap and Resin
Pale orange Plasters; 1 in 9.

Emplastrum Saponis:—Hard Soap, Resin, and Lead Plaster.

White.

,,

Brown.

Cerati Saponis:—Hard Soap boiled to dryness with a solution of Subacetate of Lead, made by boiling Litharge in vinegar; mixed with Bees-wax and Olive Oil by the aid of heat.

Mistura Cretæ:—Prepared Chalk, suspended by gum and syrup in Cinnamon water; about 13 grains in the fluid ounce.

Mistura Ferri Co:—A solution of Ferrous Sulphate in Rose-water, added to a milky fluid obtained by rubbing Potassic Carbonate, Myrrh, and sugar with Rose-water and Spirit of Nutmeg. Ferrous Carbonate is precipitated and suspended by the “resin soap” that the Potash (which is in excess) forms with the Myrrh.

Bluish green.

2½ grains of Ferri Sulphas to the fl. oz.

,, **Ferri Aromatica**:—An infusion of Cinchona, Calumba, and Cloves in Peppermint water, macerated on Iron wire, flavoured with Tr. Cardam. Co. and Tincture of Orange-peel.

Dark brown.

Contains very little iron.

Enema Magnesiae Sulphatis:—An ounce of Epsom Salts in 15 fl. oz. of Mucilage of Starch, with a fl. oz. of Olive Oil.

Suppositoria Hydrargyri:—5 grains of Ung. Hydrargyri (2½ of Bluish grey. Mercury) in each.

Suppositoria Plumbi Co. 3 grains of Pb. Acetas, and 1 of Opium in each.

Lotio Hydrargyri Flava (see p. 130):—Mercuric Oxide precipitated from Yellow. Corrosive Sublimate by Lime Water (18 grains to the half-pint).

,, **Hydrargyri Nigra** (see p. 130):—Mercurous Oxide similarly produced from Calomel (30 grains to the half-pint).

Cataplasma Sodæ Chloratæ:—A poultice made from 4 oz. of Linseed Meal, with 2 fl. oz. of Liq. Sodæ Chloratæ added.

Vapor Chlori:—The vapour arising from 2 oz. of Calx Chlorata (“bleaching powder”), moistened with water.

SYNOPSIS

OF THE

COMPOUND PREPARATIONS OF THE PHARMACOPŒIA,
AND THEIR INGREDIENTS.

POWDERS.

Pulv. Amygdalæ Co.

Sweet Almonds	3
Gum	1
Sugar	4

—
8**Pulv. Antimonialis.**

Oxide of Antimony	1
Phosphate of Lime	2

—
3**Pulv. Catechu Co.**

Catechu	4
Kino	2
Rhatany	2
Cinnamon	1
Nutmeg	1

—
10**Pulv. Cinnamomi Co.**

Cinnamon	1
Cardamoms	1
Ginger	1

—
3**Pulv. Cretæ Aromaticus.**

Cinnamon	4
Nutmeg	3
Saffron	3
Cloves	1½
Cardamoms	1
Sugar	25
Chalk	11

—
48½**Pulv. Cr. Ar. c. Opio.**

P. Cretæ Aromat.	39
Opium	1

—
40**Pulv. Elaterii Co.**

Elaterium	1
Sugar of Milk	9

—
10**Pulv. Glycyrrhizæ Co.**

Liquorice Root	1
Senna	1
Sugar	3

—
5**Pulv. Ipecacuanhæ Co. (Dover's
Powder.)**

Ipecacuanha	1
Opium	1
Sulphate of Potash	8

—
10**Pulv. Jalapæ Co.**

Jalap	5
Acid Tartrate of Potash	9
Ginger	1

—
15**Pulv. Kino Co.**

Kino	15
Opium	1
Cinnamon	4

—
20

Pulv. Opii Co.

Opium	3
Pepper	4
Ginger	10
Caraway	12
Tragacanth	1
			—
			30

Pulv. Rhei Co. (Gregory's Powder.)

Rhubarb	2
Light Magnesia	6
Ginger	1
			—
			9

Pulv. Scammonii Co.

Scammony	4
Jalap...	3
Ginger	1
			—
			8

Pulv. Tragacanthæ Co.

Tragacanth	1
Gum...	1
Starch	1
Sugar	3
			—
			6

CONFECTIONS.

Conf. Opii.

Pulv. Opii Co.	1
Syrup	...	(by weight)	3
			—
			4

Conf. Piperis.

Black Pepper	2
Caraway	3
Honey	15
			—
			20

Conf. Scammonii.

Scammony	2
Ginger	1
Oil of Caraway	$\frac{1}{2}$
„ Cloves	$\frac{1}{4}$
Syrup	2
Honey	1
			—
			$6\frac{1}{2}$

Conf. Sennæ.

Senna	$2\frac{1}{2}$
Coriander	1
Figs	4
Tamarinds	3
Cassia	3
Prunes	2
Liquorice (Extract)	$\frac{1}{4}$
Sugar	10
			—

After manufacture :—25

Conf. Sulphuris.

Sulphur	4
Acid Tartrate of Potash	1
Syrup of Orange-peel	4
			—
			9

Conf. Terebinthinæ.

Oil of Turpentine	1
Liquorice Root	1
Honey	2
			—
			4

The Confections of Roses and of Hips (Conf. Rosæ Gallicæ, Conf. Rosæ Caninæ) contain no ingredient besides the drug and the sugar.

PILL-MASSSES.

a. Aloetic. 9.**Pil. Aloes Barbadosensis.**

Barbadoes Aloes	2
Soap	1
Oil of Caraway	$\frac{1}{8}$
Confection of Roses	1
			—
			$4\frac{1}{8}$

Pil. Aloes et Ferri.

Barbadoes Aloes	2
Sulphate of Iron	$1\frac{1}{2}$
Pulv. Cinn. Co.	3
Confection of Roses	4
			—
			$10\frac{1}{2}$

Pil. Cambogiæ Co.

Gamboge	1
Barbadoes Aloes	1
Pulv. Cinn. Co.	1
Soap	2
Syrup	q.s.

6

Pil. Colocynthidis Co.

Colocynth	1
Barbadoes Aloes	2
Scammony	2
Sulphate of Potash	$\frac{1}{4}$
Oil of Cloves	$\frac{1}{4}$

 $5\frac{1}{2}$ **Pil. Col. et Hyoscyami.**

Pil. Colocynth. Co.	2
Extract of Henbane	1

3

Pil. Aloes Socotrinæ.

Socotrine Aloes	2
Soap	1
Oil of Nutmeg
Confection of Roses	1

4

Pil. Aloes et Assafoetidæ.

Socotrine Aloes	1
Assafoetida	1
Soap	1
Confection of Roses	1

4

Pil. Al. et Myrrhæ.

Socotrine Aloes	2
Myrrh	1
Saffron	$\frac{1}{2}$
Confection of Roses	$2\frac{1}{2}$

6

Pil. Rhei Co.

Rhubarb	3
Socotrine Aloes	$2\frac{1}{4}$
Myrrh	$1\frac{1}{2}$
Soap	$1\frac{1}{2}$
Oil of Peppermint
Treacle	4

 $12\frac{1}{4}$ **β. Opiate. 3.****Pil. Saponis Co.**

Opium	1
Soap*	4
Water	q.s.

5

Pil. Plumbi c. Opio.

Acetate of Lead	6
Opium	1
Confection of Roses	1

8

Pil. Ipecac. c. Scilla.

Dover's Powder (p. 152)	3
Squill	1
Ammoniac	1
Treacle	q.s.

7

γ. Squill.**Pil. Scillæ Co.**

Squill	$1\frac{1}{4}$
Ammoniac	1
Ginger	1
Soap	1
Treacle	2

 $6\frac{1}{4}$ **δ. Resinous. 2.****Pil. Assafoetidæ Co.**

Assafoetida	2
Galbanum	2
Myrrh	2
Treacle	1

7

Pil. Scammonii Co.

Scammony Resin	1
Jalap Resin	1
Curd Soap	1
Tr. Zingib. Fort.	1
Rectified Spirit	2

After manufacture:— $3\frac{1}{2}$ * Unless otherwise stated, "Soap" signifies *Castile Soap* (**Sapo Durus**).

ε. *Conium*.**Pil. Conii Co.**

Extract of Hemlock	...	5
Ipecacuanha...	...	1
Treacle	...	q.s.

7

ζ. *Mercurial*. 2.**Pil. Hydrargyri (Blue Pill).**

Mercury	...	2
Liquorice Root	...	1
Confection of Roses	...	3

6

Pil. Hydrarg. Subchloridi Co.
(*Plummer's Pill*.)

Calomel	...	1
Sulphd. Antimony	...	1
Guaiac Resin	...	2
Castor Oil	...	1

5

η. *Martial*. 2.**Pil. Ferri Carbonatis.**

Saccharated Carbonate	of	
Iron	...	4
Confection of Roses	...	1

5

Pil. Ferri Iodidi.

Iron	...	4
Iodine	...	8
Sugar	...	7
Liquorice	...	14
Water	...	5

After manufacture :—30

θ.

Pil. Phosphori.

Phosphorus	...	1
Tolu	...	60
Beeswax	...	30

91

Pil. Quiniæ.

Sulphate of Quinia...	...	3
Confection of Hips...	...	1
		4

MIXTURES.

a. *Emulsions and Suspensions*. 7.**Mist. Amygdalæ.**

Pulv. Amygd. Co. (p. 152)	...	1
Water	...	8

Mist. Ammoniaci.

Ammoniac	...	1
Water	...	32

Mist. Scammonii.

Scammony Resin	...	2 gr.
Milk	...	1 oz.

Mist. Guaiaci.

Guaiac Resin	...	2
Sugar	...	2
Gum	...	1
Cinnamon Water	...	80

* * Ammoniacum, being a gum-resin, emulsifies with water alone. The simple resins require the aid of an emulsifying agent (p. 53).

Mist. Cretæ.

Prepared Chalk	...	1
Gum	...	1
Syrup	...	2
Cinnamon Water	...	30

Mist. Ferri Co. (Griffith's Mixture.)

Ferri Sulphas	...	1½ gr.
Potassæ Carbonas	...	1½ gr.
Myrrh	...	3 gr.
Sugar	...	3 gr.
Spirit of Nutmeg	...	12 min.
Rose Water, to	...	1 oz.

Result :—Ferrous Carbonate and Myrrh, suspended by the aid of a "resin-soap" formed by the Myrrh and Potash.

Mist. Spiritus Vini Gallici
(*Egg Flip*.)

Egg Yolks	...	Two.
Sugar	...	¼ oz.
Brandy	...	4 oz.
Cinnamon Water...	...	4 oz.

*β. Solutions. 4.***Mist. Creasoti.**

Creasote	1 min.
Glacial Acetic Acid		1 min.
Spirit of Juniper...		2 min.
Syrup	30 min.
Water, to	1 oz.

Mist. Sennæ Co.

Epsom Salts ...	88 gr.
Extract of Liquorice	11 gr.
Tr. Cardam. Co. ...	$\frac{1}{2}$ drn.
Tincture of Senna	1 drn.
Infusion of Senna, to	1 oz.

Mist. Gentianæ.

Gentian.
Orange-peel.
Coriander.

A tincture of the above in proof spirit, diluted four times with water.

Mist. Ferri Aromatica.

Pale Cinchona.
Calumba.
Cloves.
Iron Wire.

An infusion of the above in Peppermint Water, with

Tincture of Orange-peel.
Tr. Cardam. Co.

Contains a minute proportion of organic salts of Iron.

EXTRACT.**Ext. Colocynthis Co.**

A spirituous extract of		
Colocynth,	6
to which is added		
Extract of Socotrine Aloes		12
Scammony Resin...	...	4
Soap	3
Cardamoms	1

After manufacture, 24

INFUSIONS.**Infus. Gentianæ Co.**

Gentian.
Orange-peel.
Lemon-peel.

Infus. Aurantii Co.

Orange-peel.
Lemon-peel.
Cloves.

Infus. Rosæ Acidum.

Red Rose Petals ...	11 gr.
Ac. Sulphuricum Dil.	6 min.
Water, to ...	1 oz.

DECOCTIONS.**Decoct. Aloes Co.**

Extract of Socotrine Aloes.
" " Liquorice.
Myrrh.
Saffron.
Carbonate of Potash.
Tr. Cardam. Co.

FLAVOURED PREPARATIONS.

Besides the above—

Infus. Catechu is flavoured with Cinnamon.

" **Lini** " Liquorice.
" **Sennæ** " Ginger.

Decoct. Hæmatoxyli ,, Cinnamon.

Decoct Sarsæ Co.

Sarsaparilla.
 Sassafras.
 Guaiac Wood.
 Mezereon.
 Liquorice.

TINCTURES.

Tr. Cardamomi Co.

Cardamoms.
 Caraway.
 Raisins.
 Cinnamon.
 Cochineal.

Tr. Chloroformi Co.

Chloroform	1
Tr. Cardam. Co.	5
Rectified Spirit	4

Tr. Lavandulæ Co.

Red Sandal Wood.
 Cinnamon.
 Nutmeg.
 Oil of Lavender.
 Oil of Rosemary (a trace).

Tr. Cinchonæ Co.

Pale Bark.
 Orange-peel.
 Serpentry.
 Saffron.
 Cochineal.

Tr. Gentianæ Co.

Gentian.
 Orange-peel.
 Cardamoms.

Tr. Benzoini Co. (Friar's Balsam.)

Benzoin	44 gr.
Storax	33 gr.
Socotrine Aloes	8 gr.
Tolu	11 gr.
				in the ounce.

Tr. Camphoræ Co. (Paregoric.)

Camphor	1½ gr.
Opium	2 gr.
Benzoic Acid	2 gr.
Oil of Anise	1½ min.
			in the ounce.

Tr. Catechu is flavoured with Cinnamon.

Tr. Aloes " Liquorice.

Tr. Quiniæ " Orange-peel; being made with Tr. Aurantii.

Tr. Sennæ " Raisins, Coriander, and Caraway.

Tr. Rhei " Coriander and Cardamoms, and coloured with Saffron.

Tr. Iodi contains Iodide of Potassium.

Vin. Opii is flavoured with Cinnamon and Cloves.

Vin. Aloes " Cardamoms and Ginger.

Vin. Rhei " Canella Alba.

Vin. Quiniæ and **Vin. Ferri Citratis**, with orange-peel, being made with Orange wine. The former contains free citric acid.

Syr. Rhamni is flavoured with Ginger and Pimento.

Syr. Rhei " } Coriander.

Syr. Sennæ " } Lemon-peel.

Syr. Ferri Phosphatis contains free Phosphoric Acid. **Syr. Scillæ** free Acetic Acid

Tr. Opii Ammoniata (*Scotch Paregoric*).

Opium	...	5 gr.
Saffron	...	9 gr.
Benzoic Acid	...	9 gr.
Oil of Anise	...	3 min.
Liq. Ammon. Fort.	96 min.	in the ounce.

*** For the Ammoniated Tinctures of Guaiac, Valerian, and Quinine, see page 6.

SPIRITS.

Sp. Armoraciæ Co.

Horse-radish.
Orange-peel.
Nutmeg.

Sp. Ammoniæ Fœtidus.

Assafoetida.
Liq. Ammon. Fort.

Sp. Amm. Aromaticus (*Sal Volatile*).

Amm. Carbonas.
Liq. Amm. Fort.
Oil of Nutmeg.
Oil of Lemon.

*** In each case the ingredients are distilled with Spirit.

LOZENGES.

Troch. Bismuthi.

Bismuthi Subnitras	...	2 gr.
Magnesiae Carbonas	...	2½ gr.
Calcis Carbonas Præcipitata	3½ gr.	
Rose water	...	q.s.
		in each lozenge.

Troch. Morphiae et Ipecacuanhæ.

Morphiae Hydrochloras	...	¾ gr.
Ipecacuanha	...	⅓ gr.
Tr. Tolutana.	...	a trace
		in each lozenge.

DILUTE ACID.

Ac. Sulphuricum Aromaticum.

Cinnamon.

Ginger.

macerated in

Ac. Sulphuricum...	3 parts.
Rectified Spirit	40 parts.

SOLUTIONS.

Liquor Calcis Saccharatus.

Slaked Lime	...	7 gr.
Sugar	...	44 gr.
		in the ounce.

Liquor Epispasticus.

Cantharides.
Acetic Acid.
Ether.

Troch. Opii are flavoured with Liquorice and Tolu.

Troch. Morphiae " }
Troch. Ac. Tannici " } Tolu.

Liquor Arsenicalis is coloured with Tr. Lavandulæ Co.

Liquor Bism. et Amm. Citr. is flavoured with Syrup of Lemons.

Liq. Arsenici Hydrochloricus.

Liq. Ferri Perchloridi (and L.F.P. Fort.).

Liq. Ferri Pernitratidis.

Liq. Hydrargyri Nitratis Acidus.

Liq. Morphiae Acetatis.

Liq. Morphiae Hydrochloratis.

Liq. Strychniæ.

} contain free acid.

Liq. Hydrarg. Perchloridi contains Chloride of Ammonium.

Liq. Iodi " Iodide of Potassium.

SUPPOSITORIES.

Suppositoria Plumbi Co.

Opium	1 gr.
Acetate of Lead	3 gr.
in each suppository.			

*** **Suppos. Hydrargyri** are made with Ung. Hydrargyri.

LINIMENTS.

a. *Strong Tinctures, made with Rectified Spirit, for painting on the skin.* 4.

To 1 fluid ounce of Rectified Spirit.

Lin. Aconiti.

Aconite Root	1 oz.
Camphor	22 gr.

Lin. Belladonnæ.

Belladonna Root	1 oz.
Camphor	22 gr.

Lin. Iodi.

Iodine	55 gr.
Iodide of Potassium	22 gr.
Camphor	11 gr.

Lin. Camphoræ Co.

Camphor	55 gr.
Liq. Amm. Fort.	2 drms.
Oil of Lavender	15 min.

b. *Oily or Saponaceous Mixtures for rubbing into the skin.* 12.

Lin. Ammoniaæ.

Liq. Ammoniaæ	1
Olive Oil	3

An Ammonia Soap is formed, which emulsifies the rest of the oil.

Lin. Calcis (Carron Oil).

Liquor Calcis.	} equal parts.	
Olive Oil.		

A similar reaction.

Lin. Camphoræ.

Camphor	1
Olive Oil	4

Solution takes place.

Lin. Crotonis.

Croton Oil	2
Oil of Cajeput	7
Rectified Spirit	7

Lin. Terebinthinæ.

Soft Soap	2
Camphor	1
Oil of Turpentine	16

Lin. Saponis (Opodeldoc).

Soap	50 gr.
Camphor	25 gr.
Oil of Rosemary	9 min.
Water	40 min.
Rectified Spirit, to	1 oz.

Lin. Pot. Iodidi cum Sapone.

Iodide of Potassium	66 gr.
Soap (<i>Castile or Curd</i>)	66 gr.
Glycerine	44 min.
Oil of Lemon	6 min.
Water, to	1 oz.

Lin. Sinapis Co.

Oil of Mustard	12 min.
Ethereal Extract	of		
Mezereon	8 gr.
Camphor	24 gr.
Castor Oil	1 drms.
Rectified Spirit, to	1 oz.

Lin. Chloroformi.

Chloroform	} equal parts.	
Camphor Liniment		

Lin. Hydrargyri.

Ung. Hydrargyri	} equal parts.	
(p. 160)		
Liquor Ammoniaæ		
Camphor Liniment		

Lin. Terebinth. Aceticum.

Oil of Turpentine	} equal parts.	
Acetic Acid		
Camphor Liniment		

Lin. Opii.

Laudanum	} equal parts.	
Soap Liniment		

OINTMENTS.

α.

Adeps Benzoatus. See p. 13.**Unguentum Simplex.**

White Wax	2
Lard	3
Almond Oil	3

β. *Made on a basis of Lard.* 10.**Ung. Aconitiæ.**,, **Atropiæ.**,, **Veratriæ.**,, **Belladonnæ.**

Made with the extract.

,, **Sabinæ.**

Contains beeswax.

,, **Iodi.**

Contains as much Iodide of Potassium as Iodine.

,, **Sulphuris Iodidi.**,, **Potassii Iodidi.**

Contains a little Pot. Carbonas.

,, **Potassæ Sulphuratæ.**,, **Hydrargyri Subchloridi.**γ. *On a basis of Benzoated Lard.* 5.**Ung. Sulphuris.**,, **Zinci.**

Oxide of Zinc.

,, **Plumbi Acetatis.**,, **Gallæ.**,, **Gallæ cum Opio.**δ. *On a basis of Simple Ointment.* 9.**Ung. Creasoti.**,, **Elemi.**,, **Resinæ.**

Contains beeswax.

,, **Antimonii Tartarati.****Ung. Cadmii Iodidi.**,, **Hydrarg. Iodidi Rubri.**,, **Hydrarg. Ammoniati.**,, **Plumbi Iodidi.**,, **Plumbi Carbonatis.**ε. *On a basis of Beeswax.* 1.**Ung. Picis Liquidæ.**ζ. *On a basis of Suet and Lard.* 1.**Ung. Hydrargyri** (*Blue Ointment*).η. *On a basis of Beeswax and Lard.* 1.**Ung. Terebinthinæ.**θ. *On a basis of Lard and Olive Oil.* 1.**Ung. Hydrargyri Nitratis** (*Citrine Ointment*).

Contains some free Nitric Acid.

ι. *On a basis of Beeswax and Olive Oil.* 2.**Ung. Cantharidis.**,, **Hydrargyri Co.** (*Scott's Dressing*).

Made with Ung. Hydrargyri and Camphor.

κ. *On a basis of Beeswax and Almond Oil.* 1.**Ung. Hydrarg. Oxidi Rubri.**λ. *On a basis of White Wax and Almond Oil.* 2.**Ung. Cetacei.**,, **Plumbi Subacetatis Co**
Contains Camphor.

PLASTERS.

α.

Empl. Plumbi (*Diachylon*).Lead Soap, *i.e.*, Oleate and Margarate of Lead (see p. 150).**Empl. Cerati Saponis.**

Acetate of Lead.

Soap.

Beeswax.

Olive Oil.

Empl. Ammoniaci c. Hydrargyro.

Ammoniac.

Mercury.

Olive Oil.

Sulphur.

Empl. Picis.

Burgundy Pitch.

Frankincense.

Rosin.

Beeswax.

Nutmeg Butter.

Olive Oil.

Empl. Cantharidis (*Blister*).

Cantharides.

Beeswax.

Suet.

Lard.

Rosin.

β. *Made on a basis of Lead Plaster.***Empl. Ferri.***With Peroxide of Iron.*

Burgundy Pitch.

Empl. Hydrargyri.*With Mercury.*

Olive Oil.

Sulphur.

Empl. Galbani.*With Galbanum.*

Ammoniac.

Beeswax.

Empl. Saponis. }**Empl. Resinæ.** }*With Soap.*

Rosin.

The former with six times as much Soap as Rosin; the latter with only half as much.

γ. *On a basis of Rosin Plaster.* 2.**Empl. Belladonnæ.***Made with equal parts of a Spirituous Extract of Extr. Belladonnæ.***Empl. Opii.***With powdered Opium, 1 in 10.*δ. *On a basis of Soap and Rosin Plasters.* 2.**Empl. Calefaciens.***With Cantharides.*

Rosin.

Beeswax.

Nutmeg Butter.

Empl. Plumbi Iodidi.

PAPERS.

Charta Sinapis.

Black Mustard Seeds in powder.

Gutta Percha.

Charta Epispastica.

Cantharides.

Canada Balsam.

Rosin.

Spermaceti.

Olive Oil.

White Wax.

* * **Hydrargyrum cum Creta** (*Grey powder*) :—see page 147.**Lotio Hydrargyri Flava** (*Yellow wash*) } see page 130.**Lotio Hydrargyri Nigra** (*Black wash*) }**Collodium Flexile** :—see page 111.

POPULAR DESIGNATIONS OF CERTAIN PHARMACOPŒIAL PREPARATIONS AND COMPOUNDS.

Aromatic Confection	of	Pulv. Cretæ Aromaticus.
Bhang ; Gunjah ; Hashish	,,	Cannabis Indica ; and its extracts.
Black Wash	,,	Lotio Hydrargyri Nigra.
Blue Pill,—Ointment... ..	,,	Pil.,—Ung. Hydrargyri.
Calomel	,,	Hydrargyri Subchloridum.
Carron Oil	,,	Linim. Calcis.
Chloric Ether	,,	Sp. Chloroformi.
Citrine, or Golden, Ointment... ..	,,	Ung. Hydrargyri Nitratis.
Corrosive Sublimate	,,	Hydrargyri Perchloridum.
Cream of Tartar	,,	Potassæ Tartras Acida.
Diachylon	,,	Emplastrum Plumbi.
Dover's Powder	,,	Pulv. Ipecac. Co.
Epsom Salts	,,	Magnesiæ Sulphas.
Flowers of Sulphur	,,	Sulphur Sublimatum.
Fowler's Solution	,,	Liq. Arsenicalis.
Friar's Balsam... ..	,,	Tr. Benzoini Co.
Glauber's Salt... ..	,,	Sodæ Sulphas.
Granular Effervescent Citrate of Magnesia	,,	Sodæ Citro-Tartras Effervescens.
Goulard's Extract	,,	Liq. Plumbi Subacetatis.
Goulard Water	,,	Liq. Plumbi Subacetatis Dil.
Gregory's Powder	,,	Pulv. Rhei Co.
Grey Powder	,,	Hydrargyrum cum Cretâ.
Griffith's Mixture	,,	Mist. Ferri Co.
Hartshorn	,,	Ammonia.
Laudanum	,,	Tr. Opii.
Lenitive Electuary	,,	Confect. Sennæ.
Litharge	,,	Plumbi Oxidum.
Liver of Sulphur	,,	Potassa Sulphurata.
Lunar Caustic... ..	,,	Argenti Nitras.
Milk of Sulphur	,,	Sulphur Præcipitatum.
Mindererus Spirit	,,	Liq. Ammonię Acetatis.
Nitre ; Saltpetre	,,	Potassæ Nitras.
Oil of Male Fern	,,	Extract. Filicis Liquidum.
Oil of Vitriol	,,	Acidum Sulphuricum.
Opodeldoc	,,	Lin. Saponis.

Paregoric Elixir	of	Tr. Camphoræ Co.
Plummer's Pill	,,	Pil. Hydrarg. Subchloridi Co.
Prussic Acid	,,	Ac. Hydrocyanicum.
Rochelle Salt	,,	Soda Tartarata.
Sal Ammoniac...	,,	Ammonii Chloridum.
Sal Volatile	,,	Sp. Ammoniæ Aromaticus.
Scotch Paregoric	,,	Tr. Opii Ammoniata.
Scott's Ointment	,,	Ung. Hydrargyri Co.
Sugar of Lead	,,	Plumbi Acetas.
Sweet Spirits of Nitre	,,	Sp. Ætheris Nitrosi.
Tartar Emetic...	,,	Antimonium Tartaratum.
Vitriol: Blue, Green, White...	,,	Cupri,—Ferri,—Zinci Sulphas.
White Arsenic...	,,	Acidum Arseniosum.
White Precipitate	,,	Hydrargyrum Ammoniatum.
Yellow Wash	,,	Lotio Hydrargyri Flava.

Substitutes for Patent Remedies are introduced in Pulvis Antimonialis, for "James's Fever Powders," and in Confectio Piperis, for "Ward's Paste."

LATIN PHRASES EMPLOYED IN PRESCRIBING.

The following are among those in most frequent use.

R, recipe. M, misce take. Mix.
Aa., ana, singulorum (—arum) of each.
Partes æquales equal parts.
Q. s., quantum sufficiat a sufficiency.
F., fiat haustus statim sumendus ...	make a draught to be taken at once.
Cujus (quorum, quarum, e quibus) sum-	
a(n)tur, capia(n)tur... of which let there be taken.
Cochleare, C. modicum, amplum a tea-, dessert-, table-spoon.
Tertiâ quâque horâ, nocte every third hour, night.
Omni, quoque die. Alternis diebus...	every day. Every other day.
Quotidie. Semel, bis, ter, quater die.	daily. Once, twice, thrice, four times
	daily.
Cras manc. Mane primo. Vespere...	to-morrow morn. The first thing in
	the morning. In the evening.
Sequenti nocte. Horâ somni the following night. At bed-time.
Post horas sex, si opus sit six hours after, if necessary.
P. r. n., pro re natâ as circumstances arise.
Alvo adstrictâ. Tusse molestâ when (if) the bowels be confined, the
	cough be troublesome.
Donec alvus deiciat until the bowels act.
Repetatur. Sæpe utendus repeat. To be used frequently.
Ejusdem mitte... send of the same.
Dimidium. Ss., Semis a half. Half.
Semihora. Uncia cum semisse half an hour. An ounce and a half.
Quadrans. Tertia, quinta pars a quarter. A third, fifth part.
Divide in partes duodecim divide into twelve parts.
Deaurentur pilulæ gild the pills.
Aqua fervens, calida, tepida, frigida...	boiling, hot, warm, cold water.
Applicetur lotio; —ntur hirudines	
undecim apply the lotion, eleven leeches.
Partibus affectis. Lateri dolenti to the affected parts, the painful side.
Femori, brachio interno, externo to the inner, outer side of the thigh,
	arm.
Tempori dextro, sinistro to the right, left temple.
Regioni hepatis, cordis to the region of the liver, heart.
Injiciatur sub cute inject subcutaneously.
Modo præscripto as directed.
Signa label as follows.

SYNOPSIS OF THE DOSES OF THE PHARMACOPŒIAL
DRUGS AND PREPARATIONS,
AND OF THE LEADING NON-OFFICINAL DRUGS.*

α. OF THE DRUGS.

Phosphorus	$\frac{1}{10}$ to $\frac{1}{10}$ grain.
Iodine	$\frac{1}{2}$ grain, gradually increased.	
Charcoal and Sulphur	a scruple to a drachm.	
<i>Iodoform</i>	1 to 5 grains.
<i>Croton-Chloral Hydrate</i>	1 to 10 grains.
Chloral Hydrate	5 to 30 grains.
<i>Nitroglycerin</i>	$\frac{1}{50}$ to $\frac{1}{20}$ minim.
Creasote	1 to 3 minims.
Amyl Nitrite (<i>inhaled</i>)	2 to 5 minims.
Chloroform	3 to 10 minims.
Ether and Acetic Ether	20 minims to a drachm.	
Glycerine	1 to 2 drachms.

Acids.

Arsenious	$\frac{1}{60}$ to $\frac{1}{12}$ grain.
Carbolic	1 to 3 grains.
Gallic and Tannic	2 to 10 grains.
Benzoic	10 to 15 grains.
Citric, Tartaric, and <i>Salicylic</i>	10 to 30 grains.
Dilute Hydrocyanic	2 to 8 minims.
Dilute Nitrohydrochloric	5 to 20 minims.
Dilute Sulphuric, Nitric, Hydrochloric and Phosphoric, and Aromatic Sulphuric	5 to 30 minims.
Sulphurous	$\frac{1}{2}$ to 1 drachm.
Dilute Acetic, also Vinegar and Oxymel	1 to 2 drachms.

Lactic Acid is chiefly used as a spray, diluted 8-16 times. Sometimes internally in $\frac{1}{2}$ —1 drachm doses.

* The names of the non-officinal drugs are printed in *italics*.

The Metals and

Arseniates	$\frac{1}{16}$ to $\frac{1}{2}$ grain.
Valerianates	1 to 5 grains.
Hypophosphites	5 to 10 grains.
Salicylates	10 to 60 grains.

Potassium Compounds 10 to 30 grains.
($\frac{1}{2}$ scruple to $\frac{1}{2}$ drachm.)

Except the following :—

The Permanganate and Potassa Sulphurata	... 1 to 5 grains.
„ Iodide 1 to 10 grains.
„ Bromide 5 to 30 grains.
„ Acetate, Citrate, and Tartrates	... 20 to 60 grains.

Lithium Compounds.

The Carbonate	3 to 6 grains.
„ Citrate	5 to 10 grains.

Iron and its Compounds 1 to 5 grains.

Except the following :—

The Dehydrated Sulphate	$\frac{1}{2}$ to 3 grains.
„ Magnetic Oxide, Phosphate, Tartrated Iron } and the Double Citrates	5 to 10 grains.
„ Hydrated Peroxide	5 to 30 grains.
„ Moist Peroxide	2 to 4 drachms.

The Arseniate:—see above.

Alum ; Bismuth and Calcium Compounds ... 5 to 20 grains.

Except—

Sulphide of Calcium	$\frac{1}{4}$ to 1 grain.
Chalk, and Precipitated Carbonate of Lime	... 10 to 60 grains.

Hypophosphite of Lime:—see above.

Salts used as Purgatives.

Calomel	2 to 5 grains.
Alum	1 scr. to 1 drachm.
Sulphate of Potash	1 scr. to 2 drachms.
Sulphate of Magnesia	1 to 4 drachms.
Tartrates of Potash, and Tartrated Soda ...	2 to 4 drachms.
Sulphate of Soda, Acetate of Potash, and Phos- phate of Soda	2 to 8 drachms.

Metallic Compounds.

Arsenious Acid	$\frac{1}{60}$ to $\frac{1}{12}$ grains.
Nitrate of Silver	$\frac{1}{4}$ to $\frac{1}{2}$ grain.
Oxide of Silver, Iodide of Lead, Acetate and Chloride of Zinc, Oxalate of Cerium, and Sulphate of Copper	$\frac{1}{4}$ to 2 grains.
Acetate of Lead	1 to 4 grains.

Sodium and Magnesium Compounds ... 20 to 60 grains.
(*A scruple to a drachm.*)

Except the following Sodium Compounds:—

The Dehydrated Carbonate...	3 to 10 grains.
„ Carbonate	5 to 30 grains.
„ Phosphate and Citro-tartrate	1 to 2 drachms.

The Arseniate, Valerianate, Hypophosphite, *Salicylate*:—see above.

Ammonium Compounds ... 5 to 20 grains.

Except the following:—

The Carbonate	3 to 10 grains.
„ Bromide	2 to 20 grains.

The *Salicylate*:—see above.

Mercury and its Compounds.

The Red Iodide, and Corrosive Sublimate	$\frac{1}{16}$ to $\frac{1}{4}$ grain.
The Green Iodide	1 to 3 grains.
Calomel <i>as an alterative</i>	$\frac{1}{2}$ to 1 grain.
Calomel <i>as a purgative</i>	2 to 5 grains.
Hydrarg. cum Cretâ...	3 to 8 grains.

Antimonial Compounds.

Tartar Emetic <i>as a diaphoretic</i>	$\frac{1}{16}$ to $\frac{1}{8}$ grain.
„ „ „ <i>depressant</i>	$\frac{1}{6}$ to 2 grains.
„ „ „ <i>emetic</i>	1 to 5 grains.
Oxide of Antimony and Sulphurated Antimony	1 to 5 grains.

Salts used as Emetics.

Tartar Emetic	1 to 5 grains.
Sulphate of Copper	5 to 10 grains.
Acetate of Zinc	10 to 20 grains.
Sulphate of Zinc	10 to 30 grains.
Carbonate of Ammonia	30 gr. freely diluted.

Animal Drugs.

Pepsin	2 to 5 grains.
Oxgall, Castor, and Musk	5 to 10 grains.
Spermaceti	1 scr. to 1 drachm.
Cod-Liver Oil	1 drm. to 1 oz.

Vegetable Drugs.

<i>Ignatia</i>	$\frac{1}{8}$ to 1 grain.
Capsicum	$\frac{1}{2}$ to 1 grain.
Digitalis and Ipecacuanha	$\frac{1}{4}$ to 2 grains.
Squill	1 to 2 grains.
Nux Vomica	1 to 3 grains.
Calabar Bean	1 to 4 grains.
Rhubarb	1 to 5 grains.
Conium, Colocynth, Colchicum	2 to 8 grains.
Savin	5 to 10 grains.
Quassia, Calumba, Gentian, and most Aromatics	5 to 20 grains.
Jalap, Ipecacuanha as an emetic, Rhubarb as a purgative...	10 to 30 grains.
Ergot	20 to 30 grains.
Cinchona Bark	20 to 60 grains.
Cusso, Cubebs, Wurrus, and <i>Guarana</i>	1 to 2 drachms.
Mustard as an emetic, and Cassia	2 drms upwards.

Resins, Gum-resins, Oleo-resins, Balsams, Gums.

Gamboge, and Jalap-resin (or " <i>Jalapine</i> ," p. 88)	1 to 5 grains.
<i>Red Gum</i>	3 to 6 grains.
Scammony-resin	3 to 8 grains.
Scammony	5 to 10 grains.
Assafoetida, Storax, and Tolu	5 to 20 grains.
Ammoniac, Benzoin, Galbanum, Guaiac, and Myrrh	10 to 30 grains.
Mastic	20 to 40 grains.
Tragacanth	20 grains upwards.
Balsam of Peru	10 to 15 minims.
„ Copaiba	$\frac{1}{2}$ to 1 drachm.

Concrete Juices, etc.

Elaterium	$\frac{1}{16}$ to $\frac{1}{2}$ grain.
Opium	$\frac{1}{2}$ to 2 grains.
Aloes	2 to 6 grains.
Camphor	2 to 10 grains.
<i>Black Catechu</i>	5 to 15 grains.
Galls	10 to 20 grains.
Catechu and Kino, <i>Lactucarium</i>	10 to 30 grains.
Manna	1 drm. to 1 oz.

Fixed Oils.

Croton Oil	$\frac{1}{3}$ to 1 minim.
Cod-Liver and Castor Oils	A drm. to an oz.

Volatile Oils **1 to 5 minims.***Except—*

The Oils of Copaiba, Cubebs, and Turpentine	5 to 20 minims.
Oil of Turpentine as an Anthelmintic	1 to 4 drachms.
<i>Oil of Eucalyptus</i> }	5 to 10 minims.
<i>Oil of Winter Green</i> }	5 to 20 minims.
<i>Oil of Chenopodium</i>	5 to 20 minims.
<i>Oil of Eucalyptus</i> as an antiperiodic	$\frac{1}{2}$ to 1 drachm.

Active Principles (officinal).

Digitaline and Atropia (or Atr. Sulph.)	$\frac{1}{60}$ to $\frac{1}{30}$ grain.
Strychnia	$\frac{1}{30}$ to $\frac{1}{12}$ grain.
Veratria	$\frac{1}{12}$ to $\frac{1}{6}$ grain.
Salts of Morphia	$\frac{1}{8}$ to $\frac{1}{2}$ grain.
Podophyllin	$\frac{1}{4}$ to 1 grain.
Santonin	2 to 6 grains.
Sulphates of Quinia and Beberia, as tonics	1 to 5 grains.
” ” ” as antiperiodics	5 to 10 grains.

(non-officinal)—

<i>Apomorphia</i>	$\frac{1}{20}$ to $\frac{1}{5}$ grain.
<i>Pilocarpin</i>	$\frac{1}{8}$ to $\frac{1}{2}$ grain.*
<i>Codeia, Convallamarin</i>	$\frac{1}{2}$ to 2 grains.
<i>Ergotin</i>	2 to 4 grains.†
<i>Caffein</i> and its <i>Citrate</i>	2 to 6 grains.
<i>Salicin</i>	5 to 30 grains.

Cinchona alkaloids, the same as Sulphate of Quinia.

* Usually administered hypodermically in doses of $\frac{1}{15}$ to $\frac{1}{5}$ grain.

† ” ” ” ” ” ” 1 to 2 grains.

β. OF THE OFFICIAL PREPARATIONS.**Extracts.**

Of Calabar Bean	$\frac{1}{16}$ to $\frac{1}{4}$ grain.
„ Stramonium	$\frac{1}{4}$ to $\frac{1}{2}$ grain.
„ Belladonna and Bhang	$\frac{1}{4}$ to 1 grain.
„ Colchicum, Nux Vom., Aconite, and Opium	$\frac{1}{2}$ to 2 grains.
„ Quassia, Poppy, Hemlock, and Aloes	2 to 5 grains.
„ Gentian, Colocynth, Chamomile, Calumba	2 to 10 grains.
„ Henbane	5 to 10 grains.
„ Rhubarb, Lettuce, Jalap, and Hop	5 to 15 grains.
„ Pareira, Krameria	5 to 20 grains.
„ Taraxacum, Logwood, and Liquorice	5 to 30 grains.

Liquid Extracts.

Of Cinchona, Fern, and Ergot	10 to 30 minims.
„ Opium	10 to 40 minims.
„ Liquorice, Bael, Pareira	1 to 2 drachms.
„ Sarsaparilla	2 to 4 drachms.

Juices.

Of Belladonna	5 to 15 minims.
„ Hemlock and Henbane	$\frac{1}{2}$ to 1 drachm.
„ Scoparium and Taraxacum..	1 to 2 drachms.

Essences of Anise and Peppermint	10 to 20 minims.
---	-----	-----	-----	-----	------------------

Infusions, Decoctions, Mixtures, and Waters, 1 to 2 ounces.*With the following exceptions.**Infusions.*

Of Digitalis	1 drm. to $\frac{1}{2}$ ounce.
„ Chamomile, Matico, Bucco, and Cloves	1 to 4 ounces.
„ Cusso	4 to 8 ounces.
„ Linseed	ad libitum.

Decoctions.

Of Aloes	$\frac{1}{2}$ to 1 ounce.
„ Scoparium, Elm, Taraxacum	2 to 4 ounces.
„ Sarsaparilla (both simple and compound)	2 to 10 ounces.

Mixtures.

Of Ammoniac and Gentian	$\frac{1}{2}$ to 1 ounce.
„ Senna	1 to $1\frac{1}{2}$ ounce.

Water.

Of Cherry-laurel	5 to 30 minims.
------------------	----	-----	-----	-----	-----------------

Tinctures $\frac{1}{2}$ to 2 drachms

With the following exceptions: namely, those

Of Aconite	5 to 15 minims.
„ Belladonna, Iodine, Nux Vomica, Cantharides,	}	5 to 20 minims.
Capsicum, Cannabis Indica, Hellebore, and		
Ginger (strong tincture)	5 to 40 minims.
„ Opium	
„ Digitalis, Colchicum, Iron-Salts, Stramony,	}	10 to 30 minims.
Sumbul, Squill, Lobelia, Larch		
„ Hemlock, Chloroform, Ergot, Savin, Camph. Co.	}	15 to 60 minims.
and Ginger...		
„ Tolu	20 to 40 minims.
„ Henbane, Castor, Assafoetida, Myrrh, Benzoin,	}	30 to 60 minims.
and the Ammoniated Tinctures of Opium,		
Guaiac, and Valerian	
„ Senna	2 to 4 drachms.
„ Rhubarb (as a purgative)	4 to 8 drachms.

Wines.

Of Antimony, Ipecac., Opium, and Colchicum	5 to 40 minims.
„ Rhubarb and Aloes	1 to 2 drachms.
„ Iron and its Citrate	1 to 4 drachms.
„ Ipecac. as an emetic	3 to 6 drachms.
„ Quinine; and Antimony, as an emetic	4 to 8 drachms.

Spirits $\frac{1}{2}$ to 1 drachm.

With the following exceptions, viz. :—

Of Camphor, Chloroform, and Rosemary	10 to 30 minims.
„ Nitrous Ether	$\frac{1}{2}$ to 2 drachms.

Syrups 1 drachm.

With the following exceptions, viz. :—

Of Squill and Iodide of Iron	$\frac{1}{2}$ to 1 drachm.
„ Rhubarb and Senna	1 to 4 drachms.

Oxymel of Squill $\frac{1}{2}$ to 1 drachm.
Vinegar of Squill 1 to 2 drachms

Solutions (Liquors).

Hypodermic Injection of Morphia	1 to 6 minims.
Arsenical Solutions	2 to 8 minims.
Liq. Strychniæ and Liq. Ammon. Fortior	5 to 10 minims.
Sols. of Ammonia, Chlorine, and Iron Salts	10 to 30 minims.
Liq. Potassæ, Liq. Sodæ, Liq. Calcis Saccharatus, and Solutions of Morphia Salts	}	15 to 60 minims.
Liq. Bismuthi et Ammoniac Citratis	30 to 60 minims.
Liq. Hydrarg. Perchloridi	30 to 120 minims.
Liq. Potassæ Permanganatis	2 to 4 drachms.
Solutions of Ammonia Salts	2 to 6 drachms.
Liq. Magnesiac Carbonatis	1 to 2 ounces.
Liq. Calcis	1 to 4 ounces.
Liq. Magnesiac Citratis and the Effervescent Solutions	5 to 10 ounces.

Oleum Phosphoratum	5 to 10 minims.
--------------------	-----	-----	-----	-----	-----------------

Powders.

Pulv. Elaterii Co.	1 to 5 grains.
„ Opii Co...	3 to 8 grains.
Hydrargyrum cum Creta	3 to 10 grains.
Pulv. Antimonialis	5 to 15 grains.
„ Ipecacuanhæ Co.	5 to 20 grains.
„ Kino Co.	10 to 20 grains.
„ Scammonii Co.	10 to 30 grains.
„ Cinnamomi Co.	10 to 40 grains.
„ Cretæ Aromaticus c. Opii	10 to 60 grains.
„ Cretæ Aromaticus	1 to 2 scruples.
„ Catechu Co.	1 scr. to 1 drachm
Comp. Powders of Jalap, Rhubarb, Liquorice, and Tragacanth	1 to 2 drachms.
Pulv. Amygdalæ Co.	

Confections 1 to 2 drachms

With the following exceptions, viz. :—

Of Opium	5 to 20 grains.
„ Scammony	10 to 30 grains.

Pills ... 5 to 10 grains.

With the following exceptions—

Pil. Saponis Co. and Pil. Pb. c. Opio	3 to 5 grains.
Pil. Phosphori	3 to 6 grains.
Pil. Hydrargyri and Pil. Ferri Iodidi	3 to 8 grains.
Pil. Scammonii Co.	5 to 15 grains.
Pil. Ferri Carbonatis	5 to 20 grains.

* * *For the doses of standard preparations of non-officinal drugs, see p. 85.*

PROPORTION OF DOSE TO AGE :

The following scale (Gaubius') is of general acceptance.

One-twelfth.	One-eighth.	One-sixth.	One-fourth.	One-third.	One-half.	Two-thirds.	Full dose.
Yrs. 1	2	3	4	7	14	21	21

Beyond 65 yrs. diminish in the inverse gradation of the above.

Children tolerate almost adult doses of *Mercury*, but only the most minute quantities of *Opium*.

In injecting subcutaneously, commence with half the dose which would be given by the mouth.

PROPORTION OF ACTIVE INGREDIENTS IN THE PHARMACOPŒIAL PREPARATIONS.*

Dilute Acids.

See page 7.

Solutions.

Amount of active ingredient in one drachm.

Liq. Calcis ; Liq. Hydrarg. Perchloridi	$\frac{1}{10}$ grain.
Liq. Pb. Subacet. Dil.	$\frac{1}{5}$ grain.
Liq. Potassæ Permanganatis ;†	Arsenical solutions and			
solutions of Alkaloids	$\frac{1}{2}$ grain.
Liq. Calcis Saccharatus	1 grain.
Liq. Magnesiæ Carbonatis	$1\frac{1}{2}$ grain.
Liq. Sodæ	$2\frac{1}{3}$ grains.
Liq. Iodi	$2\frac{1}{2}$ grains.
Liq. Potassæ	$3\frac{1}{2}$ grains.
Liq. Ferri Perchloridi	8 grains.
Liq. Plumbi Subacetatis	16 grains.
Liq. Epispasticus (of <i>Cantharides</i>)	22 grains.
Liq. F. Perchlor. Fortior	32 grains.
Liq. Zinci Chloridi‡	40 grains.

Liq. Bismuthi et Amm. Citratis contains 3 grains of Bismuthic Oxide in the drachm.

Liq. Calcis Chloratæ is made with $5\frac{1}{4}$ grains to the drachm.

Liq. Chlori contains $\frac{1}{8}$ grain of Chlorine in the drachm.

Liq. Ammonię „ $5\frac{1}{8}$ „ „ Ammonia „ „

Liq. Amm. Fortior contains nearly 16 grains of Ammonia in the drachm.

Inj. Morphię Hypodermica contains 5 grains of the Acetate in the drachm.

Oleum Phosphoratum contains $\frac{1}{10}$ grain of Phosphorus in 10 minims.

Syrupus contains 6 oz. of sugar in 7 fluid oz.

* For the proportion of Opium in its preparations, see also p. 59.

† “Condyl’s Fluid” contains 1 grain of the Permanganate in the drachm.

‡ “Burnett’s Solution” is a concentrated solution of Zincic Chloride, of sp. gr. 2.0.

Syrups.

Amount of the active drug contained in, or corresponding to, one drachm.

Syr. Ferri Phosphatis	1 grain.
Syr. Tolutanus	2 grains.
Syr. Scillæ	3 grains.
Syr. Rhei	4 grains.
Syr. Ferri Iodidi	4½ grains.
Syr. Chloral	10 grains.
Syr. Papaveris	25 grains.
Syr. Sennæ	27 grains.

Wines.

Ditto.

Vin. Quiniæ	⅛ grain.
Vin. Antimoniale (<i>of Tartar Emetic</i>)	¼ grain.
Vin. Ferri Citratis (<i>of F. et Amm. Citr.</i>)	1 grain.
Vin. Aloes	2 grains.
Vin. Ipecacuanhæ	2¾ grains.
Vin. Rhei	4 grains.
Vin. Opii (<i>of Opium</i>)	5½ grains.
Vin. Colchici	11 grains.

Tinctures.

Ditto.

Tr. Quiniæ, Q. Ammoniata	1 grain.
Tr. Iodi	1⅓ grain.
Tr. Cannabis Indicæ (<i>of the Extract</i>)	2¾ grains.
Tr. Ferri Acetatis	6 grains.
Tr. Ferri Perchloridi	8 grains.
Tr. Chloroformi Co. (<i>of Chloroform</i>)	6 minims.
Tr. Lavandulæ Co. (<i>of Ol. Lavandulæ</i>)	¼ minim.
Tr. Opii Ammoniata (<i>of Opium</i>)	½ grain.
Tr. Cantharidis ; Cardam. Co.	⅔ grain.
Tr. Aloes	1⅓ grains.
Tr. Capsici ; Quassiæ	2 grains.
Tr. Arnicæ ; Belladonnæ ; Castorei ; Croci	2¾ grains.
Tr. Opii ; Gentianæ Co.	4 grains.
Tr. Aurantii ; Benzoini Co. ; Cinchonæ Co. ; Kino ; Nucis Vomicæ ; Rhei	5½ grains.
Tr. Cinchonæ Flavæ ; Guaiaci ; Pyrethri ; Veratri Viridis	11 grains.
Tr. Ergotæ	14 grains.
Tr. Zingiberis Fortior	27 grains.
Tr. Camphoræ Co. { <i>of Opium and of Benzoic Acid</i>	⅓ grain.
Tr. Camphoræ Co. { <i>of Camphor</i>	⅓ grain.
In all other cases	7 grains.

Vinegars.*Ditto.*

Acet. Cantharidis	5 $\frac{1}{4}$ grains.
Acet. Scillæ	7 grains.

Spirits.*See page 6.***Glycerines.***See page 7.***Essences.***See page 5.***Waters.***Amount of active ingredient in 1 fluid ounce.*

Aq. Camphoræ	$\frac{1}{8}$ grain.
Aq. Chloroformi	2 $\frac{1}{2}$ minims.

*See also page 6.***Infusions and Decoctions.***Amount of the active drug to which 1 fluid ounce corresponds.*

Inf. Digitalis	3 grains.
Inf. Gentianæ ; Quassia	6 grains.
Inf. Aurantii Co. ; Chirata ; Caryophylli ; Ergotæ ; Rhei ; Rosæ Acidum* ; Serpentariæ	11 grains.
Inf. Valerianæ	12 grains.
Inf. Catechu ; Lini	16 grains.
Inf. Cusso	27 grains.
Inf. Cascarillæ ; Dulcamaræ ; Sennæ	44 grains.
Dec. Aloes Co.	4 grains.
Dec. Cinchonæ Flavæ ; Hæmatoxyli ; Quercus	27 grains.
Dec. Hordei ; Pareiræ	30 grains.
Dec. Granati Radicis ; Papaveris	44 grains.
Dec. Ulmi ; Sarsæ ; Sarsæ Co.	55 grains.

* And 6 minims of Acid, Sulphuric. Dil.

In all other cases	22 grains.
--------------------	-----	-----	-----	-----	-----	-----	------------

Mixtures.

Amount of the active drug contained in, or corresponding to, 1 fluid ounce.

Mistura Creasoti	1 minim.
Mist. Scammonii	2 grains.
Mist. Ferri Co. (of <i>Ferri Sulphas</i>)	2½ grains.
Mist. Guaiaci	11 grains.
Mist. Cretæ	13 grains.
Mist. Ammoniaci	14 grains.
Mist. Sennæ Co. (of <i>Magn. Sulphas</i>)	88 grains.
Mist. Sp. Vini Gallici (of <i>Brandy</i>)	3 drachms.

Powders.

See pages 152, 153.

Confections.

See page 153.

Pill-masses.

Amount of active ingredient in 3 grains.

Pil. Phosphori	$\frac{1}{30}$ grain.
Pil. Hydrargyri; Ferri Iodidi	1 grain.
Pil. Quiniæ	2½ grains.

Amount of active ingredient in 5 grains.

Pil. Aloes Barbadoes; A. Socotrinæ	2 grains.
Pil. Aloes et Assafoetidæ (of both)	1½ grain.
Pil. Aloes et Ferri	{ of Aloes	1 grain.
	{ of <i>F. Sulphas</i>	$\frac{3}{4}$ grain.
Pil. Aloes et Myrrhæ	{ of Aloes	1½ grain.
	{ of <i>Myrrh</i>	half as much.
Pil. Assafoetidæ Co. (of <i>Assafœtida</i> and of <i>Galbanum</i>)	1½ grain.
Pil. Cambogiæ Co. (of <i>Gamboge</i> and of <i>Aloes</i>)	1 grain.
Pil. Colocynthis Co.	{ of <i>Colocynth</i>	1 grain.
	{ of <i>Aloes</i>	2 grains.
Pil. Colocynth. et Hyoscyami	half the above.
Pil. Conii Co. (of <i>Extract of Hemlock</i>)	3½ grains.
Pil. Ferri Carbonatis (of <i>F. Carb. Sacch.</i>)	4 grains.

Pil. Hydrargyri Subchloridi Co. (of Calomel and of Sulphd. Antimony)	1 grain.
Pil. Ipecac. cum Scilla (of Opium)	$\frac{1}{6}$ grain.
Pil. Plumbi cum Opio { of Pb. Acetas of Opium	3 $\frac{3}{4}$ grains. $\frac{5}{8}$ grain.
Pil. Rhei Co. { of Rhubarb of Aloes	1 $\frac{1}{4}$ grain. 1 grain.
Pil. Saponis Co. (of Opium)	1 grain.
Pil. Scammonii Co. (of Scammony and of Jalap Resins)... ..	1 $\frac{1}{2}$ grains.
Pil. Scillæ Co. (of Squill)	1 grain.

Lozenges.*Amount of active ingredient in each Lozenge.*

Troch. Morphiæ	$\frac{1}{30}$ grain.
Troch. Morph. c. Ipecac. { of Morphia of Ipecacuanha	$\frac{1}{2}$ grain. $\frac{1}{10}$ grain.
Troch. Opii (of Extract of Opium)	$\frac{1}{4}$ grain.
Troch. Ipecacuanhæ	$\frac{1}{2}$ grain.
Troch. Acidi Tannici	1 grain.
Troch. Catechu ; Ferri Redacti	2 grains.
Troch. Bismuthi (of B. Subnitras)... ..	5 grains.
Troch. Pot. Chloratis ; Sodæ Bicarbonatis	

Suppositories.*Amount of active ingredient in each Suppository.*

Suppositories of Morphia	$\frac{1}{2}$ grain.
Suppositories of Carbolic Acid and Opium	1 grain.
Supp. Plumbi Co. { of Opium of Pb. Acetas	1 grain. 3 grains.
Suppositories of Tannic Acid	3 grains.
Mercurial Suppository	5 grains.

Enemas.*See page 9.***Inhalations.***See page 8.***Lotions.***See page 151.*

Poultices.*See page 9.***Liniments.***See page 159.***Ointments.***Amount of active ingredient in 1 drachm.*

Ung. Aconitiæ, Atropiæ, Veratriæ	1 grain.
Ung. Plumbi Acetatis	1 $\frac{3}{4}$ grain.
Ung. Iodi	2 grains.
Ung. Plumbi Subacetatis	3 grains.
Ung. Pot. Sulphuratæ ; Sulphuris Iodidi	4 grains.
Ung. Creasoti ; Pot. Iodidi	6 $\frac{2}{3}$ grains.
Ung. Cadmii ; Cantharidis ; Plumbi Carbonatis ; Pb. Iodidi	7 $\frac{1}{2}$ grains.
Ung. Belladonnæ ; Gallæ ; Zinci (<i>of Z. Oxidum</i>)	9 grains.
Ung. Antim. Tartarati ; Cetacei ; Elemi ; Sulphuris	12 grains.
Ung. Resinæ ; Sabinæ	17 grains.
Ung. Hydrargyri Iodidi Rubri	2 grains.
Ung. H. Nitratis	4 grains.
Ung. H. Oxidi Rubri ; H. Ammoniati	7 $\frac{1}{2}$ grains.
Ung. H. Subchloridi	9 grains.
Ung. Hydrargyri Co. (<i>of Mercury</i>)	13 grains.
Ung. Hydrargyri	1 part in 2.
Ung. Terebinthinæ	1 part in 2.
Ung. Picis Liquidæ	5 parts in 8.
Ung. Gallæ cum Opio { <i>of Galls</i>	9 grains (nearly).
{ <i>of Opium</i>	4 grains.

Extracts.*Proportion yielded by—*

Quassia	2 per cent.
Belladonna, Colchicum	4 " "
Physostigma, Lettuce, Hemlock, Henbane, Hop, Colchi- } cum (<i>Acetic Extract</i>)	5 " "
Nux Vomica	6 " "
Aconite	(variable)	7 " "
Indian Hemp, Dandelion	8 " "
Stramony	12 " "
Calumba	15 " "
Pareira	20 " "

Rhatany, Mezereon	30 per cent.
Chamomile, Rhubarb	40 „ „
Socotrine Aloes, Opium, Gentian, Jalap	50 „ „
Barbadoes Aloes	75 „ „

Colocynth (*of Compound Extract*)—four times the Colocynth.

Liquid Extracts—

Fern	9 per cent.
Cinchona	one-fourth.
Sarsaparilla	one-half.
Ergot, Bael, Pareira	equal quantities.
Opium	ten times.

These numbers are all subject to variations.

Juices

Ditto. (The numbers are approximative.)

Taraxacum	30 p. c.		Hemlock and Henbane ...	50 p. c.
Scoparium	40 „		Belladonna ...	60 „

EFFERVESCENT DRAUGHTS.

Used to neutralise 20 grains of—

			<i>Ac. Citricum.</i>	<i>Ac. Tartaricum.</i>
Potass. Bicarbonas	14 grains.	15 grains.
Sodæ Bicarbonas	17 grains.	19 grains.

To neutralize 15 grains of—

Amm. Carbonas	17 grains.	20 grains.
Magn. Carbonas	19 grains.	23 grains.

To neutralize 10 grains of—

Lithiæ Carbonas	20 grains.	24 grains.
---------------------	-----	-----	------------	------------

Half an ounce of fresh Lemon-juice contains 16 grains of Acidum Citricum.

INCOMPATIBILITY.

Most cases of chemical incompatibility between drugs treated of in the foregoing pages will be obvious to anyone possessing a rudimentary knowledge of chemistry. The following points it may be as well to direct especial attention to :—

- 1.—Permanganate of Potash is decomposed in presence of organic substances. Nitrate of Silver and Corrosive Sublimate are liable to decomposition in similar circumstances. Nitrate of Silver forms an explosive compound with Creasote.
- 2.—Acidum Nitrohydrochloricum Dil. contains free Chlorine, which is liable to decompose organic colouring and other matters.
- 3.—Spiritus Ætheris Nitrosi decomposes the Iodides and Bromides, blackens solutions of Ferri Sulphas, and decomposes with Guaiac Resin.
- 4.—Alkaloids are for the most part precipitated from solutions of their salts by caustic alkalies and their carbonates.* The various solutions, infusions, etc., contain the alkaloids in the form of salts. The volatile alkaloids, such as Conia, are liberated by caustic alkalies.
- 5.—Many alkaloids are precipitated by the alkaline iodides.*
- 6.—Alkaloids are for the most part precipitated by Tannic and Gallic Acids,* or by preparations of drugs containing them.
- 7.—Tannic Acid precipitates with alkalies and their carbonates, with Ferric salts, most metallic salts, Arsenious Acid, Arsenites,* and Tartar Emetic. Gallic Acid blackens solutions of Ferric salts.
- 8.—Preparations of Opium precipitate metallic (not alkaline) salts, and Arsenious Acid.*
- 9.—Solutions of Potassæ Tartras precipitate the Acid Tartrate when acidulated.
- 10.—Gum Acacia forms insoluble compounds with Subacetate of Lead and with Borax. It decomposes Ferric Salts.
- 11.—Resins are precipitated from spirituous solutions on dilution with water (see p. 48).

The presence of free acid in certain preparations should be especially noted.

* It must not, however, be concluded that these precipitates are therapeutically inactive.

INDEX.

The "Popular Designations" given alphabetically on pp. 162, 163, are not repeated in the Index.

- Acaciæ Gummi, 52, 74
Acetic Acid, 100, 106
Acetic Ether, 100
Acetum, 100
Acids, Dilute, 7, 158
Acids, Mineral, 96
Acids, Organic, 61, 88, 105
Acidum Aceticum, 106
Acidum Arseniosum, 141
Acidum Benzoicum, 61, 78
Acidum Carbolicum, 107
Acidum Citricum, 109
Acidum Gallicum, 109
Acidum Hydrochloricum, 96
Acidum Hydrocyanicum Dil., 110
Acidum Lacticum, 109
Acidum Nitricum, 96
Acidum Nitrohydrochloricum Dil., 97
Acidum Phosphoricum Dil., 97
Acidum Salicylicum, 108
Acidum Sulphuricum, 96
Acidum Sulphurosum, 97
Acidum Tannicum, 109
Acidum Tartaricum, 108
Aconiti Folia, 25, 72
Aconiti Radix, 28, 72
Aconitia, 62, 65, 72
Actæa Racemosa, 84
Active Principles of Definite Composition, 60, 88
Adeps, 12
Æther, 100
Æther Aceticus, 100
Albumen Ovi, 12
Alcohol, 99
Alcohol Amylicum, 101
Alkaloids, 62, 89
 " Extraction of, 65
Almond Oil, 42, 75
Almonds, 39, 75
Aloes, 55, 82
Alumen, 143
Ammonia, 123
Ammoniacum, 53, 76
Ammoniated Mercury, 130
Ammoniated Tinctures, 6
Ammonium Compounds, 123
Amygdala, 39, 75
Amyl Nitris, 101
Amylic Alcohol, 101
Amylum, 60, 83
Amyridaceæ, 74
Anacardiaceæ, 74
Anethi Fructus, 34, 76
Anethi Oleum, 45, 76
Anethol, 44
Angustura Bark, 23, 74
Animal Kingdom, drugs furnished by, 12
Anisi Oleum, 47, 72, 76
Anthemidis Flores, 20, 77
Anthemidis Oleum, 46, 77
Antimonial Compounds, 139
Apomorphia, 90
Aqua, 92
Aquæ, 6
Arabin, 52
Areca, 39, 82
Argentum, 145
Argol, 116
Aristolochiaceæ, 80
Armoracia, 28, 73

- Arnica, 28, 77
 Arsenical Compounds, 141
 Arsenious Acid, 141
 Artanthic Acid, 25
 Asclepiadaceæ, 78
 Assafœtida, 53, 76
 Atropaceæ, 79
 Atropia, 62, 65, 79
 Atropiæ Sulphas, 62, 66, 79
 Attar of Roses, 20
 Aurantiaceæ, 73
 Aurantii Floris Aqua, 20, 73
 Aurantii Fructus, 34, 73

 Bael Fruit, 36, 73
 Balsams, 51
 Balsamum Dipterocarpi, 86
 Balsamum Peruvianum, 51, 74
 Balsamum Tolutanum, 51, 74
 Barks, 13
 Barley, Pearl, 39, 83
 Bearberry, 26, 78
 Bebeeru, 24, 80
 Beberia Sulphas, 62, 66, 80
 Bela, 36, 73
 Belladonnæ Folia, 25, 79
 Belladonnæ Radix, 20, 79
 Benzoic Acid, 51, 61, 78
 Benzoinum, 51, 78
 Berberia, 28, 84
 Betel-Nut, 39, 82
 Bhang, 19, 81
 Bismuth Compounds, 138
 Black Oxide of Manganese, 146
 Blisters, 15
 Bone-ash, 93, 132
 Bone-black, 92
 Borax, 120
 Bornéene, 30
 Brandy, 104
 Bread-crumbs, 112
 Bromum (*Bromine*), 94
 Broom, 18, 75
 Brucia, 40, 84
 Buchu, 25, 74
 Buckthorn, 37, 74
 Buds, 21
 Burgundy Pitch, 48, 81
 Burnett's Solution, 174
 Byttneriaceæ, 73

 Cadmii Iodidum, 146

 Caffein, 89
 Cajeput Oil, 46, 75
 Calabar Bean, 40, 75
 Calamine, 134
 Calcium Compounds, 132
 Calumba, 28, 72
 Calx, 132
 Calx Chlorata, 133
 Cambogia, 53, 73
 Camphora, 46, 80
 Canada Balsam, 50, 131
 Canella Alba, 23, 73
 Canellaceæ, 73
 Cannabinaceæ, 81
 Cannabis Indica, 19, 81
 Cantharis, 14
 Caprifoliaceæ, 77
 Capsicum, 36, 79
 Capsules, 11
 Caraway, 35, 76
 Carbo Animalis, 92
 Carbo Ligni, 92
 Carbohydrates, 60
 Carbolic Acid, 107
 Cardamomum, 39, 82
 Carmine, 14
 Carui Fructus, 35, 76
 Carui Oleum, 45, 76
 Carvol, 44
 Caryophylli Oleum, 45, 76
 Caryophyllum, 21, 76
 Cascarella, 23, 80
 Cassia, 36, 74
 Castoreum, 14
 Castor Oil, 42, 80
 Cataplasmata, 9
 Catechu Nigrum, 86
 Catechu Pallidum, 56, 77
 Cathartin, 26
 Cephælic Acid, 29
 Cera, 12
 Cerevisiæ Fermentum, 17, 83
 Cerii Oxalas, 146
 Cetaceum, 14
 Cetraria, 17, 83
 Cevadilla, 37, 82
 Chalk, 132
 Chamomile, 20, 77
 Charcoal, animal, 92
 „ vegetable, 92
 Charta Epispastica, 15
 Chartæ, 10

- Cherry Laurel, 25, 75
 Chimaphila, 84
 Chirata, 18, 78
 Chloral Hydras, 101
 Chloroforms, 10
 Chloroformum, 101
 Chlorum (*Chlorine*), 94
 Chrysophanic Acid, 29, 88
 Cimicifugin, 84
 Cinchona Flava, Pallida, Rubra, 23,
 77; C. Lancifolia, 62, 77
 Cinchona Alkaloids, 89
 Cinnamic Acid, 51
 Cinnamomi Cortex, 23, 80
 Cinnamomi Oleum, 45, 80
 Citric Acid, 109
 Cloves, 21, 76
 Clysters, 9
 Coccus (*Cochineal*), 14
 Cocoa-butter, 43, 73
 Cod-Liver Oil, 14
 Codeia, 58, 89
 Colchici Cormus, 28, 82
 „ Semina, 39, 82
 Collodium, 111
 Collyria, 10
 Colocynthis, 36, 76
 Compositæ, 77
 Compound Preparations, Synopsis of,
 152
 Concrete Juices, 55, 86
 Confections, 8
 Coniferae, 81
 Conii Folia, 25, 76
 „ Fructus, 36, 76
 Convallaria, 84
 Convolvulaceæ, 78
 Copaiba, 50, 75
 Copaibæ Oleum, 46, 75
 Copper, 146
 Coriandri Fructus, 34, 76
 „ Oleum, 45, 76
 Cotton-wool, 40, 73
 Creasotum, 106
 Creta, 132
 Crocus, 20, 82
 Croton-Chloral Hydrate, 101
 Crotonis Oleum, 42, 80
 Cruciferae, 73
 Cryptopia, 58
 Cubeba, 36, 80
 Cubebæ Oleum, 46, 80
 Cucurbitaceæ, 76
 Cupri Sulphas, 146
 Cupuliferae, 81
 Curd Soap, 120
 Cusparia, 23, 74
 Cusso, 19, 75
 Cymene, 44
 Dandelion, 38, 77
 Datura Tatula, 84
 Daturia, 26, 40
 Deadly Nightshade, 79
 Decoctions, 5, 156
 Digitalinum, 61, 70, 79
 Digitalis Folia, 25, 79
 Dill, 34, 76
 Discs, 11
 „ Ophthalmic, 11
 Dog-rose, 35, 75
 Dolomite, 142
 Dose, Proportion of, to Age, 173
 Doses, Synopsis of, 165
 „ of Non-official Preparations,
 85
 Duboisia, 84
 Dulcamara, 18, 78
 Ecbalii Fructus, 37, 76
 Egg, 12
 Elaterium, 57, 76
 Elder Flower, 20, 77
 Electuaries, 8
 Elemi, 50, 74
 Elm-bark, 24, 81
 Embrocations, 9, 159
 Emetia, 29
 Emplastra, 10, 161
 Emplastrum Calefaciens, 15
 Emulsin, 39
 Enemas, 9
 Ergota, 17, 83
 Ergotine, 88
 Ericaceæ, 78
 Eserin, 40, 89
 Essences, 5
 Ether, 100
 Eucalyptol, 87
 Eucalyptus, 84
 Euphorbiaceæ, 80
 Extracts, 3, 156
 Eye-washes, 10

- Farina Tritici, 39, 83
 Farina Lini, 40, 73
 Fel Bovinum, 14
 Fennel Fruit, 35, 76
 Fern, 28, 83
 Ferrum, 125
 Ferrum Redactum, 128
 Ficus (*Fig*), 17, 80
 Filices, 83
 Filix Mas, 28, 83
 Fixed Oils, 42, 87
 Flax, 40, 73
 Flexile Collodion, 111
 Flour, 39
 Flowering Tops, 19
 Flowers, 20
 Fœniculi Fructus, 35, 76
 Frankincense, 50, 81
 Fruits, 34
 Fungi and Lichens, 17, 83
 Fusel Oil, 101

 Galbanum, 53, 76
 Galla (*Galls*), 56, 81
 Gallic Acid, 109
 Gamboge, 53, 73
 Gargles, 10
 Gelsemium, 84
 Gentiana, 28, 78
 Gentianaceæ, 78
 Ginger, 31, 82
 Glacial Acetic Acid, 106
 Glonoin, 112
 Glycerines, 7
 Glycerinum, 111
 Glycyrrhiza, 28, 75
 Gossypium, 40, 73
 Graminaceæ, 83
 Granati Radix, 23, 76
 Granulated Sulphate of Iron, 125
 Granulated Zinc, 134
 Guaiaci Lignum, 22, 74
 " Resina, 48, 74
 Guarana, 84
 Gum Arabic, 52, 74
 Gum Tragacanth, 52, 75
 Gummi Rubrum, 86
 Gum-resins, 53
 Gums, 52
 Gun-Cotton, 111
 Gunjah, 19, 81
 Gurjun Oil, 86

 Gutta Percha, 57, 78
 Guttiferæ, 73

 Hæmatoxyli Lignum, 22, 75
 Hamamelis, 84
 Hashish, 19, 81
 Hellebore, 30, 82
 Hemidesmus, 29, 78
 Hemlock Leaves, 25, 76
 " Fruit, 36, 76
 Hemp, 19, 81
 Henbane, 25, 79
 Hips, 35, 75
 Hirudo, 13
 Honey, 7, 12
 Hop, 19, 81
 Hordeum Decorticatum, 39, 83
 Horse-radish, 28, 73
 Humulin, 19
 Hydrargyrum, 129
 Hydrargyrum cum Creta, 147
 Hydrastis, 84
 Hydrochloric Acid, 96
 Hydrocyanic Acid, 110
 Hyoseyami Folia, 25, 79
 Hypodermic Injection, 9

 Iceland Moss, 17, 83
 Igasuric Acid, 40
 Ignatia, 84
 Illicium Anisatum, 72
 Incompatibility, 181
 Indian Hemp, 19, 81
 Indian Tobacco, 18, 77
 Infusions, 5, 156
 Inhalations, 8
 Injections (Hypodermic), 9
 Iodoform, 101
 Iodum (*Iodine*), 94
 Ipecacuanha, 29, 77
 Iridaceæ, 82
 Iron Alum, 127
 Iron Compounds, 125

 Jalapa, 29, 78
 Jalapæ Resina, 48, 78
 Jalapine, 88
 James' Powder, 163
 Juices, 5
 Juniperi Oleum, 46, 81

 Kamala, 37, 80

- Kelp, 94
 Kino, 56, 75
 Koussou, 19, 75
 Krameria, 29, 73

 Labiatae, 79
 Lac, 12
 Lactic Acid, 109
 Lactuca, 18, 77
 Lactucarium, 86
 Lactucin, Lactucone, 86
 Lamels, 11
 Lard, 12
 Larix (*Larch*), 23, 81
 Latin Phrases, 164
 Lauraceae, 80
 Lauro-cerasi Folia, 25, 75
 Lavandulæ (*Lavander*) Oleum, 45, 79
 Lead Compounds, 136
 Leaves, 25
 Leeches, 13
 Leguminosae, 74
 Lemon, 34, 73
 Lettuce, 18, 77
 Lichens, 17, 83
 Liliaceae, 82
 Lime, 132
 Limonis Cortex, Succus, 34, 73
 „ Oleum, 45, 73
 Linaceae, 73
 Linctuses, 10
 Lini Farina, 40, 73
 „ Oleum, 42, 73
 „ Semina, 40, 73
 Liniments, 9, 159
 Linseed, 40, 73
 Meal, 40, 73
 „ Oil, 42, 73
 Liquidambaraceae, 81
 Liquor Epispasticus, 15
 Liquores, 6, 158
 Liquorice, 28, 75
 Litharge, 136
 Lithium Compounds, 144
 Lobelia, 18, 77
 Lobeliaceae, 77
 Loganiaceae, 78
 Logwood, 22, 75
 Lotions, 9, 130
 Lozenges, 8, 158
 Lupulus, 19, 81

 Magnesium Compounds, 142
 Magnetic Oxide of Iron, 128
 Magnoliaceae, 72
 Male Fern, 28, 83
 Malvaceae, 73
 Manganesi Oxidum Nigrum, 146
 Manna, 57, 78
 Margarin, 42
 Marmor Album (*Marble*), 133
 Mastiche, 48, 74
 Maticæ Folia, 25, 80
 Meconic Acid, 58
 Meconin, 58
 Mel, 7, 12
 Melanthaceae, 82
 Menispermaceae, 72
 Menthae Piperitæ Oleum, 45, 79
 „ Viridis Oleum, 45, 79
 Menthol, 44, 88
 Mercurial Compounds, 129
 Metarabin, 52
 Methyl-Conia, 25, 36
 Mezerei Cortex, 23, 79
 Mica Panis, 112
 Milk, 12
 Milk Sugar, 15
 Mineral Acids, 96
 Minerals and Manufactured Products, 91
 Mixtures, 8, 155
 Momordicin, 57
 Moraceae, 80
 Mori Succus, 34, 80
 Morphia, 62, 67, 72
 Moschus, 14
 Mucilages, 7
 Mulberry, 34, 80
 Musk, 14
 Mustard, 40, 73
 Myristica, 40, 80
 Myristicaceae, 80
 Myristicæ Oleum, 45, 80
 „ „ Expressum, 42, 80
 Myristicol, 44
 Myronic Acid, 40
 Myrosin, 40
 Myrrha, 53, 74
 Myrtaceae, 76

 Narceia, 58
 Narcotin, 58
 Nectandræ Cortex, 24, 80

- Nicotia, 26
 Nitre, 114
 Nitric Acid, 96
 Nitrite of Amyl, 101
 Nitroglycerin, 112
 Nitro-hydrochloric Acid, 97
 Nitrous Ether, 100
 Nitrous Oxide, 95
 Nutmeg, 40, 80
 Nutmeg-butter, 42
 Nux Vomica, 40, 78

 Oak Bark, 24, 81
 Oil of Bitter Almonds, 39
 Oil of Cade, 105
 Oil of Male Fern, 31
 Oil of Winter Green, 87
 Oils, 7
 Oils, Fixed, 42, 87
 " Volatile, 44, 87
 Ointments, 9, 159
 Oleaceæ, 78
 Olein, 42
 Oleo-resins, 50, 86
 Oleum Amygdalæ, 42, 75
 " Anethi, 45, 76
 " Anisi, 45, 72, 76
 " Anthemidis, 46, 77
 " Cadinum, 105
 " Cajuputi, 46, 76
 " Carui, 45, 76
 " Caryophylli, 45, 76
 " Chenopodii, 87
 " Cinnamomi, 45, 80
 " Copaibæ, 46, 75
 " Coriandri, 45, 76
 " Crotonis, 42, 80
 " Cubebæ, 46, 80
 " Eucalypti, 87
 " Gualtheriæ, 87
 " Gynocardia, 87
 " Juniperi, 46, 81
 " Lavandulæ, 45, 79
 " Limonis, 45, 73
 " Lini, 42, 73
 " Menthæ Piperitæ, 45, 79
 " Menthæ Viridis, 45, 79
 " Morrhuæ, 14
 " Myristicæ, 45, 80
 " Expressum, 42, 80
 " Olivæ, 32, 78
 " Phosphoratum, 95

 Oleum Pimentæ, 45, 76
 " Ricini, 42, 80
 " Rosmarini, 45, 79
 " Rutæ, 46, 74
 " Sabinæ, 47, 81
 " Santali Flavi, 87
 " Sinapis, 47, 73
 " Staphisagriæ, 87
 " Terebinthinæ, 47, 81
 " Theobromæ, 43, 73
 Olivæ Oleum, 32, 78
 Opium, 58, 72
 Orange, 34, 73
 Orange Flower Water, 20, 73
 Orange Wine, 104
 Os Ustum, 93, 132
 Ovi Albumen, 12
 " Vitellus, 12
 Ox Bile, 14
 Oxymer, 7, 13

 Palmaceæ, 82
 Papaver Rhæas, 20, 72
 Papaveraceæ, 72
 Papaverin, 58
 Papaveris Capsulæ, 37, 72
 Papers, 10
 Pareira, 26, 72
 Pearl Barley, 39, 83
 Pellitory, 29, 77
 Pelosia, 29
 Pepper, 37, 80
 Peppermint, Oil of, 45, 79
 Pepsin, 14
 Peruvian Balsam, 51, 74
 Pessaries, 10
 Pharmacopœia, the, 1
 Phenol, 107
 Phosphoric Acid, 97
 Phosphorus, 93
 Phrases used in Prescribing, 164
 Physostigma, 40, 75
 Pill-Masses, 8, 153
 Pilocarpin, 90
 Pimaric Acid, 49
 Pimenta, 35, 76
 Pimentæ Oleum, 45, 76
 Pinic Acid, 49
 Piper Nigrum, 37, 80
 Piperaceæ, 80
 Pix Burgundica, 48, 81
 Pix Liquida (*Pitch*), 105

- Plants Entire, 18
 Plasters, 10, 161
 Plumbum, 136
 Podophylli Radix, 29, 72
 „ Resina, 49, 72
 Polycroite, 20
 Polygalaceæ, 73
 Polygalic Acid, 30
 Polygonaceæ, 79
 Pomegranate, 23, 76
 Poppy Capsules, 37, 72
 Popular Designations, 162
 Posological Tables, 165
 Potassium Compounds, 114
 Poultrices, 9
 Powders, Compound, 8, 152
 Preparations, the Pharmacopœial, 2
 „ Synopsis of Compound, 154
 Prescribing, Latin Phrases used in, 152
 Proof Spirit, 99
 Proportion of active ingredients in the Pharmacopœial Preparations, 174
 Proportion of Dose to Age, 173
 Prunum, 37, 75
 Prussic Acid, 25, 110
 Pterocarpi Lignum, 22, 75
 Pulveres Compositæ, 8
 Punica Granatum, 23, 76
 Pyrethrum, 29, 77
 Pyroxylin, 111

 Quassia, 46, 74
 Quercus, 24, 81
 Quinæ Sulphas, 62, 68, 77
 Quinidin, 23, 89

 Raisins, 35, 73
 Ranunculaceæ, 72
 Rectified Spirit, 99
 Red Gum, 86
 Red Poppy, 20, 72
 Resina, 49, 81
 Resins, 48
 Rhamnaceæ, 74
 Rhamni Succus, 37, 74
 Rhamnus Frangula, 85
 Rhatany, 29, 73
 Rheum, 29, 79
 Rhœados Petala, 20, 72
 Rhubarb, 29, 79
 Ricini Oleum, 42, 80

 Roots and Rhizomes, 28
 Rosa Canina (*Hips*), 35, 75
 Rosa Centifolia (*Cabbage Rose*), 20, 75
 Rosa Gallica (*Red Rose*), 20, 75
 Rosaceæ, 75
 Rosin, 49, 81
 Rosmarini (*Rosemary*) Oleum, 45, 79
 Rubiaceæ, 77
 Rutaceæ, 74
 Rutæ (*Rue*) Oleum, 46, 74

 Sabadilla, 37, 82
 Sabinæ Cacumina, 13, 81
 Sabinæ Oleum, 47, 81
 Saccharum Lactis, 15
 Saccharum Purificatum, 60, 83
 Saffron, 20, 82
 St. Ignatius' Bean, 84
 Salicin, 88
 Salicylic Acid, 108
 Saltpetre, 114
 Sambuci Flores, 20, 77
 Sandal-wood, Red, 22, 75
 Santali Flavi Oleum, 87
 Santalin, 22
 Santonica, 21, 77
 Santoninum, 60, 70, 77
 Sapo Animalis, 120
 Sapo Durus, 120
 Sapo Mollis, 116
 Sapotaceæ, 78
 Sarsæ Radix (*Sarsaparilla*), 29, 82
 Sassafras, 29, 80
 Savin Tops, 18, 81
 Scammoniae Radix, 30, 78
 „ Resina, 44, 78
 Scammonium, 53, 78
 Scilla, 30, 82
 Scoparii Cacumina, 18, 75
 Scrophulariaceæ, 79
 Seeds, 39
 Senega, 30, 73
 Senna, 26, 75
 Serpentaria, 30, 80
 Sevum, 12
 Sherry, 104
 Silver Compounds, 145
 Simarubaceæ, 74
 Sinalbin, 40
 Sinapis, 40, 73
 Sinapis Oleum, 47, 73
 Smilaceæ, 82

- Smilacin, 29
 Soap, Animal or Curd, 120
 " Castile or Hard, 120
 " Soft, 116
 Sodinm Compounds, 118
 Solanaceæ, 79
 Solania, 18
 Solutions, 6, 158
 Spanish Fly, 14
 Sparteia, 18
 Spearmint, Oil of, 45, 79
 Spermaceti, 14
 Spigelia, 85
 Spirits, 6, 158
 Spiritus Ætheris Nitrosi, 100
 Spiritus Rectificatus, Tenuior, 99
 Spiritus Vini Gallici, 104
 Squill, 30, 82
 Squinting Cucumber, 37, 76
 Staphisagriæ Oleum, 87
 Star Anise, 72
 Starch, 60, 83
 Stavesacre, Oil of, 87
 Storax, 51, 81
 Stramonii Folia, 26, 79
 " Semina, 40, 79
 Strobile, 19
 Strychia, 62, 68, 78, 84
 Styraaceæ, 78
 Styracin, 51
 Styra, 51, 81
 Subcutaneous Injection, 9
 Succî, 5
 Snet, 12
 Sugar, Cane, 60, 83
 " Milk, 15
 Sulphur, 93
 Sulphuric Acid, 96
 Sulphuris Iodidum, 93
 Sulphurous Acid, 97
 Sumbul, 30, 76
 Suppositories, 9, 159
 " Urethral, 10
 Sylvic Acid, 49
 Symbols, 11
 Synopsis, Botanical, 72
 " of Compound Preparations, 174
 " of Doses, 165
 Syrups, 7, 157
 Syrupus, 7, 61
 Tabaci Folia, 26, 79
 Table of Doses, 85, 165
 " Latin Phrases used in Pre-
 scribing, 146
 " Popular Designations, 162
 " Proportion of Active Ingre-
 dients in the Pharmacopœial Pre-
 parations, 174
 Tablets, 11
 Tamarindus, 18, 75
 Tannic Acid, 109
 Tar, Wood, 105
 " Coal, 107
 Taraxacum, 38, 77
 Tartar Emetic, 139
 Tartaric Acid, 108
 Terebinthina Canadensis, 50, 81
 Terebinthinæ Oleum, 47, 81
 Thebaia, 58
 Thebolactic Acid, 58
 Theobromæ Oleum, 43, 73
 Theriaca, 83, 112
 Thns Americanum, 50, 81
 Thymelaceæ, 79
 Thymol, 88
 Tinctures, 5, 157
 Tobacco, 26, 79
 Toln, 51, 74
 Tops, 18, 19
 Tragacantha, 52, 75
 Treacle, 83, 112
 Triticum, 39, 83
 Trochisci, 8
 Turpentine, 50, 81
 " Oil of, 47, 81
 Twigs, 18
 Ulmaceæ, 81
 Ulmi Cortex, 24, 81
 Umbelliferæ, 76
 Unguenta, 9
 Unguentum Simplex, 9, 13
 Uvæ, 35, 73
 Uvæ Ursæ Folia, 26, 78
 Valeriana, 30, 77
 Valerianaceæ, 77
 Valerianic Acid, 30, 120
 Valerol, 30
 Vapores, 8
 Vegetable Kingdom, officinal drugs
 furnished by the, 16

- | | |
|---|--------------------------|
| Vegetable Kingdom, non-official drugs
furnished by the, 84 | Weights and Measures, 11 |
| Veratri Viridis Radix, 30, 82 | Wheat Flour, 39, 83 |
| Veratria, 62, 69, 82 | Wines, 6, 157 |
| Veratroidea, 30 | Wood-ashes, 114 |
| Vina, 6, 157 | Wood Tar, 105 |
| Vinegar, 100 | Woods, 22 |
| Vinegars, 6 | Woody Nightshade, 18, 79 |
| Vinum Aurantii, 104 | Worm Seed, 21, 77 |
| Vinum Xericum, 104 | Wurrus, 37, 80 |
| Viridia, 30 | |
| Vitaceæ, 73 | Yeast, 17, 83 |
| Volatile Oils, 44, 87 | |
| | Zinc Compounds, 134 |
| Water, 92 | Zingiber, 31, 82 |
| Waters, 6 | Zingiberaceæ, 82 |
| Wax, 12 | Zygophyllaceæ, 74 |



J. & A. CHURCHILL'S
MEDICAL CLASS BOOKS.

ANATOMY.

BRAUNE.—An Atlas of Topographical Anatomy, after Plane Sections of Frozen Bodies. By WILHELM BRAUNE, Professor of Anatomy in the University of Leipzig. Translated by EDWARD BELLAMY, F.R.C.S., and Member of the Board of Examiners; Surgeon to Charing Cross Hospital, and Lecturer on Anatomy in its School. With 34 Photo-lithographic Plates and 46 Woodcuts. Large Imp. 8vo, 40s.

FLOWER.—Diagrams of the Nerves of the Human Body, exhibiting their Origin, Divisions, and Connexions, with their Distribution to the various Regions of the Cutaneous Surface, and to all the Muscles. By WILLIAM H. FLOWER, F.R.C.S., F.R.S., Hunterian Professor of Comparative Anatomy, and Conservator of the Museum, Royal College of Surgeons. Third Edition, containing 6 Plates. Royal 4to, 12s.

GODLEE.—An Atlas of Human Anatomy: illustrating most of the ordinary Dissections and many not usually practised by the Student. By RICKMAN J. GODLEE, M.S., F.R.C.S., Assistant-Surgeon to University College Hospital, and Senior Demonstrator of Anatomy in University College. With 48 Imp. 4to Coloured Plates, containing 112 Figures, and a Volume of Explanatory Text, with many Engravings. 8vo, £4 14s. 6d.

HEATH.—Practical Anatomy: a Manual of Dissections. By CHRISTOPHER HEATH, F.R.C.S., Holme Professor of Clinical Surgery in University College and Surgeon to the Hospital. Fifth Edition. With 24 Coloured Plates and 269 Engravings. Crown 8vo, 15s.

ANATOMY—*continued.*

HOLDEN.—A Manual of the Dissection of the Human Body. By LUTHER HOLDEN, F.R.C.S., Consulting-Surgeon to St. Bartholomew's Hospital. Fourth Edition. Revised by the Author and JOHN LANGTON, F.R.C.S., Surgeon and Lecturer on Anatomy at St. Bartholomew's Hospital. With Engravings. 8vo, 16s.

By the same Author.

Human Osteology: comprising a Description of the Bones, with Delineations of the Attachments of the Muscles, the General and Microscopical Structure of Bone and its Development. Sixth Edition, revised by the Author and JAMES SHUTER, F.R.C.S., Assistant-Surgeon to St. Bartholomew's Hospital. With 61 Lithographic Plates and 89 Engravings. Royal 8vo, 16s.

ALSO,

Landmarks, Medical and Surgical. Third Edition. 8vo, 3s. 6d.

MORRIS.—The Anatomy of the Joints of Man.

By HENRY MORRIS, M.A., F.R.C.S., Surgeon to, and Lecturer on Anatomy and Practical Surgery at, the Middlesex Hospital. With 44 Plates (19 Coloured) and Engravings. 8vo, 16s.

The Anatomical Remembrancer; or, Complete Pocket Anatomist. Eighth Edition. 32mo, 3s. 6d.

WAGSTAFFE.—The Student's Guide to Human

Osteology. By WM. WARWICK WAGSTAFFE, F.R.C.S., late Assistant-Surgeon to, and Lecturer on Anatomy at, St. Thomas's Hospital. With 23 Plates and 66 Engravings. Fcap. 8vo, 10s. 6d.

WILSON — BUCHANAN — CLARK.—Wilson's

Anatomist's Vade-Mecum: a System of Human Anatomy. Tenth Edition, by GEORGE BUCHANAN, Professor of Clinical Surgery in the University of Glasgow, and HENRY E. CLARK, M.R.C.S., Lecturer on Anatomy in the Glasgow Royal Infirmary School of Medicine. With 450 Engravings, including 26 Coloured Plates. Crown 8vo, 18s.

11, NEW BURLINGTON STREET.

BOTANY.

BENTLEY.—A Manual of Botany. By Robert

BENTLEY, F.L.S., M.R.C.S., Professor of Botany in King's College and to the Pharmaceutical Society. With 1185 Engravings. Fourth Edition. Crown 8vo, 15s.

BENTLEY AND TRIMEN.—Medicinal Plants:

being descriptions, with original Figures, of the Principal Plants employed in Medicine, and an account of their Properties and Uses. By ROBERT BENTLEY, F.L.S., and HENRY TRIMEN, M.B., F.L.S. In 4 Vols., large 8vo, with 306 Coloured Plates, bound in half morocco, gilt edges, £11 11s.

CHEMISTRY.

BERNAYS.—Notes for Students in Chemistry;

being a Syllabus of Chemistry compiled mainly from the Manuals of Fownes-Watts, Miller, Wurz, and Schorlemmer. By ALBERT J. BERNAYS, Ph.D., Professor of Chemistry at St. Thomas's Hospital. Sixth Edition. Fcap. 8vo, 3s. 6d.

By the same Author.

Skeleton Notes on Analytical Chemistry,

for Students in Medicine. Fcap. 8vo, 2s. 6d.

BLOXAM.—Chemistry, Inorganic and Organic;

with Experiments. By CHARLES L. BLOXAM, Professor of Chemistry in King's College. Fifth Edition. With nearly 300 Engravings. 8vo, 16s.

By the same Author.

Laboratory Teaching; or, Progressive

Exercises in Practical Chemistry. Fourth Edition. With 83 Engravings. Crown 8vo, 5s. 6d.

CHEMISTRY—*continued.*

BOWMAN AND BLOXAM.—Practical Chemistry, including Analysis. By JOHN E. BOWMAN, formerly Professor of Practical Chemistry in King's College, and CHARLES L. BLOXAM, Professor of Chemistry in King's College. With 98 Engravings. Seventh Edition. Fcap. 8vo, 6s. 6d.

BROWN.—Practical Chemistry: Analytical Tables and Exercises for Students. By J. CAMPBELL BROWN, D.Sc. Lond., Professor of Chemistry in University College, Liverpool. Second Edition. 8vo, 2s. 6d.

CLOWES.—Practical Chemistry and Qualitative Inorganic Analysis. An Elementary Treatise, specially adapted for use in the Laboratories of Schools and Colleges, and by Beginners. By FRANK CLOWES, D.Sc., Professor of Chemistry in University College, Nottingham. Third Edition. With 47 Engravings. Post 8vo, 7s. 6d.

FOWNES AND WATTS.—Physical and Inorganic Chemistry. Twelfth Edition. By GEORGE FOWNES, F.R.S., and HENRY WATTS, B.A., F.R.S. With 154 Engravings, and Coloured Plate of Spectra. Crown 8vo, 8s. 6d.

By the same Authors.

Chemistry of Carbon - Compounds, or
Organic Chemistry. Twelfth Edition. With Engravings. Crown 8vo, 10s.

LUFF.—An Introduction to the Study of Chemistry. Specially designed for Medical and Pharmaceutical Students. By A. P. LUFF, F.I.C., F.C.S., Lecturer on Chemistry in the Central School of Chemistry and Pharmacy. Crown 8vo, 2s. 6d.

TIDY.—A Handbook of Modern Chemistry, Inorganic and Organic. By C. MEYMOTT TIDY, M.B., Professor of Chemistry and Medical Jurisprudence at the London Hospital. 8vo, 16s.

VACHER.—A Primer of Chemistry, including Analysis. By ARTHUR VACHER. 18mo, 1s.

11, NEW BURLINGTON STREET.

CHEMISTRY—*continued.*

VALENTIN.—Introduction to Inorganic Chemistry. By WILLIAM G. VALENTIN, F.C.S. Third Edition. With 82 Engravings. Svo, 6s. 6d.

By the same Author.

A Course of Qualitative Chemical Analysis.

Fifth Edition. By W. R. HODGKINSON, Ph.D. (Würzburg), Demonstrator of Practical Chemistry in the Science Training Schools. With Engravings. Svo, 7s. 6d.

ALSO,

Chemical Tables for the Lecture-room and Laboratory. In Five large Sheets, 5s. 6d.

CHILDREN, DISEASES OF.

DAY.—A Treatise on the Diseases of Children.

For Practitioners and Students. By WILLIAM H. DAY, M.D., Physician to the Samaritan Hospital for Women and Children. Crown Svo, 12s. 6d.

ELLIS.—A Practical Manual of the Diseases

of Children. By EDWARD ELLIS, M.D., late Senior Physician to the Victoria Hospital for Sick Children. With a Formulary. Fourth Edition. Crown Svo, 10s.

SMITH.—Clinical Studies of Disease in

Children. By EUSTACE SMITH, M.D., F.R.C.P., Physician to H.M. the King of the Belgians, and to the East London Hospital for Children. Second Edition. Post Svo. *[In preparation.]*

By the same Author.

On the Wasting Diseases of Infants and Children. Third Edition. Post Svo, 8s. 6d.

STEINER.—Compendium of Children's Dis-

eases; a Handbook for Practitioners and Students. By JOHANN STEINER, M.D. Translated by LAWSON TAIT, F.R.C.S., Surgeon to the Birmingham Hospital for Women, &c. Svo, 12s. 6d.

DENTISTRY.

SEWILL.—The Student's Guide to Dental

Anatomy and Surgery. By HENRY E. SEWILL, M.R.C.S., L.D.S., late Dental Surgeon to the West London Hospital. Second Edition. With 78 Engravings. Fcap. 8vo, 5s. 6d.

STOCKEN.—Elements of Dental Materia Medica

and Therapeutics, with Pharmacopœia. By JAMES STOCKEN, L.D.S.R.C.S., late Lecturer on Dental Materia Medica and Therapeutics and Dental Surgeon to the National Dental Hospital, assisted by THOMAS GADDES, L.D.S. Eng. and Edin. Third Edition. Fcap. 8vo, 7s. 6d.

TAFT.—A Practical Treatise on Operative

Dentistry. By JONATHAN TAFT, D.D.S., Professor of Operative Surgery in the Ohio College of Dental Surgery. Third Edition. With 134 Engravings. 8vo, 18s.

TOMES (C. S.).—Manual of Dental Anatomy,

Human and Comparative. By CHARLES S. TOMES, M.A., F.R.S. Second Edition. With 191 Engravings. Crown 8vo, 12s. 6d.

TOMES (J. and C. S.).—A Manual of Dental

Surgery. By JOHN TOMES, M.R.C.S., F.R.S., and CHARLES S. TOMES, M.A., M.R.C.S., F.R.S.; Lecturer on Anatomy and Physiology at the Dental Hospital of London. Second Edition. With 262 Engravings. Fcap. 8vo, 14s.

EAR, DISEASES OF.

BURNETT.—The Ear: its Anatomy, Physio-

logy, and Diseases. A Practical Treatise for the Use of Medical Students and Practitioners. By CHARLES H. BURNETT, M.D., Aural Surgeon to the Presbyterian Hospital, Philadelphia. With 87 Engravings. 8vo, 18s.

DALBY.—On Diseases and Injuries of the Ear.

By WILLIAM B. DALBY, F.R.C.S., Aural Surgeon to, and Lecturer on Aural Surgery at, St. George's Hospital. Second Edition. With Engravings. Fcap. 8vo, 6s. 6d.

EAR, DISEASES OF—continued.

JONES.—A Practical Treatise on Aural Surgery. By H. MACNAUGHTON JONES, M.D., Professor of the Queen's University in Ireland, Surgeon to the Cork Ophthalmic and Aural Hospital. Second Edition. With 63 Engravings. Crown 8vo, 8s. 6d.

By the same Author.

Atlas of the Diseases of the Membrana Tympani. In Coloured Plates, containing 59 Figures. With Explanatory Text. Crown 4to, 21s.

FORENSIC MEDICINE.

OGSTON.—Lectures on Medical Jurisprudence.

By FRANCIS OGSTON, M.D., Professor of Medical Jurisprudence and Medical Logic in the University of Aberdeen. Edited by FRANCIS OGSTON, Jun., M.D., Assistant to the Professor of Medical Jurisprudence and Lecturer on Practical Toxicology in the University of Aberdeen. With 12 Plates. 8vo, 18s.

TAYLOR.—The Principles and Practice of

Medical Jurisprudence. By ALFRED S. TAYLOR, M.D., F.R.S. Third Edition, revised by THOMAS STEVENSON, M.D., F.R.C.P., Lecturer on Chemistry at Guy's Hospital, &c. With 188 Engravings. 2 Vols. 8vo, 31s. 6d.

By the same Author.

A Manual of Medical Jurisprudence.

Tenth Edition. With 55 Engravings. Crown 8vo, 14s.

ALSO,

On Poisons, in relation to Medical Jurisprudence and Medicine. Third Edition. With 104 Engravings. Crown 8vo, 16s.

TIDY AND WOODMAN.—A Handy-Book of

Forensic Medicine and Toxicology. By C. MEYMOTT TIDY, M.B.; and W. BATHURST WOODMAN, M.D., F.R.C.P. With 8 Lithographic Plates and 116 Wood Engravings. 8vo, 31s. 6d.

HYGIENE.

PARKES.—A Manual of Practical Hygiene.

By EDMUND A. PARKES, M.D., F.R.S. Sixth Edition by F. DE CHAUMONT, M.D., F.R.S., Professor of Military Hygiene in the Army Medical School. With 9 Plates and 103 Engravings. 8vo, 18s.

WILSON.—A Handbook of Hygiene and Sanitary Science.

By GEORGE WILSON, M.A., M.D., Medical Officer of Health for Mid Warwickshire. Fourth Edition. With Engravings. Crown 8vo, 10s. 6d.

MATERIA MEDICA AND THERAPEUTICS.

BINZ AND SPARKS.—The Elements of Therapeutics: a Clinical Guide to the Action of Medicines.

By C. BINZ, M.D., Professor of Pharmacology in the University of Bonn. Translated and Edited with Additions, in conformity with the British and American Pharmacopœias, by EDWARD I. SPARKS, M.A., M.B., F.R.C.P. Lond. Crown 8vo, 8s. 6d.

OWEN.—Tables of Materia Medica; comprising the Contents, Doses, Proportional Composition, and Methods

of Manufacture of Pharmacopœial Preparations. By ISAMBARD OWEN, M.B., M.R.C.P., Lecturer on Materia Medica at St. George's Hospital. Fifth Edition. Crown 8vo, 2s. 6d.

ROYLE AND HARLEY.—A Manual of Materia

Medica and Therapeutics. By J. FORBES ROYLE, M.D., F.R.S., and JOHN HARLEY, M.D., F.R.C.P., Physician to, and Joint Lecturer on Clinical Medicine at, St. Thomas's Hospital. Sixth Edition. With 139 Engravings. Crown 8vo, 15s.

THOROWGOOD.—The Student's Guide to

Materia Medica and Therapeutics. By JOHN C. THOROWGOOD, M.D., F.R.C.P., Lecturer on Materia Medica at the Middlesex Hospital. Second Edition. With Engravings. Fcap. 8vo, 7s.

WARING.—A Manual of Practical Therapeutics.

By EDWARD J. WARING, C.B., M.D., F.R.C.P. Third Edition. Fcap. 8vo, 12s. 6d.

MEDICINE.

BARCLAY.—A Manual of Medical Diagnosis.

By A. WHYTE BARCLAY, M.D., F.R.C.P., late Physician to, and Lecturer on Medicine at, St. George's Hospital. Third Edition. Fcap. 8vo, 10s. 6d.

CHARTERIS.—The Student's Guide to the

Practice of Medicine. By MATTHEW CHARTERIS, M.D., Professor of Materia Medica, University of Glasgow; Physician to the Royal Infirmary. With Engravings on Copper and Wood. Third Edition. Fcap. 8vo, 7s.

FENWICK.—The Student's Guide to Medical

Diagnosis. By SAMUEL FENWICK, M.D., F.R.C.P., Physician to the London Hospital. Fifth Edition. With 111 Engravings. Fcap. 8vo, 7s.

By the same Author.

The Student's Outlines of Medical Treatment. Second Edition. Fcap. 8vo, 7s.

FLINT.—Clinical Medicine: a Systematic Treatise

on the Diagnosis and Treatment of Disease. By AUSTIN FLINT, M.D., Professor of the Principles and Practice of Medicine, &c., in Bellevue Hospital Medical College. 8vo, 20s.

HALL.—Synopsis of the Diseases of the Larynx,

Lungs, and Heart: comprising Dr. Edwards' Tables on the Examination of the Chest. With Alterations and Additions. By F. DE HAVILLAND HALL, M.D., F.R.C.P., Assistant-Physician to the Westminster Hospital. Royal 8vo, 2s. 6d.

SANSOM.—Manual of the Physical Diagnosis

of Diseases of the Heart, including the use of the Sphygmograph and Cardiograph. By A. E. SANSOM, M.D., F.R.C.P., Assistant-Physician to the London Hospital. Third Edition. With 47 Woodcuts. Fcap. 8vo, 7s. 6d.

WARNER.—Student's Guide to Medical Case-

Taking. By FRANCIS WARNER, M.D., Assistant-Physician to the London Hospital. Fcap. 8vo, 5s.

WEST.—How to Examine the Chest: being a

Practical Guide for the Use of Students. By SAMUEL WEST, M.D., M.R.C.P., Physician to the City of London Hospital for Diseases of the Chest, &c. With 42 Engravings. Fcap. 8vo, 5s.

MEDICINE—*continued.*

WHITTAKER.—Students' Primer on the Urine.

By J. TRAVIS WHITTAKER, M.D., Clinical Demonstrator at the Royal Infirmary, Glasgow. With Illustrations, and 16 Plates etched on Copper. Post 8vo, 4s. 6d.

MIDWIFERY.

BARNES.—Lectures on Obstetric Operations,

including the Treatment of Hæmorrhage, and forming a Guide to the Management of Difficult Labour. By ROBERT BARNES, M.D., F.R.C.P., Obstetric Physician to, and Lecturer on Diseases of Women, &c., at, St. George's Hospital. Third Edition. With 124 Engravings. 8vo, 18s.

CLAY.—The Complete Handbook of Obstetric

Surgery; or, Short Rules of Practice in every Emergency, from the Simplest to the most formidable Operations connected with the Science of Obstetrics. By CHARLES CLAY, M.D., late Senior Surgeon to, and Lecturer on Midwifery at, St. Mary's Hospital, Manchester. Third Edition. With 91 Engravings. Fcap. 8vo, 6s. 6d.

RAMSBOTHAM.—The Principles and Practice

of Obstetric Medicine and Surgery. By FRANCIS H. RAMSBOTHAM, M.D., formerly Obstetric Physician to the London Hospital. Fifth Edition. With 120 Plates, forming one thick handsome volume. 8vo, 22s.

REYNOLDS.—Notes on Midwifery: specially

designed to assist the Student in preparing for Examination. By J. J. REYNOLDS, M.R.C.S. Fcap. 8vo, 4s.

ROBERTS.—The Student's Guide to the Practice

of Midwifery. By D. LLOYD ROBERTS, M.D., F.R.C.P., Physician to St. Mary's Hospital, Manchester. Second Edition. With 111 Engravings. Fcap. 8vo, 7s.

SCHROEDER.—A Manual of Midwifery; includ-

ing the Pathology of Pregnancy and the Puerperal State. By KARL SCHROEDER, M.D., Professor of Midwifery in the University of Erlangen. Translated by CHARLES H. CARTER, M.D. With Engravings. 8vo, 12s. 6d.

11, NEW BURLINGTON STREET.

MIDWIFERY—*continued.*

SWAYNE.—Obstetric Aphorisms for the Use of Students commencing Midwifery Practice. By JOSEPH G. SWAYNE, M.D., Lecturer on Midwifery at the Bristol School of Medicine. Seventh Edition. With Engravings. Fcap. 8vo, 3s. 6d.

MICROSCOPY.

CARPENTER.—The Microscope and its Revelations. By WILLIAM B. CARPENTER, C.B., M.D., F.R.S. Sixth Edition. With 26 Plates, a Coloured Frontispiece, and more than 500 Engravings. Crown 8vo, 16s.

MARSH.—Microscopical Section-Cutting: a Practical Guide to the Preparation and Mounting of Sections for the Microscope, special prominence being given to the subject of Animal Sections. By Dr. SYLVESTER MARSH. Second Edition. With 17 Engravings. Fcap. 8vo, 3s. 6d.

MARTIN.—A Manual of Microscopic Mounting By JOHN H. MARTIN, Member of the Society of Public Analysts, &c. Second Edition. With several Plates and 144 Engravings. 8vo, 7s. 6d.

OPHTHALMOLOGY.

DAGUENET.—A Manual of Ophthalmoscopy for the Use of Students. By Dr. DAGUENET. Translated by C. S. JEAFFRESON, Surgeon to the Newcastle-on-Tyne Eye Infirmary. With Engravings. Fcap. 8vo, 5s.

HIGGENS.—Hints on Ophthalmic Out-Patient Practice. By CHARLES HIGGENS, F.R.C.S., Ophthalmic Surgeon to, and Lecturer on Ophthalmology at, Guy's Hospital. Second Edition. Fcap. 8vo, 3s.

JONES.—A Manual of the Principles and Practice of Ophthalmic Medicine and Surgery. By T. WHARTON JONES, F.R.C.S., F.R.S., late Ophthalmic Surgeon and Professor of Ophthalmology to University College Hospital. Third Edition. With 9 Coloured Plates and 173 Engravings. Fcap. 8vo, 12s. 6d.

OPHTHALMOLOGY—*continued.*

MACNAMARA.—A Manual of the Diseases of the Eye. By CHARLES MACNAMARA, F.R.C.S., Surgeon to, and Lecturer on Surgery at, Westminster Hospital. Fourth Edition. With 4 Coloured Plates and 66 Engravings. Fcap. 8vo, 10s. 6d.

NETTLESHIP.—The Student's Guide to Diseases of the Eye. By EDWARD NETTLESHIP, F.R.C.S., Ophthalmic Surgeon to, and Lecturer on Ophthalmic Surgery at, St. Thomas's Hospital. Second Edition. With 137 Engravings, and a Set of Coloured Papers illustrating Colour-blindness. Fcap. 8vo, 7s. 6d.

WOLFE.—On Diseases and Injuries of the Eye : a Course of Systematic and Clinical Lectures to Students and Medical Practitioners. By J. R. WOLFE, M.D., F.R.C.S.E., Senior Surgeon to the Glasgow Ophthalmic Institution, Lecturer on Ophthalmic Medicine and Surgery in Anderson's College. With 10 Coloured Plates, and 120 Wood Engravings, 8vo.

PATHOLOGY.

JONES AND SIEVEKING.—A Manual of Pathological Anatomy. By C. HANDFIELD JONES, M.B., F.R.S., and EDWARD H. SIEVEKING, M.D., F.R.C.P. Second Edition. Edited, with considerable enlargement, by J. F. PAYNE, M.B., Assistant-Physician and Lecturer on General Pathology at St. Thomas's Hospital. With 195 Engravings. Crown 8vo, 16s.

LANCEREAUX.—Atlas of Pathological Anatomy. By Dr. LANCEREAUX. Translated by W. S. GREENFIELD, M.D., Professor of Pathology in the University of Edinburgh. With 70 Coloured Plates. Imperial 8vo, £5 5s.

VIRCHOW.—Post-Mortem Examinations: a Description and Explanation of the Method of Performing them, with especial reference to Medico-Legal Practice. By Professor RUDOLPH VIRCHOW, Berlin Charité Hospital. Translated by Dr. T. B. SMITH. Second Edition, with 4 Plates. Fcap. 8vo, 3s. 6d.

WILKS AND MOXON.—Lectures on Pathological Anatomy. By SAMUEL WILKS, M.D., F.R.S., Physician to, and late Lecturer on Medicine at, Guy's Hospital; and WALTER MOXON, M.D., F.R.C.P., Physician to, and Lecturer on the Practice of Medicine at, Guy's Hospital. Second Edition. With 7 Steel Plates. 8vo, 18s.

PSYCHOLOGY.

BUCKNILL AND TUKE.—*A Manual of Psychological Medicine: containing the Lunacy Laws, Nosology, Ætiology, Statistics, Description, Diagnosis, Pathology, and Treatment of Insanity, with an Appendix of Cases.* By JOHN C. BUCKNILL, M.D., F.R.S., and D. HACK TUKE, M.D., F.R.C.P. Fourth Edition, with 12 Plates (30 Figures). Svo, 25s.

PHYSIOLOGY.

CARPENTER.—*Principles of Human Physiology.* By WILLIAM B. CARPENTER, C.B., M.D., F.R.S. Ninth Edition. Edited by Henry Power, M.B., F.R.C.S. With 3 Steel Plates and 377 Wood Engravings. Svo, 31s. 6d.

DALTON.—*A Treatise on Human Physiology: designed for the use of Students and Practitioners of Medicine.* By JOHN C. DALTON, M.D., Professor of Physiology and Hygiene in the College of Physicians and Surgeons, New York. Seventh Edition. With 252 Engravings. Royal Svo, 20s.

FREY.—*The Histology and Histo-Chemistry of Man. A Treatise on the Elements of Composition and Structure of the Human Body.* By HEINRICH FREY, Professor of Medicine in Zurich. Translated by ARTHUR E. BARKER, Assistant-Surgeon to the University College Hospital. With 608 Engravings. Svo, 21s.

SANDERSON.—*Handbook for the Physiological Laboratory: containing an Exposition of the fundamental facts of the Science, with explicit Directions for their demonstration.* By J. BURDON SANDERSON, M.D., F.R.S., Jodrell Professor of Physiology in University College; E. KLEIN, M.D., F.R.S., Assistant-Professor in the Brown Institution; MICHAEL FOSTER, M.D., F.R.S., Prælector of Physiology at Trinity College, Cambridge; and T. LAUDER BRUNTON, M.D., F.R.S., Lecturer on Materia Medica at St. Bartholomew's Hospital Medical College. 2 Vols., with 123 Plates. Svo, 24s.

SURGERY.

BRYANT.—*A Manual for the Practice of Surgery.* By THOMAS BRYANT, F.R.C.S., Surgeon to, and Lecturer on Surgery at, Guy's Hospital. Third Edition. With 672 Engravings (nearly all original, many being coloured). 2 vols. Crown Svo, 28s.

SURGERY—*continued.*

BELLAMY.—The Student's Guide to Surgical

Anatomy; a Description of the more important Surgical Regions of the Human Body, and an Introduction to Operative Surgery. By EDWARD BELLAMY, F.R.C.S., and Member of the Board of Examiners; Surgeon to, and Lecturer on Anatomy at, Charing Cross Hospital. Second Edition. With 76 Engravings. Fcap. 8vo, 7s.

CLARK AND WAGSTAFFE. — Outlines of

Surgery and Surgical Pathology. By F. LE GROS CLARK, F.R.C.S., F.R.S., Consulting Surgeon to St. Thomas's Hospital. Second Edition. Revised and expanded by the Author, assisted by W. W. WAGSTAFFE, F.R.C.S., Assistant-Surgeon to St. Thomas's Hospital. Svo, 10s. 6d.

DRUITT. — The Surgeon's Vade-Mecum; a

Manual of Modern Surgery. By ROBERT DRUITT, F.R.C.S. Eleventh Edition. With 369 Engravings. Fcap. 8vo, 14s.

FERGUSON.—A System of Practical Surgery.

By Sir WILLIAM FERGUSON, Bart., F.R.C.S., F.R.S., late Surgeon and Professor of Clinical Surgery to King's College Hospital. With 463 Engravings. Fifth Edition. Svo, 21s.

HEATH.—A Manual of Minor Surgery and

Bandaging, for the use of House-Surgeons, Dressers, and Junior Practitioners. By CHRISTOPHER HEATH, F.R.C.S., Holme Professor of Clinical Surgery in University College and Surgeon to the Hospital. Sixth Edition. With 115 Engravings. Fcap. 8vo. 5s. 6d.

By the same Author.

A Course of Operative Surgery: with

Twenty Plates drawn from Nature by M. LÉVEILLÉ, and Coloured by hand under his direction. Large Svo, 40s.

ALSO,

The Student's Guide to Surgical Diag-

nosis. Second Edition. Fcap. 8vo, 6s. 6d.

MAUNDER.—Operative Surgery. By Charles

F. MAUNDER, F.R.C.S., late Surgeon to, and Lecturer on Surgery at, the London Hospital. Second Edition. With 164 Engravings. Post 8vo, 6s.

SURGERY—continued.

PIRRIE.—The Principles and Practice of Surgery. By WILLIAM PIRRIE, F.R.S.E., late Professor of Surgery in the University of Aberdeen. Third Edition. With 490 Engravings. 8vo, 28s.

SOUTHAM.—Regional Surgery: including Surgical Diagnosis. A Manual for the use of Students. Part I., the Head and Neck. By FREDERICK A. SOUTHAM, M.A., M.B. Oxon, F.R.C.S., Assistant-Surgeon to the Royal Infirmary, and Assistant-Lecturer on Surgery in the Owen's College School of Medicine, Manchester. Crown 8vo, 6s. 6d.

TERMINOLOGY.

DUNGLISON.—Medical Lexicon: a Dictionary of Medical Science, containing a concise Explanation of its various Subjects and Terms, with Accentuation, Etymology, Synonymes, &c. By ROBLEY DUNGLISON, M.D. New Edition, thoroughly revised by RICHARD J. DUNGLISON, M.D. Royal 8vo, 28s.

MAYNE.—A Medical Vocabulary: being an Explanation of all Terms and Phrases used in the various Departments of Medical Science and Practice, giving their Derivation, Meaning, Application, and Pronunciation. By ROBERT G. MAYNE, M.D., LL.D., and JOHN MAYNE, M.D., L.R.C.S.E. Fifth Edition. Fcap. 8vo, 10s. 6d.

WOMEN, DISEASES OF.

BARNES.—A Clinical History of the Medical and Surgical Diseases of Women. By ROBERT BARNES, M.D., F.R.C.P., Obstetric Physician to, and Lecturer on Diseases of Women, &c., at, St. George's Hospital. Second Edition. With 181 Engravings. 8vo, 28s.

COURTY.—Practical Treatise on Diseases of the Uterus, Ovaries, and Fallopian Tubes. By Professor COURTY, Montpellier. Translated from the Third Edition by his Pupil, AGNES M'LAREN, M.D., M.K.Q.C.P. With Preface by Dr. MATTHEWS DUNCAN. With 424 Engravings. 8vo, 24s.

DUNCAN.—Clinical Lectures on the Diseases of Women. By J. MATTHEWS DUNCAN, M.D., F.R.C.P., Obstetric Physician to St. Bartholomew's Hospital. Second Edition, with Appendices. 8vo, 14s.

WOMEN, DISEASES OF—*continued.*

EMMET.—The Principles and Practice of Gynæcology. By THOMAS ADDIS EMMET, M.D., Surgeon to the Woman's Hospital of the State of New York. With 130 Engravings. Royal 8vo, 24s.

GALABIN.—The Student's Guide to the Diseases of Women. By ALFRED L. GALABIN, M.D., F.R.C.P., Obstetric Physician and Joint Lecturer on Obstetric Medicine at Guy's Hospital. Second Edition. With 70 Engravings. Fcap. 8vo, 7s. 6d.

REYNOLDS.—Notes on Diseases of Women. Specially designed for Students preparing for Examination. By J. J. REYNOLDS, M.R.C.S. Fcap. 8vo, 2s. 6d.

SMITH.—Practical Gynæcology: a Handbook of the Diseases of Women. By HEYWOOD SMITH, M.D., Physician to the Hospital for Women and to the British Lying-in Hospital. With Engravings. Second Edition. Crown 8vo. [*In preparation.*]

WEST AND DUNCAN.—Lectures on the Diseases of Women. By CHARLES WEST, M.D., F.R.C.P. Fourth Edition. Revised and in part re-written by the Author, with numerous additions, by J. MATTHEWS DUNCAN, M.D., F.R.C.P., Obstetric Physician to St. Bartholomew's Hospital. 8vo, 16s.

ZOOLOGY.

CHAUVEAU AND FLEMING.—The Comparative Anatomy of the Domesticated Animals. By A. CHAUVEAU, Professor at the Lyons Veterinary School; and GEORGE FLEMING, Veterinary Surgeon, Royal Engineers. With 450 Engravings. 8vo, 31s. 6d.

HUXLEY.—Manual of the Anatomy of Invertebrated Animals. By THOMAS H. HUXLEY, LL.D., F.R.S. With 156 Engravings. Fcap. 8vo, 16s.

By the same Author.

Manual of the Anatomy of Vertebrated Animals. With 110 Engravings. Post 8vo, 12s.

WILSON.—The Student's Guide to Zoology: a Manual of the Principles of Zoological Science. By ANDREW WILSON, Lecturer on Natural History, Edinburgh. With Engravings. Fcap. 8vo, 6s. 6d.

11, NEW BURLINGTON STREET.



